

X(4660) $I^G(J^{PC}) = ?^?(1^{--})$

Seen in radiative return from $e^+ e^-$ collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review under the $X(3872)$ particle listings. (See the index for the page number.)

X(4660) MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-----------------------|----------|---|
| 4643± 9 OUR AVERAGE | | | | Error includes scale factor of 1.2. |
| 4652±10±11 | 279 | ¹ WANG | 15A BELL | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |
| 4669±21± 3 | 37 | ² LEES | 14F BABR | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |
| 4634 ^{+ 8 + 5} _{- 7 - 8} | 142 | ³ PAKHLOVA | 08B BELL | $e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|----|------------------|----------|---|
| 4661 ^{+ 9 ± 6} _{- 8 ± 6} | 44 | ⁴ LIU | 08H RVUE | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |
| 4664±11± 5 | 44 | WANG | 07D BELL | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |

¹ From a two-resonance fit. Supersedes WANG 07D.

² From a two-resonance fit.

³ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

⁴ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

X(4660) WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-----------------------|----------|---|
| 72±11 OUR AVERAGE | | | | |
| 68±11± 5 | 279 | ¹ WANG | 15A BELL | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |
| 104±48±10 | 37 | ² LEES | 14F BABR | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |
| 92 ^{+40 +10} _{-24 -21} | 142 | ³ PAKHLOVA | 08B BELL | $e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|----|------------------|----------|---|
| 42 ^{+17 ± 6} _{-12 ± 6} | 44 | ⁴ LIU | 08H RVUE | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |
| 48±15± 3 | 44 | WANG | 07D BELL | $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$ |

¹ From a two-resonance fit. Supersedes WANG 07D.

² From a two-resonance fit.

³ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

⁴ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

X(4660) DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|-----------------------------------|--------------------------------|
| $\Gamma_1 e^+ e^-$ | |
| $\Gamma_2 \psi(2S)\pi^+\pi^-$ | seen |
| $\Gamma_3 J/\psi\eta$ | |
| $\Gamma_4 D^0 D^{*-}\pi^+$ | |
| $\Gamma_5 \chi_{c1}\gamma$ | |
| $\Gamma_6 \chi_{c2}\gamma$ | |
| $\Gamma_7 \Lambda_c^+\Lambda_c^-$ | |

X(4660) $\Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_1/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------------|----------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.0 \pm 0.3 \pm 0.2 | 279 | ¹ WANG | 15A BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 8.1 \pm 1.1 \pm 1.0 | 279 | ² WANG | 15A BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 2.7 \pm 1.3 \pm 0.5 | 37 | ³ LEES | 14F BABR | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 7.5 \pm 1.7 \pm 0.7 | 37 | ⁴ LEES | 14F BABR | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 2.2 $^{+0.7}_{-0.6}$ | 44 | ⁵ LIU | 08H RVUE | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 5.9 \pm 1.6 | 44 | ⁶ LIU | 08H RVUE | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 3.0 \pm 0.9 \pm 0.3 | 44 | ³ WANG | 07D BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 7.6 \pm 1.8 \pm 0.8 | 44 | ⁴ WANG | 07D BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |

¹ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

² Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

³ Solution I of two equivalent solutions in a fit using two interfering resonances.

⁴ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁵ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

⁶ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_1/\Gamma$$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------|----------|---------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.94 | 90 | WANG | 13B BELL | $e^+e^- \rightarrow J/\psi\eta\gamma$ |

$$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_5\Gamma_1/\Gamma$$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------|-----|------------------|---------|--|
| <0.45 | 90 | ¹ HAN | 15 BELL | 10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$ |

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

| $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | $\Gamma_6\Gamma_1/\Gamma$ | | | |
|--|---------------------------|-------------|------|---|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
| <2.1 | 90 | 1 HAN | 15 | BELL $10.58 e^+e^- \rightarrow \chi_{c2}\gamma$ |
| ¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$. | | | | |

X(4660) BRANCHING RATIOS

| $\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S) \pi^+ \pi^-)$ | Γ_4/Γ_2 | | | |
|---|---------------------|-------------|------|---------------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <10 | 90 | PAKHLOVA 09 | BELL | $e^+e^- \rightarrow D^0 D^{*-} \pi^+$ |

| $\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | $\Gamma_4/\Gamma \times \Gamma_1/\Gamma$ | | | |
|--|--|---------------|------|---------------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| $<0.37 \times 10^{-6}$ | 90 | 1 PAKHLOVA 09 | BELL | $e^+e^- \rightarrow D^0 D^{*-} \pi^+$ |

¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of X(4660).

| $\Gamma(\Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | $\Gamma_7/\Gamma \times \Gamma_1/\Gamma$ | | | |
|---|--|----------------|------|--|
| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
| $0.68^{+0.16+0.29}_{-0.15-0.30}$ | 142 | 1 PAKHLOVA 08B | BELL | $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ |

¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

X(4660) REFERENCES

| | | | | |
|----------|-----|----------------|-------------------------------|-----------------|
| HAN | 15 | PR D92 012011 | Y.L. Han <i>et al.</i> | (BELLE Collab.) |
| WANG | 15A | PR D91 112007 | X.L. Wang <i>et al.</i> | (BELLE Collab.) |
| LEES | 14F | PR D89 111103 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| WANG | 13B | PR D87 051101 | X.L. Wang <i>et al.</i> | (BELLE Collab.) |
| PAKHLOVA | 09 | PR D80 091101 | G. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| LIU | 08H | PR D78 014032 | Z.Q. Liu, X.S. Qin, C.Z. Yuan | |
| PAKHLOVA | 08B | PRL 101 172001 | C. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| AUBERT | 07S | PRL 98 212001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| WANG | 07D | PRL 99 142002 | X.L. Wang <i>et al.</i> | (BELLE Collab.) |