

$\chi_{b1}(3P)$ $I^G(J^{PC}) = 0^+(1^{++})$

Observed in the radiative decay to $\gamma(1S, 2S, 3S)$, therefore $C = +$.
 J needs confirmation.

 $\chi_{b1}(3P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10512.1 ± 2.1 ± 0.9	351	1 AAIJ	14BG LHCb	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10515.7 ± 2.2 ± 1.5	169	2 AAIJ	14BG LHCb	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$
10511.3 ± 1.7 ± 2.5	182	3 AAIJ	14BI LHCb	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$
10530 ± 5 ± 9		4 AAD	12A ATLAS	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$
10551 ± 14 ± 17		4 ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$

¹ The mass of the $\chi_{b1}(3P)$ state obtained by combining the results of AAIJ 14BG with that of AAIJ 14BI. The first uncertainty is experimental and the second attributable to the unknown mass splitting, assumed to be $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

² From $\chi_{b1}(3P) \rightarrow \gamma(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for ±30% variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

³ From $\chi_{b1}(3P) \rightarrow \gamma(3S)\gamma$ transition assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

⁴ The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.

 $\chi_{b1}(3P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma(1S)\gamma$	seen
$\Gamma_2 \quad \gamma(2S)\gamma$	seen
$\Gamma_3 \quad \gamma(3S)\gamma$	seen

 $\chi_{b1}(3P)$ BRANCHING RATIOS

$\Gamma(\gamma(1S)\gamma)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_1/Γ
seen	169	5 AAIJ	14BG LHCb	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen		AAD	12A ATLAS	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$
seen		ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma\mu^+\mu^-X$

⁵ From $\chi_{b1}(3P) \rightarrow \gamma(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for ±30% variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_2/Γ</u>
seen	169	⁶ AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen		AAD	12A ATLAS	$p p \rightarrow \gamma \mu^+ \mu^- X$	
⁶ From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.					

 $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_3/Γ</u>
seen	182	AAIJ	14BI LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$	

 $\chi_{b1}(3P)$ REFERENCES

AAIJ	14BG JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BI EPJ C74 3092	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	12A PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV	12Q PR D86 031103	V.M. Abazov <i>et al.</i>	(D0 Collab.)