

$\eta_b(1S)$ $I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.


 $\eta_b(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
9399.0± 2.3 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.			
9400.7± 1.7± 1.6	33.1k	TAMPONI	15	BELL $e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$	
9402.4± 1.5± 1.8	34k	¹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$	
9391.8± 6.6± 2.0	2.3k	² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$	
9394.2± 4.8± 2.0	13k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$	
9388.9± 3.1± 2.7	19k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
9393.2± 3.4± 2.3	10	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$	
9300 ±20 ±20		HEISTER	02D ALEP	181–209 $e^+ e^-$	

¹ With floating width. Not independent of the corresponding mass difference measurement.


² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.

³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.



$m\gamma(1S) - m_{\eta_b}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3±3.2 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
57.9±1.5±1.8	34k	⁴ MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^-$ + hadrons
68.5±6.6±2.0	2.3 ± 0.5k	⁵ BONVICINI	10 CLEO	$\gamma(3S) \rightarrow \gamma X$
66.1 ^{+4.8} _{-4.9} ±2.0	13 ± 5k	⁵ AUBERT	09AQ BABR	$\gamma(2S) \rightarrow \gamma X$
71.4 ^{+2.3} _{-3.1} ±2.7	19 ± 3k	⁵ AUBERT	08V BABR	$\gamma(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1±3.4±2.3	10 ⁺⁵ ₋₄	^{5,6} DOBBS	12	$\gamma(2S) \rightarrow \gamma$ hadrons

⁴ With floating width. Not independent of the corresponding mass measurement.⁵ Assuming $\Gamma_{\eta_b}(1S) = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.⁶ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration. γ ENERGY IN $\gamma(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
920.6^{+2.8}_{-3.2} OUR AVERAGE				
918.6±6.0±1.9	2.3 ± 0.5k	⁷ BONVICINI	10 CLEO	$\gamma(3S) \rightarrow \gamma X$
921.2 ^{+2.1} _{-2.8} ±2.4	19 ± 3k	⁷ AUBERT	08V BABR	$\gamma(3S) \rightarrow \gamma X$

⁷ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
609.3^{+4.6}_{-4.5}^{±1.9}	$13 \pm 5k$	⁸ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$

⁸ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

$\eta_b(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10⁺⁵₋₄ OUR AVERAGE				
8 ⁺⁶ ₋₅ ± 5	33.1k	⁹ TAMPONI	15	BELL $e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
10.8 ^{+4.0} _{-3.7} ^{+4.5} _{-2.0}	34k	⁹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$

⁹ With floating mass.

$\eta_b(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+ 3h^-$	not seen	
Γ_3 $2h^+ 2h^-$	not seen	
Γ_4 $4h^+ 4h^-$		
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+ \mu^-$	$< 9 \times 10^{-3}$	90%
Γ_7 $\tau^+ \tau^-$	$< 8\%$	90%

$\eta_b(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(3h^+ 3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_5/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<470	95	ABDALLAH	06	DLPH 161–209 $e^+ e^-$
<132	95	HEISTER	02D	ALEP 181–209 $e^+ e^-$

$\Gamma(2h^+ 2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3 \Gamma_5/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<190	95	ABDALLAH	06	DLPH 161–209 $e^+ e^-$
< 48	95	HEISTER	02D	ALEP 181–209 $e^+ e^-$

$\Gamma(4h^+4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_5/\Gamma$
<u>VALUE</u> (eV)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<660	95	ABDALLAH 06	DLPH	161–209 $e^+ e^-$	

 $\eta_b(1S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	34k	MIZUK	12	BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$					Γ_6/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<9 \times 10^{-3}$	90	¹⁰ AUBERT	09Z	BABR	$e^+ e^- \rightarrow \gamma(2S, 3S) \rightarrow \gamma\eta_b$
10 Obtained using $B(\gamma(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$ and $B(\gamma(3S) \rightarrow \gamma\eta_b) = (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$. This limit is equivalent to $B(\eta_b \rightarrow \mu^+\mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.					
$<8 \times 10^{-2}$	90	AUBERT	09P	BABR	$e^+ e^- \rightarrow \gamma\tau^+\tau^-$

 $\eta_b(1S)$ REFERENCES

TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)