

**$f_4(2300)$**  $I^G(J^{PC}) = 0^+(4^{++})$ 

## OMITTED FROM SUMMARY TABLE

This entry was previously called  $U_0(2350)$ . Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also  $\rho(2150)$ ,  $f_2(2150)$ ,  $\rho_3(2250)$ ,  $\rho_5(2350)$ .

 **$f_4(2300)$  MASS** **$\bar{p}p \rightarrow \pi\pi$  or  $\bar{K}K$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
~ 2314	HASAN	94	$\bar{p}p \rightarrow \pi\pi$
~ 2300	<sup>1</sup> MARTIN	80B	RVUE
~ 2300	<sup>1</sup> MARTIN	80C	RVUE
~ 2340	<sup>2</sup> CARTER	78B	CNTR 0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2330	DULUDE	78B	OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$
~ 2310	<sup>3</sup> CARTER	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

<sup>1</sup>  $I(J^P) = 0(4^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .<sup>2</sup>  $I(J^P) = 0(4^+)$  from Barrelet-zero analysis.<sup>3</sup>  $I(J^P) = 0(4^+)$  from amplitude analysis.**S-CHANNEL  $\bar{p}p$  or  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
2283 ± 17	<sup>4</sup> ANISOVICH	00J	SPEC
~ 2380	<sup>5</sup> CUTTS	78B	CNTR 0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2345 ± 15	<sup>5,6</sup> COUPLAND	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2359 ± 2	<sup>5,7</sup> ALSPECTOR	73	CNTR $\bar{p}p$ S channel
2375 ± 10	ABRAMS	70	CNTR S channel $\bar{N}N$

<sup>4</sup> From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on  $\bar{p}p \rightarrow \eta \pi^0 \pi^0$ ,  $\pi^0 \pi^0$ ,  $\eta \eta$ ,  $\eta \eta'$ ,  $\pi^+ \pi^-$ .<sup>5</sup> Isospins 0 and 1 not separated.<sup>6</sup> From a fit to the total elastic cross section.<sup>7</sup> Referred to as  $U$  or  $U$  region by ALSPECTOR 73. **$\pi^- p \rightarrow \eta \pi\pi n$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
2330 ± 20 ± 40	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

***pp* CENTRAL PRODUCTION**

VALUE (MeV)	DOCUMENT ID	COMMENT
<b><math>2320 \pm 60</math> OUR ESTIMATE</b>		
• • • We do not use the following data for averages, fits, limits, etc. • • •		
2332 $\pm$ 15	BARBERIS 00F 450 $pp \rightarrow p_f \omega \omega p_s$	

 **$f_4(2300)$  WIDTH** **$\bar{p}p \rightarrow \pi\pi$  or  $\bar{K}K$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$\sim 278$	HASAN 94	RVUE	$\bar{p}p \rightarrow \pi\pi$
$\sim 200$	8 MARTIN 80C	RVUE	
$\sim 150$	9 CARTER 78B	CNTR	$0.7\text{--}2.4 \bar{p}p \rightarrow K^- K^+$
$\sim 210$	10 CARTER 77	CNTR	$0.7\text{--}2.4 \bar{p}p \rightarrow \pi\pi$
$8 I(J^P) = 0(4^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$ .			
$9 I(J^P) = 0(4^+)$ from Barrelet-zero analysis.			
$10 I(J^P) = 0(4^+)$ from amplitude analysis.			

 **$S$ -CHANNEL  $\bar{p}p$  or  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$310 \pm 25$	11 ANISOVICH 00J	SPEC	
$135^{+150}_{-65}$	12,13 COUPLAND 77	CNTR	$0.7\text{--}2.4 \bar{p}p \rightarrow \bar{p}p$
$165^{+18}_{-8}$	13 ALSPECTOR 73	CNTR	$\bar{p}p$ <i>S</i> channel
$\sim 190$	ABRAMS 70	CNTR	<i>S</i> channel $\bar{N}N$
11 From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on $\bar{p}p \rightarrow \eta\pi^0\pi^0$ , $\pi^0\pi^0$ , $\eta\eta$ , $\eta\eta'$ , $\pi^+\pi^-$ .			
12 From a fit to the total elastic cross section.			
13 Isospins 0 and 1 not separated.			

 **$\pi^- p \rightarrow \eta\pi\pi n$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$235 \pm 50 \pm 40$	AMELIN 00 VES 37	$\pi^- p \rightarrow \eta\pi^+\pi^- n$	

***pp* CENTRAL PRODUCTION**

VALUE (MeV)	DOCUMENT ID	COMMENT
<b><math>250 \pm 80</math> OUR ESTIMATE</b>		
• • • We do not use the following data for averages, fits, limits, etc. • • •		
260 $\pm$ 57	BARBERIS 00F 450 $pp \rightarrow p_f \omega \omega p_s$	

**$f_4(2300)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\rho$	seen
$\Gamma_2$ $\omega\omega$	seen
$\Gamma_3$ $\eta\pi\pi$	seen
$\Gamma_4$ $\pi\pi$	seen
$\Gamma_5$ $K\bar{K}$	seen
$\Gamma_6$ $N\bar{N}$	seen

 **$f_4(2300)$  BRANCHING RATIOS**

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$	$\Gamma_1/\Gamma_2$	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$		
$2.8 \pm 0.5$	BARBERIS 00F	$450 \text{ pp} \rightarrow p_f \omega\omega p_s$

 **$f_4(2300)$  REFERENCES**

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE	78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)