

$\omega(1420)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\omega(1420)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
(1400–1450) OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1470 ± 50	13.1k	¹ AULCHENKO	15A SND	$1.05\text{--}1.80 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1382 ± 23 ± 70		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
1350 ± 20 ± 20		AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
1400 ± 50 ± 130	1.2M	² ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1450 ± 10		³ HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
1373 ± 70	177	⁴ AKHMETSHIN	00D CMD2	$1.2\text{--}1.38 e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1370 ± 25	5095	ANISOVICH	00H SPEC	$0.0 p\bar{p} \rightarrow \omega \pi^0 \pi^0 \pi^0$
1400^{+100}_{-200}		⁵ ACHASOV	98H RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
~ 1400		⁶ ACHASOV	98H RVUE	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
~ 1460		⁷ ACHASOV	98H RVUE	$e^+ e^- \rightarrow K^+ K^-$
1440 ± 70		⁸ CLEGG	94 RVUE	
1419 ± 31	315	⁹ ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+ e^- \rightarrow \rho \pi$

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.³ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.⁴ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho \pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.⁵ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.⁶ Using the data from ANTONELLI 92.⁷ Using the data from IVANOV 81 and BISELLO 88B.⁸ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.⁹ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed $(+, -, +)$ phases. **$\omega(1420)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
(180–250) OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
880 ± 170	13.1k	¹⁰ AULCHENKO	15A SND	$1.05\text{--}1.80 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
130 ± 50 ± 100		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
450 ± 70 ± 70		AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
$870^{+500}_{-300} \pm 450$	1.2M	¹¹ ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
199 ± 15		¹² HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$

188 ± 45	177	¹³ AKHMETSHIN 00D	CMD2	1.2–1.38	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
360^{+100}_{-60}	5095	ANISOVICH	00H	SPEC	$0.0 p\bar{p} \rightarrow \omega \pi^0 \pi^0 \pi^0$
240 ± 70		¹⁴ CLEGG	94	RVUE	
174 ± 59	315	¹⁵ ANTONELLI	92	DM2	$1.34–2.4 e^+ e^- \rightarrow \rho \pi$

¹⁰ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

¹¹ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

¹² Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

¹³ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho \pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

¹⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹⁵ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed $(+, -, +)$ phases.

$\omega(1420)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho \pi$	dominant
$\Gamma_2 \omega \pi \pi$	seen
$\Gamma_3 b_1(1235) \pi$	seen
$\Gamma_4 e^+ e^-$	seen
$\Gamma_5 \pi^0 \gamma$	

$$\omega(1420) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$$

$\Gamma(\rho \pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma \times \Gamma_4/\Gamma$
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.73 ± 0.08	13.1k ¹⁶ AULCHENKO 15A SND $1.05–1.80 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.82 $\pm 0.05 \pm 0.06$	AUBERT,B 04N BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
0.65 $\pm 0.13 \pm 0.21$	1.2M ^{17,18} ACHASOV 03D RVUE $0.44–2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.625 ± 0.160	19,20 CLEGG 94 RVUE
0.466 ± 0.178	21,22 ANTONELLI 92 DM2 $1.34–2.4 e^+ e^- \rightarrow \rho \pi$

¹⁶ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

¹⁷ Calculated by us from the cross section at the peak.

¹⁸ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

¹⁹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

²⁰ From the partial and leptonic width given by the authors.

²¹ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed $(+, -, +)$ phases.

²² From the product of the leptonic width and partial branching ratio given by the authors.

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma \times \Gamma_4/\Gamma$
--	--

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma \times \Gamma_4/\Gamma$		
<hr/>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
19.7 \pm 5.7	AUBERT 07AU BABR	10.6	$e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1.9 \pm 1.9	23 AKHMETSHIN 00D	CMD2	1.2–2.4 $e^+e^- \rightarrow \omega\pi^+\pi^-$
23 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.			

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma \times \Gamma_4/\Gamma$
---	--

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma \times \Gamma_4/\Gamma$		
<hr/>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.03 $^{+0.70}_{-0.75}$	24 AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
24 Using 1420 MeV and 220 MeV for the $\omega(1420)$ mass and width.			

$\omega(1420)$ BRANCHING RATIOS

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
--	-------------------

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$	Γ_2/Γ		
<hr/>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.301 \pm 0.029 possibly seen	25 HENNER 02 AKHMETSHIN 00D	RVUE CMD2	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$	Γ_2/Γ_3
---	---------------------

$\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$	Γ_2/Γ_3			
<hr/>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.60 \pm 0.16	5095	ANISOVICH 00H	SPEC	$0.0 p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
---	-------------------

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	Γ_1/Γ		
<hr/>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.699 \pm 0.029	25 HENNER 02	RVUE	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$	Γ_4/Γ
--	-------------------

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$	Γ_4/Γ			
<hr/>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 6.6	1.2M	26,27 ACHASOV 03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

23 ± 1 25 HENNER 02 RVUE $1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

25 Assuming that the $\omega(1420)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

26 Calculated by us from the cross section at the peak.

27 Assuming that the $\omega(1420)$ decays into $\rho\pi$ only.

$\omega(1420)$ REFERENCES

AULCHENKO	15A	JETP 121 27 Translated from ZETF 148 34.	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164 Translated from ZETFP 46 132.	L.M. Barkov <i>et al.</i>	(NOVO)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)