

# $\Upsilon(2S)$

 $I^G(J^{PC}) = 0^-(1^{--})$ 

## $\Upsilon(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10023.26±0.31 OUR AVERAGE</b>			
10023.5 ± 0.5	1 ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
10023.1 ± 0.4	BARBER 84	REDE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10023.6 ± 0.5	2,3 BARU	86B REDE	$e^+ e^- \rightarrow$ hadrons
1 Reanalysis of BARU 86B using new electron mass (COHEN 87).			
2 Reanalysis of ARTAMONOV 84.			
3 Superseded by ARTAMONOV 00.			

## $m\Upsilon(3S) - m\Upsilon(2S)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>331.50±0.02±0.13</b>	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

## $\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID
<b>31.98±2.63 OUR EVALUATION</b>	See the Note on "Width Determinations of the $\Upsilon$ States"

## $\Upsilon(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1 \Upsilon(1S)\pi^+\pi^-$	(17.85 ± 0.26) %	
$\Gamma_2 \Upsilon(1S)\pi^0\pi^0$	( 8.6 ± 0.4 ) %	
$\Gamma_3 \tau^+\tau^-$	( 2.00 ± 0.21 ) %	
$\Gamma_4 \mu^+\mu^-$	( 1.93 ± 0.17 ) %	S=2.2
$\Gamma_5 e^+e^-$	( 1.91 ± 0.16 ) %	
$\Gamma_6 \Upsilon(1S)\pi^0$	< 4 $\times 10^{-5}$	CL=90%
$\Gamma_7 \Upsilon(1S)\eta$	( 2.9 ± 0.4 ) $\times 10^{-4}$	S=2.0
$\Gamma_8 J/\psi(1S)$ anything	< 6 $\times 10^{-3}$	CL=90%
$\Gamma_9 J/\psi(1S)\eta_c$	< 5.4 $\times 10^{-6}$	CL=90%
$\Gamma_{10} J/\psi(1S)\chi_{c0}$	< 3.4 $\times 10^{-6}$	CL=90%
$\Gamma_{11} J/\psi(1S)\chi_{c1}$	< 1.2 $\times 10^{-6}$	CL=90%
$\Gamma_{12} J/\psi(1S)\chi_{c2}$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{13} J/\psi(1S)\eta_c(2S)$	< 2.5 $\times 10^{-6}$	CL=90%
$\Gamma_{14} J/\psi(1S)X(3940)$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{15} J/\psi(1S)X(4160)$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{16} \psi(2S)\eta_c$	< 5.1 $\times 10^{-6}$	CL=90%

$\Gamma_{17}$	$\psi(2S)\chi_{c0}$	< 4.7	$\times 10^{-6}$	CL=90%
$\Gamma_{18}$	$\psi(2S)\chi_{c1}$	< 2.5	$\times 10^{-6}$	CL=90%
$\Gamma_{19}$	$\psi(2S)\chi_{c2}$	< 1.9	$\times 10^{-6}$	CL=90%
$\Gamma_{20}$	$\psi(2S)\eta_c(2S)$	< 3.3	$\times 10^{-6}$	CL=90%
$\Gamma_{21}$	$\psi(2S)X(3940)$	< 3.9	$\times 10^{-6}$	CL=90%
$\Gamma_{22}$	$\psi(2S)X(4160)$	< 3.9	$\times 10^{-6}$	CL=90%
$\Gamma_{23}$	$\overline{^2H}$ anything	( 2.78 $\pm$ 0.30 )	$\times 10^{-5}$	S=1.2
$\Gamma_{24}$	hadrons	(94 $\pm$ 11 )	%	
$\Gamma_{25}$	$ggg$	(58.8 $\pm$ 1.2 )	%	
$\Gamma_{26}$	$\gamma gg$	( 1.87 $\pm$ 0.28 )	%	
$\Gamma_{27}$	$\phi K^+K^-$	( 1.6 $\pm$ 0.4 )	$\times 10^{-6}$	
$\Gamma_{28}$	$\omega\pi^+\pi^-$	< 2.58	$\times 10^{-6}$	CL=90%
$\Gamma_{29}$	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	( 2.3 $\pm$ 0.7 )	$\times 10^{-6}$	
$\Gamma_{30}$	$\phi f'_2(1525)$	< 1.33	$\times 10^{-6}$	CL=90%
$\Gamma_{31}$	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%
$\Gamma_{32}$	$\rho(770)a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%
$\Gamma_{33}$	$K^*(892)^0 \overline{K}_2^*(1430)^0 + \text{c.c.}$	( 1.5 $\pm$ 0.6 )	$\times 10^{-6}$	
$\Gamma_{34}$	$K_1(1270)^\pm K^\mp$	< 3.22	$\times 10^{-6}$	CL=90%
$\Gamma_{35}$	$K_1(1400)^\pm K^\mp$	< 8.3	$\times 10^{-7}$	CL=90%
$\Gamma_{36}$	$b_1(1235)^\pm \pi^\mp$	< 4.0	$\times 10^{-7}$	CL=90%
$\Gamma_{37}$	$\rho\pi$	< 1.16	$\times 10^{-6}$	CL=90%
$\Gamma_{38}$	$\pi^+\pi^-\pi^0$	< 8.0	$\times 10^{-7}$	CL=90%
$\Gamma_{39}$	$\omega\pi^0$	< 1.63	$\times 10^{-6}$	CL=90%
$\Gamma_{40}$	$\pi^+\pi^-\pi^0\pi^0$	( 1.30 $\pm$ 0.28 )	$\times 10^{-5}$	
$\Gamma_{41}$	$K_S^0 K^+ \pi^- + \text{c.c.}$	( 1.14 $\pm$ 0.33 )	$\times 10^{-6}$	
$\Gamma_{42}$	$K^*(892)^0 \overline{K}^0 + \text{c.c.}$	< 4.22	$\times 10^{-6}$	CL=90%
$\Gamma_{43}$	$K^*(892)^- K^+ + \text{c.c.}$	< 1.45	$\times 10^{-6}$	CL=90%
$\Gamma_{44}$	Sum of 100 exclusive modes	( 2.90 $\pm$ 0.30 )	$\times 10^{-3}$	

**Radiative decays**

$\Gamma_{45}$	$\gamma \chi b_1(1P)$	( 6.9 $\pm$ 0.4 )	%
$\Gamma_{46}$	$\gamma \chi b_2(1P)$	( 7.15 $\pm$ 0.35 )	%
$\Gamma_{47}$	$\gamma \chi b_0(1P)$	( 3.8 $\pm$ 0.4 )	%
$\Gamma_{48}$	$\gamma f_0(1710)$	< 5.9	$\times 10^{-4}$
$\Gamma_{49}$	$\gamma f'_2(1525)$	< 5.3	$\times 10^{-4}$
$\Gamma_{50}$	$\gamma f_2(1270)$	< 2.41	$\times 10^{-4}$
$\Gamma_{51}$	$\gamma f_J(2220)$		
$\Gamma_{52}$	$\gamma \eta_c(1S)$	< 2.7	$\times 10^{-5}$
$\Gamma_{53}$	$\gamma \chi c_0$	< 1.0	$\times 10^{-4}$
$\Gamma_{54}$	$\gamma \chi c_1$	< 3.6	$\times 10^{-6}$
$\Gamma_{55}$	$\gamma \chi c_2$	< 1.5	$\times 10^{-5}$
$\Gamma_{56}$	$\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi$	< 8	$\times 10^{-7}$
$\Gamma_{57}$	$\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	< 2.4	$\times 10^{-6}$

$\Gamma_{58}$	$\gamma X(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%
$\Gamma_{59}$	$\gamma X(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%
$\Gamma_{60}$	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%
$\Gamma_{61}$	$\gamma \eta_b(1S)$	( 3.9 $\pm$ 1.5 )	$\times 10^{-4}$	
$\Gamma_{62}$	$\gamma \eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7	$\times 10^{-6}$	CL=90%
$\Gamma_{63}$	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9	$\times 10^{-6}$	CL=90%
$\Gamma_{64}$	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%
$\Gamma_{65}$	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%
$\Gamma_{66}$	$\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$	< 8.3	$\times 10^{-6}$	CL=90%

**Lepton Family number (*LF*) violating modes**

$\Gamma_{67}$	$e^\pm \tau^\mp$	<i>LF</i>	< 3.2	$\times 10^{-6}$	CL=90%
$\Gamma_{68}$	$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.3	$\times 10^{-6}$	CL=90%

[a] 1.5 GeV  $< m_X <$  5.0 GeV

---

**CONSTRAINED FIT INFORMATION**

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 11.8$  for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$\begin{matrix} x_7 & \boxed{\phantom{00}2} \\ & x_1 \end{matrix}$$


---

 **$\Gamma(2S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$** 

$$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_4 \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>6.5 <math>\pm</math> 1.5 <math>\pm</math> 1.0</b>	KOBEL	92	$e^+ e^- \rightarrow \mu^+ \mu^-$

$$\Gamma(\Gamma(1S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_1 \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>105.4 <math>\pm</math> 1.0 <math>\pm</math> 4.2</b>	11.8K	<sup>1</sup> AUBERT	08BP BABR	$10.58 \quad e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$

<sup>1</sup> Using  $B(\Gamma(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$  and  $B(\Gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{24}\Gamma_5/\Gamma$ 

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.577±0.009 OUR AVERAGE</b>			
0.581±0.004±0.009	<sup>1</sup> ROSNER 06	CLEO	$10.0 \text{ } e^+ e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	<sup>1</sup> BARU 96	MD1	$e^+ e^- \rightarrow \text{hadrons}$
0.54 ±0.04 ±0.02	<sup>1</sup> JAKUBOWSKI 88	CBAL	$e^+ e^- \rightarrow \text{hadrons}$
0.58 ±0.03 ±0.04	<sup>2</sup> GILES 84B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$
0.60 ±0.12 ±0.07	<sup>2</sup> ALBRECHT 82	DASP	$e^+ e^- \rightarrow \text{hadrons}$
0.54 ±0.07 +0.09 -0.05	<sup>2</sup> NICZYPORUK 81C	LENA	$e^+ e^- \rightarrow \text{hadrons}$
0.41 ±0.18	<sup>2</sup> BOCK 80	CNTR	$e^+ e^- \rightarrow \text{hadrons}$

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85. $\gamma(2S) \text{ PARTIAL WIDTHS}$  $\Gamma(e^+ e^-)$  $\Gamma_5$ 

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
<b>0.612±0.011 OUR EVALUATION</b>	

 $\gamma(2S) \text{ BRANCHING RATIOS}$  $\Gamma(\gamma(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ Abbreviation MM in the *COMMENT* field below stands for missing mass.

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>17.85±0.26 OUR FIT</b>				
<b>17.92±0.26 OUR AVERAGE</b>				
16.8 ±1.1 ±1.3	906k	<sup>1</sup> LEES 11c	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$
17.80±0.05±0.37	170k	<sup>2</sup> LEES 11L	BABR	$\gamma(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
18.02±0.02±0.61	851k	<sup>3</sup> BHARI 09	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- \text{MM}$
17.22±0.17±0.75	11.8K	<sup>4</sup> AUBERT 08BP	BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$
19.2 ±0.2 ±1.0	52.6k	<sup>5</sup> ALEXANDER 98	CLE2	$\pi^+ \pi^- \ell^+ \ell^-, \pi^+ \pi^- \text{MM}$
18.1 ±0.5 ±1.0	11.6k	ALBRECHT 87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \text{MM}$
16.9 ±4.0		GELPHMAN 85	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
19.1 ±1.2 ±0.6		BESSON 84	CLEO	$\pi^+ \pi^- \text{MM}$
18.9 ±2.6		FONSECA 84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
21 ±7	7	NICZYPORUK 81B	LENA	$e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$

<sup>1</sup> LEES 11c reports  $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

<sup>3</sup> A weighted average of the inclusive and exclusive results.

<sup>4</sup> Using  $B(\gamma(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$ ,  $B(\gamma(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$  and,  $\Gamma_{ee}(\gamma(2S)) = 0.612 \pm 0.011 \text{ keV}$ .

<sup>5</sup> Using  $B(\gamma(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$  and  $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$ .

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.6 ± 0.4 OUR AVERAGE</b>				
8.43 ± 0.16 ± 0.42	38k	<sup>1</sup> BHARI 09	CLEO	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.2 ± 0.6 ± 0.8	275	<sup>2</sup> ALEXANDER 98	CLE2	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.5 ± 1.9 ± 1.9	25	ALBRECHT 87	ARG	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
8.0 ± 1.5		GELPHMAN 85	CBAL	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
10.3 ± 2.3		FONSECA 84	CUSB	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$

<sup>1</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$ .

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_2/\Gamma_1$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
0.462 ± 0.037	<sup>1</sup> BHARI 09	CLEO	$e^+ e^- \rightarrow \Upsilon(2S)$

<sup>1</sup> Not independent of other values reported by BHARI 09.

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.00 ± 0.21 OUR AVERAGE</b>				
2.00 ± 0.12 ± 0.18	22k	<sup>1</sup> BESSON 07	CLEO	$e^+ e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+ \tau^-$
1.7 ± 1.5 ± 0.6		HAAS 84B	CLEO	$e^+ e^- \rightarrow \tau^+ \tau^-$

<sup>1</sup> BESSON 07 reports  $[\Gamma(\Upsilon(2S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+ \mu^-)] = 1.04 \pm 0.04 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 


<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0193 ± 0.0017 OUR AVERAGE</b>			Error includes scale factor of 2.2. See the ideogram below.		
0.0203 ± 0.0003 ± 0.0008		120k	ADAMS 05	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0122 ± 0.0028 ± 0.0019			<sup>1</sup> KOBEL 92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0138 ± 0.0025 ± 0.0015			KAARSBERG 89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.009 ± 0.006 ± 0.006			<sup>2</sup> ALBRECHT 85	ARG	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.018 ± 0.008 ± 0.005			HAAS 84B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$

**• • • We do not use the following data for averages, fits, limits, etc. • • •**

<0.038 90 NICZYPORUK 81C LENA  $e^+ e^- \rightarrow \mu^+ \mu^-$

<sup>1</sup> Taking into account interference between the resonance and continuum.

<sup>2</sup> Re-evaluated using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 0.026$ .



### $\Gamma(\tau^+ \tau^-)/\Gamma(\mu^+ \mu^-)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_3/\Gamma_4$
<b>1.04±0.04±0.05</b>	22k	BESSON	07	CLEO $e^+ e^- \rightarrow \gamma(2S)$	

### $\Gamma(\gamma(1S)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_6/\Gamma$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
< 4	90	<sup>1</sup> TAMPONI	13	BELL $e^+ e^- \rightarrow \gamma(1S)\pi^0$	
< 18	90	<sup>2</sup> HE	08A	CLEO $e^+ e^- \rightarrow \ell^+\ell^-\gamma\gamma$	
<110	90	ALEXANDER	98	CLE2 $e^+ e^- \rightarrow \ell^+\ell^-\gamma\gamma$	
<800	90	LURZ	87	CBAL $e^+ e^- \rightarrow \ell^+\ell^-\gamma\gamma$	

<sup>1</sup> TAMPONI 13 reports  $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^0)/\Gamma_{\text{total}}] / [\mathcal{B}(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)]$

<  $2.3 \times 10^{-4}$  which we multiply by our best value  $\mathcal{B}(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$ .

<sup>2</sup> Authors assume  $\mathcal{B}(\gamma(1S) \rightarrow e^+ e^-) + \mathcal{B}(\gamma(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .

### $\Gamma(\gamma(1S)\pi^0)/\Gamma(\gamma(1S)\pi^+\pi^-)$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_6/\Gamma_1$
<b>&lt;2.3</b>	90	TAMPONI	13	BELL $e^+ e^- \rightarrow \gamma(1S)\pi^0$	

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**2.9 ± 0.4 OUR FIT** Error includes scale factor of 2.0.**2.9 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.


$2.39 \pm 0.31 \pm 0.14$	112	<sup>1</sup> LEES	11L BABR	$\Upsilon(2S) \rightarrow \ell^+ \ell^- \eta$
$2.1 \begin{array}{l} +0.7 \\ -0.6 \end{array} \pm 0.3$	14	<sup>2</sup> HE	08A CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$

• • • We use the following data for averages but not for fits. • • •

$3.55 \pm 0.32 \pm 0.05$	241	<sup>3</sup> TAMPONI	13 BELL	$e^+ e^- \rightarrow \Upsilon(1S)\eta$
--------------------------	-----	----------------------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	1,4 AUBERT	08BP BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$
< 28	90	ALEXANDER98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$
< 50	90	ALBRECHT	87 ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$
< 70	90	LURZ	87 CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84 CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$
< 20	90	FONSECA	84 CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+ \pi^- \pi^0)$

<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .<sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .<sup>3</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = (17.85 \pm 0.26) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.<sup>4</sup> Using  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011 \text{ keV}$ .

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$  $\Gamma_7/\Gamma_1$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.64±0.25 OUR FIT</b>					Error includes scale factor of 2.0.
<b>1.99±0.14±0.11</b>	241	TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.35 \pm 0.17 \pm 0.08$	1	LEES	11L	BABR	$\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
$< 5.2$	90	2	AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

<sup>1</sup> Not independent of other values reported by LEES 11L.<sup>2</sup> Not independent of other values reported by AUBERT 08BP. $\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$  $\Gamma_6/\Gamma_7$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.13$	90	TAMPONI	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.006</b>	90	MASCHMANN	90	CBAL $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 5.4 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 3.4 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 1.2 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.0 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.5 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.0 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.0 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<5.1 \times 10^{-6}$	90

 $\Gamma_{16}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<4.7 \times 10^{-6}$	90

 $\Gamma_{17}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<2.5 \times 10^{-6}$	90

 $\Gamma_{18}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<1.9 \times 10^{-6}$	90

 $\Gamma_{19}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<3.3 \times 10^{-6}$	90

 $\Gamma_{20}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<3.9 \times 10^{-6}$	90

 $\Gamma_{21}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<3.9 \times 10^{-6}$	90

 $\Gamma_{22}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG 14	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>
<b><math>2.78^{+0.30}_{-0.26}</math> OUR AVERAGE</b>	Error includes scale factor of 1.2.

 $\Gamma_{23}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
LEES 14G	BABR	$e^+ e^- \rightarrow \overline{2H} X$

$2.64 \pm 0.11^{+0.26}_{-0.21}$	
$3.37 \pm 0.50 \pm 0.25$	58

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ASNER 07	CLEO	$e^+ e^- \rightarrow \overline{2H} X$

 $\Gamma(ggg)/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>
<b><math>58.8 \pm 1.2</math></b>	6M

 $\Gamma_{25}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<sup>1</sup> BESSON 06A	CLEO	$\gamma(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$  from BESSON 06A and PDG 08 values of  $B(\pi^+ \pi^- \gamma(1S)) = (18.1 \pm 0.4)\%$ ,  $B(\pi^0 \pi^0 \gamma(1S)) = (8.6 \pm 0.4)\%$ ,  $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$ , and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma gg)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

 $\Gamma(\gamma gg)/\Gamma(ggg)$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>
<b><math>3.18 \pm 0.04 \pm 0.47</math></b>	6M

 $\Gamma_{26}/\Gamma_{25}$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
BESSON 06A	CLEO	$\gamma(2S) \rightarrow (\gamma +) \text{ hadrons}$

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>
<b><math>1.58 \pm 0.33 \pm 0.18</math></b>	58

 $\Gamma_{27}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;2.58</b>	90

 $\Gamma_{28}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>
<b><math>2.32 \pm 0.40 \pm 0.54</math></b>	135

 $\Gamma_{29}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;1.33</b>	90

 $\Gamma_{30}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;0.57</b>	90

 $\Gamma_{31}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;0.88</b>	90

 $\Gamma_{32}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>
<b><math>1.53 \pm 0.52 \pm 0.19</math></b>	32

 $\Gamma_{33}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;3.22</b>	90

 $\Gamma_{34}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;0.83</b>	90

 $\Gamma_{35}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(b_1(1235)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;0.40</b>	90

 $\Gamma_{36}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>
<b>&lt;1.16</b>	90

 $\Gamma_{37}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
<b>&lt;0.80</b>	90

 $\Gamma_{38}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	$\gamma(2S) \rightarrow \pi^+\pi^-\pi^0$

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
<b>&lt;1.63</b>	90

 $\Gamma_{39}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	$\gamma(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>EVTS</u>
<b><math>13.0 \pm 1.9 \pm 2.1</math></b>	$261 \pm 37$

 $\Gamma_{40}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	$\gamma(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$

 $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>EVTS</u>
<b><math>1.14 \pm 0.30 \pm 0.13</math></b>	$40 \pm 10$	

 $\Gamma_{41}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2	90	<sup>1</sup> DOBBS	12A	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$
------	----	--------------------	-----	--

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
<b>&lt;4.22</b>	90

 $\Gamma_{42}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
<b>&lt;1.45</b>	90

 $\Gamma_{43}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>CL%</u>
<b>0.29 <math>\pm 0.03</math></b>	

 $\Gamma_{44}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1,2 DOBBS	12A	$\gamma(2S) \rightarrow \text{hadrons}$

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>EVTS</u>
<b>0.069 <math>\pm 0.004</math> OUR AVERAGE</b>	
0.0693 $\pm 0.0012 \pm 0.0041$	407k
0.069 $\pm 0.005 \pm 0.009$	
0.091 $\pm 0.018 \pm 0.022$	
0.065 $\pm 0.007 \pm 0.012$	
0.080 $\pm 0.017 \pm 0.016$	
0.059 $\pm 0.014$	

 $\Gamma_{45}/\Gamma$ 

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ARTUSO	05	$e^+e^- \rightarrow \gamma X$
EDWARDS	99	$\gamma(2S) \rightarrow \gamma\chi(1P)$
ALBRECHT	85E	$e^+e^- \rightarrow \gamma \text{conv. } X$
NERNST	85	$e^+e^- \rightarrow \gamma X$
HAAS	84	$e^+e^- \rightarrow \gamma \text{conv. } X$
KLOPFEN...	83	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$ 

VALUE	EVTS
<b>0.0715±0.0035 OUR AVERAGE</b>	
0.0724±0.0011±0.0040	410k
0.074 ± 0.005 ± 0.008	
0.098 ± 0.021 ± 0.024	
0.058 ± 0.007 ± 0.010	
0.102 ± 0.018 ± 0.021	
0.061 ± 0.014	

 $\Gamma_{46}/\Gamma$ 

DOCUMENT ID	TECN	COMMENT
ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
HAAS 84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ 

VALUE	EVTS
<b>0.038 ±0.004 OUR AVERAGE</b>	

 $\Gamma_{47}/\Gamma$ 

DOCUMENT ID	TECN	COMMENT
ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
HAAS 84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%
<59	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.9	90	2 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma\pi^+\pi^-$
-------	----	---------------	-----	---

<sup>1</sup> Re-evaluated assuming  $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$ .

<sup>2</sup> Includes unknown branching ratio of  $f_0(1710) \rightarrow \pi^+\pi^-$ .

 $\Gamma_{48}/\Gamma$  $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%
<53	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$

<sup>1</sup> Re-evaluated assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ .

 $\Gamma_{49}/\Gamma$  $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%
<24.1	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma\pi^+\pi^-$

<sup>1</sup> Using  $B(f_2(1270) \rightarrow \pi\pi) = 0.84$ .

 $\Gamma_{50}/\Gamma$  $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%
<6.8	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$

<sup>1</sup> Includes unknown branching ratio of  $f_J(2220) \rightarrow K^+ K^-$ .

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<2.7 \times 10^{-5}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{52}/\Gamma$  $\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<1.0 \times 10^{-4}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{53}/\Gamma$  $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<3.6 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{54}/\Gamma$  $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<1.5 \times 10^{-5}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{55}/\Gamma$  $\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<0.8 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{56}/\Gamma$  $\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<2.4 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{57}/\Gamma$  $\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<2.8 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{58}/\Gamma$  $\Gamma(\gamma X(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<1.2 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{59}/\Gamma$  $\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<1.3 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
WANG	11B	BELL

 $\Gamma_{60}/\Gamma$  $\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>
$3.9 \pm 1.1^{+1.1}_{-0.9}$	13 ± 5k	

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<sup>1</sup> AUBERT	09AQ BABR	$\gamma(2S) \rightarrow \gamma X$

 $\Gamma_{61}/\Gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<21$	90	LEES	11J	BABR	$\gamma(2S) \rightarrow X \gamma$
$< 8.4$	90	<sup>1</sup> BONVICINI	10	CLEO	$\gamma(2S) \rightarrow \gamma X$
$< 5.1$	90	<sup>2</sup> ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV.

<sup>2</sup> Superseded by BONVICINI 10.

$\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.7 \times 10^{-6}$	90	SANDILYA	13	$\gamma(2S) \rightarrow \gamma$ hadrons

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 4.9	90		SANDILYA	13	$\gamma(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$46.2^{+29.7}_{-14.2} \pm 10.6$	10	<sup>1</sup> DOBBS	12	$\gamma(2S) \rightarrow \gamma$ hadrons
---------------------------------	----	--------------------	----	---

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$   
( $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ )

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.95	95	ROSNER	07A	CLEO $e^+ e^- \rightarrow \gamma X$

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$   
( $0.3 \text{ GeV} < m_{A^0} < 7 \text{ GeV}$ )

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	<sup>1</sup> LEES	11H	BABR $\gamma(2S) \rightarrow \gamma$ hadrons

<sup>1</sup> For a narrow scalar or pseudoscalar  $A^0$ , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $1 \times 10^{-6}$  to  $8 \times 10^{-5}$ .

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	<sup>1</sup> AUBERT	09Z	BABR $e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

<sup>1</sup> For a narrow scalar or pseudoscalar  $a_1^0$  with mass in the range 212–9300 MeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{a_1^0}$  range from  $0.26 - 8.3 \times 10^{-6}$ .

---

— LEPTON FAMILY NUMBER (*LF*) VIOLATING MODES — $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	LEES	10B	BABR $e^+ e^- \rightarrow e^\pm \tau^\mp$

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 3.3	90	LEES	10B	BABR $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14.4	95	LOVE	08A	CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$
-------	----	------	-----	---

---

**$\Upsilon(2S)$  Cross-Particle Branching Ratios**

$$\mathbf{B}(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times \mathbf{B}(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.78 \pm 0.02 \pm 0.11</math></b>	906k	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

 **$\Upsilon(2S)$  REFERENCES**

LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	(BELLE Collab.)
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	(BELLE Collab.)
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(NOVO)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(Crystal Ball Collab.)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUELL...	88	HE $e^+ e^-$ Physics 412 Editors: A. Ali and P. Soeding, World Scientific, Singapore	W. Buchmuller, S. Cooper	(HANN, DESY, MIT)
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelfman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
NERNST	85	Translated from YAF 41 733. PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	(DESY, ARGUS Collab.+)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)

KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)

---