

$\Upsilon(3S)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\Upsilon(3S)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10355.2±0.5	1 ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10355.3±0.5	2,3 BARU	86B REDE	$e^+ e^- \rightarrow$ hadrons
1 Reanalysis of BARU 86B using new electron mass (COHEN 87).			
2 Reanalysis of ARTAMONOV 84.			
3 Superseded by ARTAMONOV 00.			

 $m_{\Upsilon(3S)} - m_{\Upsilon(2S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

 $\Upsilon(3S)$ WIDTH

VALUE (keV)	DOCUMENT ID
20.32±1.85 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

 $\Upsilon(3S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \Upsilon(2S)$ anything	(10.6 ± 0.8) %	
$\Gamma_2 \Upsilon(2S)\pi^+\pi^-$	(2.82 ± 0.18) %	S=1.6
$\Gamma_3 \Upsilon(2S)\pi^0\pi^0$	(1.85 ± 0.14) %	
$\Gamma_4 \Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
$\Gamma_5 \Upsilon(2S)\pi^0$	< 5.1 × 10 ⁻⁴	CL=90%
$\Gamma_6 \Upsilon(1S)\pi^+\pi^-$	(4.37 ± 0.08) %	
$\Gamma_7 \Upsilon(1S)\pi^0\pi^0$	(2.20 ± 0.13) %	
$\Gamma_8 \Upsilon(1S)\eta$	< 1 × 10 ⁻⁴	CL=90%
$\Gamma_9 \Upsilon(1S)\pi^0$	< 7 × 10 ⁻⁵	CL=90%
$\Gamma_{10} h_b(1P)\pi^0$	< 1.2 × 10 ⁻³	CL=90%
$\Gamma_{11} h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 ± 1.4) × 10 ⁻⁴	
$\Gamma_{12} h_b(1P)\pi^+\pi^-$	< 1.2 × 10 ⁻⁴	CL=90%
$\Gamma_{13} \tau^+\tau^-$	(2.29 ± 0.30) %	
$\Gamma_{14} \mu^+\mu^-$	(2.18 ± 0.21) %	S=2.1
$\Gamma_{15} e^+e^-$	seen	
Γ_{16} hadrons		
$\Gamma_{17} ggg$	(35.7 ± 2.6) %	
$\Gamma_{18} \gamma gg$	(9.7 ± 1.8) × 10 ⁻³	
$\Gamma_{19} {}^2H$ anything	(2.33 ± 0.33) × 10 ⁻⁵	
Radiative decays		
$\Gamma_{20} \gamma\chi_{b2}(2P)$	(13.1 ± 1.6) %	S=3.4
$\Gamma_{21} \gamma\chi_{b1}(2P)$	(12.6 ± 1.2) %	S=2.4

Γ_{22}	$\gamma\chi_{b0}(2P)$	(5.9 \pm 0.6) %	S=1.4
Γ_{23}	$\gamma\chi_{b2}(1P)$	(9.9 \pm 1.3) $\times 10^{-3}$	S=2.0
Γ_{24}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8 $\times 10^{-5}$	CL=90%
Γ_{25}	$\gamma\chi_{b1}(1P)$	(9 \pm 5) $\times 10^{-4}$	S=1.9
Γ_{26}	$\gamma\chi_{b0}(1P)$	(2.7 \pm 0.4) $\times 10^{-3}$	
Γ_{27}	$\gamma\eta_b(2S)$	< 6.2 $\times 10^{-4}$	CL=90%
Γ_{28}	$\gamma\eta_b(1S)$	(5.1 \pm 0.7) $\times 10^{-4}$	
Γ_{29}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 $\times 10^{-4}$	CL=95%
Γ_{30}	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	< 5.5 $\times 10^{-6}$	CL=90%
Γ_{31}	$\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[b] < 1.6 $\times 10^{-4}$	CL=90%

Lepton Family number (*LF*) violating modes

Γ_{32}	$e^\pm\tau^\mp$	<i>LF</i>	< 4.2 $\times 10^{-6}$	CL=90%
Γ_{33}	$\mu^\pm\tau^\mp$	<i>LF</i>	< 3.1 $\times 10^{-6}$	CL=90%

[a] 1.5 GeV $< m_X <$ 5.0 GeV[b] For $m_{\tau^+\tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. **$\Upsilon(3S)\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$**

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_{16}\Gamma_{15}/\Gamma$
<i>VALUE</i> (keV)	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.414 \pm 0.007 OUR AVERAGE			
0.413 \pm 0.004 \pm 0.006	ROSNER	06	CLEO $10.4 e^+e^- \rightarrow$ hadrons
0.45 \pm 0.03 \pm 0.03	⁴ GILES	84B	CLEO $e^+e^- \rightarrow$ hadrons

⁴ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_6\Gamma_{15}/\Gamma$
<i>VALUE</i> (eV)	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
18.46 \pm 0.27 \pm 0.77	6.4K	⁵ AUBERT	08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
5 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.			

 $\Upsilon(3S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$			Γ_{15}
<i>VALUE</i> (keV)	<i>DOCUMENT ID</i>		
0.443 \pm 0.008 OUR EVALUATION			

 $\Upsilon(3S)$ BRANCHING RATIOS

$\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}}$			Γ_1/Γ
<i>VALUE</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
0.106 \pm 0.008 OUR AVERAGE			
0.1023 \pm 0.0105	4625	6,7,8 BUTLER	94B CLE2 $e^+e^- \rightarrow \ell^+\ell^-X$
0.111 \pm 0.012	4891	7,8, ⁹ BROCK	91 CLEO $e^+e^- \rightarrow \pi^+\pi^-X, \pi^+\pi^-\ell^+\ell^-$

⁶ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

⁸ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$.

⁹ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.82 ± 0.18 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.

$3.00 \pm 0.02 \pm 0.14$	543k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
$2.40 \pm 0.10 \pm 0.26$	800	¹⁰ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-e^+e^-$
3.12 ± 0.49	980	^{11,12} BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
2.13 ± 0.38	974	¹³ BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X, \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.82 \pm 0.65 \pm 0.53$	138	¹³ WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.1 ± 2.0	5	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$


¹⁰ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

¹¹ From the exclusive mode.

¹² Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

¹³ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

WEIGHTED AVERAGE
 2.82 ± 0.18 (Error scaled by 1.6)



$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ (units 10^{-2})

$\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.85 ± 0.14 OUR AVERAGE				
1.82 $\pm 0.09 \pm 0.12$	4391	¹⁴ BHARI	09	CLEO $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
2.16 ± 0.39		^{15,16} BUTLER	94B	CLE2 $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
1.7 $\pm 0.5 \pm 0.2$	10	¹⁷ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
¹⁴ Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.06\%$.				
¹⁵ $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.				
¹⁶ From the exclusive mode.				
¹⁷ $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.				

 $\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0502 ± 0.0069	¹⁸ BUTLER	94B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$

¹⁸ From the exclusive mode.

 $\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.51	90	¹⁹ HE	08A	CLEO $e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$

¹⁹ Authors assume $B(\Upsilon(2S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.06\%$.

 $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

Abbreviation MM in the *COMMENT* field below stands for missing mass.

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.37 ± 0.08 OUR AVERAGE				
4.32 $\pm 0.07 \pm 0.13$	90k	²⁰ LEES	11L	BABR $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$
4.46 $\pm 0.01 \pm 0.13$	190k	²¹ BHARI	09	CLEO $e^+ e^- \rightarrow \pi^+ \pi^-$ MM
4.17 $\pm 0.06 \pm 0.19$	6.4K	²² AUBERT	08BP	BABR $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$
4.52 ± 0.35	11830	²³ BUTLER	94B	CLE2 $e^+ e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$
4.46 $\pm 0.34 \pm 0.50$	451	²³ WU	93	CUSB $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$
4.46 ± 0.30	11221	²³ BROCK	91	CLEO $e^+ e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9 ± 1.0	22	GREEN	82	CLEO $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$
3.9 ± 1.3	26	MAGERAS	82	CUSB $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$

²⁰ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

²¹ A weighted average of the inclusive and exclusive results.

²² Using $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

²³ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_6

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.577 $\pm 0.026 \pm 0.060$	800	24 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
24 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.				

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.20 ± 0.13 OUR AVERAGE				
2.24 $\pm 0.09 \pm 0.11$	6584	25 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99 ± 0.34	56	26 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 $\pm 0.4 \pm 0.3$	33	27 HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
25 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$. 26 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality. 27 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.				

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_6

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.501 ± 0.043	28 BHARI	09 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$
28 Not independent of other values reported by BHARI 09.			

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.1	90	29 LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.8	90	29,30 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
<0.18	90	31 HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
<2.2	90	BROCK	91 CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
29 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$. 30 Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV. 31 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.				

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.23	90	32 LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.9	90	33 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$
32 Not independent of other values reported by LEES 11L. 33 Not independent of other values reported by AUBERT 08BP.				

$\Gamma(\gamma(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.07	90	34 HE	08A CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$

³⁴ Authors assume $B(\gamma(1S) \rightarrow e^+ e^-) + B(\gamma(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

 $\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2 × 10⁻³	90	35 GE	11 CLEO	$\gamma(3S) \rightarrow \pi^0$ anything

³⁵ Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.3 ± 1.1 ± 0.9	LEES	11k BABR	$\gamma(3S) \rightarrow \eta_b \gamma \pi^0$

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.2	90	36 LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 18	36 BUTLER	94B CLE2	$e^+ e^- \rightarrow \pi^+ \pi^- X$
< 15	36 BROCK	91 CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- X$

³⁶ For $M(h_b(1P)) = 9900$ MeV.

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.29 ± 0.21 ± 0.22	15k	37 BESSON	07 CLEO	$e^+ e^- \rightarrow \gamma(3S) \rightarrow \tau^+ \tau^-$

³⁷ BESSON 07 reports $[\Gamma(\gamma(3S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}] / [B(\gamma(3S) \rightarrow \mu^+ \mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\gamma(3S) \rightarrow \mu^+ \mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.


 $\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_{13}/Γ_{14}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.05 ± 0.08 ± 0.05	15k	BESSON	07 CLEO	$e^+ e^- \rightarrow \gamma(3S)$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0218±0.0021 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.			
0.0239±0.0007±0.0010	81k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0202±0.0019±0.0033		CHEN	89B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0173±0.0015±0.0011		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
0.033 ± 0.013 ± 0.007	1096	ANDREWS	83	CLEO $e^+e^- \rightarrow \mu^+\mu^-$

WEIGHTED AVERAGE
0.0218±0.0021 (Error scaled by 2.1)



$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$$

 $\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
35.7±2.6	3M	38 BESSON	06A CLEO	$\gamma(3S) \rightarrow \text{hadrons}$

³⁸ Calculated using BESSON 06A value of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\gamma(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+\pi^-\gamma(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0\pi^0\gamma(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma\chi b_2(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma\chi b_1(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma\chi b_0(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma\chi b_0(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+\mu^-) = (2.18 \pm 0.21)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ BESSON 06A value.

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.18	60k	39 BESSON	06A CLEO	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$

³⁹ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(ggg)/\Gamma_{\text{total}}$ BESSON 06A value.

$\Gamma(\gamma gg)/\Gamma(ggg)$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>
$2.72 \pm 0.06 \pm 0.49$	3M

 Γ_{18}/Γ_{17}

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
BESSON	06A	CLEO $\Upsilon(3S) \rightarrow (\gamma +)$ hadrons

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>
$2.33 \pm 0.15^{+0.31}_{-0.28}$

 Γ_{19}/Γ


<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
LEES	14G	BABR $e^+ e^- \rightarrow \overline{2H} X$

 $\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.131 ± 0.016 OUR AVERAGE	Error includes scale factor of 3.4. See the ideogram below.			

0.1579 $\pm 0.0017 \pm 0.0073$	568k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.111 $\pm 0.005 \pm 0.004$	10319	⁴⁰ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$
0.135 $\pm 0.003 \pm 0.017$	30741	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$

40 Supersedes NARAIN 91.


WEIGHTED AVERAGE
 0.131 ± 0.016 (Error scaled by 3.4) $\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$ $\Gamma(\gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.126 ± 0.012 OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.			

0.1454 $\pm 0.0018 \pm 0.0073$	537k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.115 $\pm 0.005 \pm 0.005$	11147	⁴¹ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$
0.105 $\pm 0.003 \pm 0.013$	25759	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$


41 Supersedes NARAIN 91.

 Γ_{21}/Γ



VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
0.059 ± 0.006 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.				
0.0677 ± 0.0020 ± 0.0065	225k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$	
0.060 ± 0.004 ± 0.006	4959	42 HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$	
0.049 +0.003 -0.004 ± 0.006	9903	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$	

⁴² Supersedes NARAIN 91.



\$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}	\$\Gamma_{23}/\Gamma\$
---	------------------------

\$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}	\$\Gamma_{23}/\Gamma\$				
9.9±1.3 OUR AVERAGE			Error includes scale factor of 2.0.		
7.5±1.2±0.5	126	43, ⁴⁴	KORNICER	11	CLEO \$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-
10.5±0.3 ^{+0.7} _{-0.6}	9.7k	LEES		11J	BABR \$\Upsilon(3S) \rightarrow X\gamma\$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19 seen	90	45	ASNER	08A	CLEO \$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}\$
		46	HEINTZ	92	CSB2 \$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-

⁴³ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

⁴⁴ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S))]$ $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S)) = (19.1 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁵ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))]$ $< 27.1 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

⁴⁶ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+\ell^-$.

\$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}	\$\Gamma_{25}/\Gamma\$
---	------------------------

\$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}	\$\Gamma_{25}/\Gamma\$				
0.9±0.5 OUR AVERAGE			Error includes scale factor of 1.9.		
1.6±0.5±0.1	50	47, ⁴⁸	KORNICER	11	CLEO \$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-
0.5±0.3 ^{+0.2} _{-0.1}		LEES		11J	BABR \$\Upsilon(3S) \rightarrow X\gamma\$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7 seen	90	49	ASNER	08A	CLEO \$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}\$
		50	HEINTZ	92	CSB2 \$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-

⁴⁷ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

⁴⁸ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S))]$ $= (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S)) = (33.9 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁹ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))]$ $< 2.5 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

⁵⁰ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+\ell^-$.

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.27±0.04 OUR AVERAGE					
0.27±0.04±0.02	2.3k	LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$
0.30±0.04±0.10	8.7k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.8	90	51 ASNER	08A	CLEO	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$
51 ASNER 08A reports $[\Gamma(\gamma(3S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma\chi_{b0}(1P))]$ < 21.9×10^{-2} which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$.					

 $\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 6.2				
< 6.2	90	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<19	90	LEES	11J	BABR $\gamma(3S) \rightarrow X\gamma$

 $\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.1±0.7 OUR AVERAGE					
7.1±1.8±1.3	2.3 ± 0.5k	52 BONVICINI	10	CLEO	$\gamma(3S) \rightarrow \gamma X$
4.8±0.5±0.6	19 ± 3k	52 AUBERT	09AQ	BABR	$\gamma(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8.5	90	LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$
4.8±0.5±1.2	19 ± 3k	52,53 AUBERT	08V	BABR	$\gamma(3S) \rightarrow \gamma X$
<4.3	90	54 ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$

52 Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

53 Systematic error re-evaluated by AUBERT 09AQ.

54 Superseded by BONVICINI 10.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{29}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.2	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.5	90	55 AUBERT	09Z	BABR $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$

55 For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from $0.27\text{--}5.5 \times 10^{-6}$.

$\Gamma(\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-)/\Gamma_{\text{total}}$	Γ_{31}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-4}$	90	56 AUBERT	09P BABR	$e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$

56 For a narrow scalar or pseudoscalar a_1^0 with $M(\tau^+\tau^-)$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of $M(\tau^+\tau^-)$ range from $1.5-16 \times 10^{-5}$.

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$	Γ_{24}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-5}$	90	57 LEES	11H BABR	$\Upsilon(3S) \rightarrow \gamma \text{ hadrons}$

57 For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

— LEPTON FAMILY NUMBER (LF) VIOLATING MODES —

$\Gamma(e^\pm\tau^\mp)/\Gamma_{\text{total}}$	Γ_{32}/Γ			
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.2	90	LEES	10B BABR	$e^+e^- \rightarrow e^\pm\tau^\mp$

$\Gamma(\mu^\pm\tau^\mp)/\Gamma_{\text{total}}$	Γ_{33}/Γ			
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	LEES	10B BABR	$e^+e^- \rightarrow \mu^\pm\tau^\mp$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<20.3	95	LOVE	08A CLEO	$e^+e^- \rightarrow \mu^\pm\tau^\mp$

$\Upsilon(3S)$ REFERENCES

LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)

ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+ e^-$ Physics 412	W. Buchmuller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)