# 8. NAMING SCHEME FOR HADRONS

Revised 2004 by M. Roos (University of Finland) and C.G. Wohl (LBNL).

### 8.1. Introduction

We introduced in the 1986 edition [1] a new naming scheme for the hadrons. Changes from older terminology affected mainly the heavier mesons made of the light (u, d, and s) quarks. Old and new names were listed alongside until 1994. Names also change from edition to edition because some characteristic like mass or spin changes. The Summary Tables give both the new and old names whenever a change occurred.

## 8.2. "Neutral-flavor" mesons (S = C = B = T = 0)

Table 8.1 shows the names for mesons having the strangeness and all heavy-flavor quantum numbers equal to zero. The scheme is designed for all ordinary non-exotic mesons, but it will work for many exotic types too, if needed.

	$J^{PC} =$	$ \begin{cases} 0^{-+} \\ 2^{-+} \\ \vdots \end{cases} $	$^{1^{+-}}_{3^{+-}}$	$2^{}$	$0^{++}$ $1^{++}$ $\vdots$
$q\overline{q}$ content <sup>2S</sup>	$^{+1}L_J =$	$^{1}(L \operatorname{even})_{J}$	$^{1}(L \operatorname{odd})_{J}$	$^{3}(L \operatorname{even})_{J}$	$^{3}(L \mathrm{odd})_{J}$
$u\overline{d}, u\overline{u} - d\overline{d}, d\overline{u}$	(I=1)	$\pi$	b	ρ	a
$\left. \begin{array}{c} d\overline{d} + u\overline{u} \\ and/or \ s\overline{s} \end{array} \right\}$	(I=0)	$\eta,\eta^\prime$	h, h'	$\omega,\phi$	f, f'
$c\overline{c}\ b\overline{b}\ t\overline{t}$		$\eta_c$ $\eta_b$ $\eta_t$	$egin{array}{c} h_c \ h_b \ h_t \end{array}$	$\psi^\dagger \ \Upsilon \  heta$	$egin{array}{c} \chi_c \ \chi_b \ \chi_t \end{array}$

**Table 8.1:** Symbols for mesons with the strangeness and all heavy-flavor quantumnumbers equal to zero.

<sup>†</sup>The  $J/\psi$  remains the  $J/\psi$ .

First, we assign names to those states with quantum numbers compatible with being  $q\bar{q}$  states. The rows of the Table give the possible  $q\bar{q}$  content. The columns give the possible parity/charge-conjugation states,

PC = -+, +-, --, and ++;

these combinations correspond one-to-one with the angular-momentum state  ${}^{2S+1}L_J$  of the  $q\overline{q}$  system being

$$^{1}(L \text{ even})_{J}, \, ^{1}(L \text{ odd})_{J}, \, ^{3}(L \text{ even})_{J}, \text{ or } ^{3}(L \text{ odd})_{J}$$

Here S, L, and J are the spin, orbital, and total angular momenta of the  $q\overline{q}$  system. The quantum numbers are related by  $P = (-1)^{L+1}$ ,  $C = (-1)^{L+S}$ , and G parity  $= (-1)^{L+S+I}$ , where of course the C quantum number is only relevant to neutral mesons.

C. Patrignani et al. (Particle Data Group), Chin. Phys. C, 40, 100001 (2016)

October 1, 2016 19:59

#### 2 8. Naming scheme for hadrons

The entries in the Table give the meson names. The spin J is added as a subscript except for pseudoscalar and vector mesons, and the mass is added in parentheses for mesons that decay strongly. However, for the lightest meson resonances, we omit the mass.

Measurements of the mass, quark content (where relevant), and quantum numbers I, J, P, and C (or G) of a meson thus fix its symbol. Conversely, these properties may be inferred unambiguously from the symbol.

If the main symbol cannot be assigned because the quantum numbers are unknown, X is used. Sometimes it is not known whether a meson is mainly the isospin-0 mix of  $u\overline{u}$  and  $d\overline{d}$  or is mainly  $s\overline{s}$ . A prime (or pair  $\omega$ ,  $\phi$ ) may be used to distinguish two such mixing states.

We follow custom and use spectroscopic names such as  $\Upsilon(1S)$  as the primary name for most of those  $\psi$ ,  $\Upsilon$ , and  $\chi$  states whose spectroscopic identity is known. We use the form  $\Upsilon(9460)$  as an alternative, and as the primary name when the spectroscopic identity is not known.

Names are assigned for  $t\bar{t}$  mesons, although the top quark is evidently so heavy that it is expected to decay too rapidly for bound states to form.

Gluonium states or other mesons that are not  $q\overline{q}$  states are, if the quantum numbers are *not* exotic, to be named just as are the  $q\overline{q}$  mesons. Such states will probably be difficult to distinguish from  $q\overline{q}$  states and will likely mix with them, and we make no attempt to distinguish those "mostly gluonium" from those "mostly  $q\overline{q}$ ."

An "exotic" meson with  $J^{PC}$  quantum numbers that a  $q\bar{q}$  system cannot have, namely  $J^{PC} = 0^{--}, 0^{+-}, 1^{-+}, 2^{+-}, 3^{-+}, \cdots$ , would use the same symbol as does an ordinary meson with all the same quantum numbers as the exotic meson except for the *C* parity. But then the *J* subscript may still distinguish it; for example, an isospin-0  $1^{-+}$  meson could be denoted  $\omega_1$ .

## 8.3. Mesons with nonzero S, C, B, and/or T

Since the strangeness or a heavy flavor of these mesons is nonzero, none of them are eigenstates of charge conjugation, and in each of them one of the quarks is heavier than the other. The rules are:

1. The main symbol is an upper-case italic letter indicating the heavier quark as follows:

 $s \to \overline{K} \qquad c \to D \qquad b \to \overline{B} \qquad t \to T \;.$ 

We use the convention that the flavor and the charge of a quark have the same sign. Thus the strangeness of the s quark is negative, the charm of the c quark is positive, and the bottom of the b quark is negative. In addition,  $I_3$  of the u and d quarks are positive and negative, respectively. The effect of this convention is as follows: Any flavor carried by a charged meson has the same sign as its charge. Thus the  $K^+$ ,  $D^+$ , and  $B^+$  have positive strangeness, charm, and bottom, respectively, and all have positive  $I_3$ . The  $D_s^+$  has positive charm and strangeness. Furthermore, the  $\Delta$ (flavor) =  $\Delta Q$  rule, best known for the kaons, applies to every flavor.

- 2. If the lighter quark is not a u or a d quark, its identity is given by a subscript. The  $D_s^+$  is an example.
- 3. If the spin-parity is in the "normal" series,  $J^P = 0^+, 1^-, 2^+, \cdots$ , a superscript "\*" is added.
- 4. The spin is added as a subscript except for pseudoscalar or vector mesons.

### 8.4. Ordinary (3-quark) baryons

The symbols N,  $\Delta$ ,  $\Lambda$ ,  $\Sigma$ ,  $\Xi$ , and  $\Omega$  used for more than 30 years for the baryons made of light quarks (u, d, and s quarks) tell the isospin and quark content, and the same information is conveyed by the symbols used for the baryons containing one or more heavy quarks (c and b quarks). The rules are:

- 1. Baryons with three u and/or d quarks are N's (isospin 1/2) or  $\Delta$ 's (isospin 3/2).
- 2. Baryons with two u and/or d quarks are  $\Lambda$ 's (isospin 0) or  $\Sigma$ 's (isospin 1). If the third quark is a c, b, or t quark, its identity is given by a subscript.
- 3. Baryons with one u or d quark are  $\Xi$ 's (isospin 1/2). One or two subscripts are used if one or both of the remaining quarks are heavy: thus  $\Xi_c$ ,  $\Xi_{cc}$ ,  $\Xi_b$ , etc.\*
- 4. Baryons with no u or d quarks are  $\Omega$ 's (isospin 0), and subscripts indicate any heavy-quark content.
- 5. A baryon that decays strongly has its mass as part of its name. Thus  $p, \Sigma^-, \Omega^-, \Lambda_c^+, etc.$ , but  $\Delta(1232)^0, \Sigma(1385)^-, \Xi_c(2645)^+, etc.$

In short, the number of u plus d quarks together with the isospin determine the main symbol, and subscripts indicate any content of heavy quarks. A  $\Sigma$  always has isospin 1, an  $\Omega$  always has isospin 0, *etc*.

#### 8.5. Exotic baryons

In 2003, several experiments reported finding a strangeness S = +1, charge Q = +1baryon, and one experiment reported finding an S = -2, Q = -2 baryon. Baryons with such quantum numbers cannot be made from three quarks, and thus they are exotic. The S = +1 baryon, which once would have been called a Z, was quickly dubbed the  $\Theta(1540)^+$ , and we proposed to name the S = -2 baryon the  $\Phi(1860)$ . However, these "discoveries" were then completely ruled out by many experiments with far larger statistics: See our 2008 Review [2].

# 4 8. Naming scheme for hadrons

#### Footnote and Reference:

- \* Sometimes a prime is necessary to distinguish two  $\Xi_c$ 's in the same SU(n) multiplet. See the "Note on Charmed Baryons" in the Charmed Baryon Listings.
- 1. Particle Data Group: M. Aguilar-Benitez et al., Phys. Lett. 170B (1986).
- 2. Particle Data Group: C. Amsler et al., Phys. Lett. B667, 1 (2008).