

$K_2(1820)$

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this *Review* (PDG 04) under $K_2(1770)$.

 $K_2(1820)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1819 ± 12 OUR AVERAGE				
$1853 \pm 27^{+18}_{-35}$	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
1816 ± 13		² ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1840		³ DAUM	81C CNTR	$63K^- p \rightarrow K^- 2\pi p$
¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0σ .				
² From a partial wave analysis of the $K^- \omega$ system.				
³ From a partial wave analysis of the $K^- 2\pi$ system.				

 $K_2(1820)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
264 ± 34 OUR AVERAGE				
$167 \pm 58^{+82}_{-72}$	4289	⁴ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
276 ± 35		⁵ ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 230		⁶ DAUM	81C CNTR	$63K^- p \rightarrow K^- 2\pi p$
⁴ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0σ .				
⁵ From a partial wave analysis of the $K^- \omega$ system.				
⁶ From a partial wave analysis of the $K^- 2\pi$ system.				

 $K_2(1820)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\pi\pi$	
$\Gamma_2 K_2^*(1430)\pi$	seen
$\Gamma_3 K^*(892)\pi$	seen
$\Gamma_4 Kf_2(1270)$	seen
$\Gamma_5 K\omega$	seen
$\Gamma_6 K\phi$	seen

 $K_2(1820)$ BRANCHING RATIOS

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.77	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K}2\pi p$

$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$				Γ_3/Γ_1
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 0.05	DAUM	81C	CNTR	$63K^- p \rightarrow \bar{K}2\pi p$
$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$				
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 0.18	DAUM	81C	CNTR	$63K^- p \rightarrow \bar{K}2\pi p$
$\Gamma(K\phi)/\Gamma_{\text{total}}$				
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	4289	7 AAIJ	17C	LHCb $B^+ \rightarrow J/\psi\phi K^+$
7 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 3.0 σ .				

K₂(1820) REFERENCES

AAIJ Also	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	04	PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
ASTON	93	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
DAUM	81C	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
		NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)