

X(10650) $^\pm$ $I^G(J^P) = ?^+(1^+)$

OMITTED FROM SUMMARY TABLE

Observed by BONDAR 12 in $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

X(10650) $^\pm$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10652.2\pm1.5	¹ BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10656.7 \pm 5.0 ^{+1.1} _{-3.1}	² GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10650.7 \pm 1.5 ^{+0.5} _{-0.2}	² GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10651.2 \pm 1.0 ^{+0.4} _{-0.3}	² GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10657 \pm 6 \pm 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10651 \pm 2 \pm 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10652 \pm 1 \pm 2	³ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10654 \pm 3 \pm 1	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 \pm 2 \pm 3	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹ Average of the BONDAR 12 measurements in separate channels.

² Correlated with the corresponding result from BONDAR 12.

³ Superseded by the average measurement of BONDAR 12.

X(10650) $^\pm$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
11.5\pm 2.2	⁴ BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
12.1 ^{+11.3 + 2.7} _{-4.8 - 0.6}	⁵ GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
14.2 \pm 3.7 ^{+ 0.9} _{- 0.4}	⁵ GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
9.3 \pm 2.2 ^{+ 0.3} _{- 0.5}	⁵ GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
16.3 \pm 9.8 ^{+ 6.0} _{- 2.0}	⁶ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
13.3 \pm 3.3 ^{+ 4.0} _{- 3.0}	⁶ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
8.4 \pm 2.0 \pm 2.0	⁶ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
20.9 ^{+ 5.4 + 2.1} _{- 4.7 - 5.7}	⁶ BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
19 \pm 7 \pm 11	⁶ BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

⁴ Average of the BONDAR 12 measurements in separate channels.

⁵ Correlated with the corresponding result from BONDAR 12.

⁶ Superseded by the average measurement of BONDAR 12.

$X(10650)^+$ DECAY MODES $X(10650)^-$ decay modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \gamma(1S)\pi^+$	($1.7^{+0.8}_{-0.6}$) $\times 10^{-3}$
$\Gamma_2 \gamma(2S)\pi^+$	($1.4^{+0.6}_{-0.4}$) %
$\Gamma_3 \gamma(3S)\pi^+$	($1.6^{+0.7}_{-0.5}$) %
$\Gamma_4 h_b(1P)\pi^+$	($8.4^{+2.9}_{-2.4}$) %
$\Gamma_5 h_b(2P)\pi^+$	(15 ± 4) %
$\Gamma_6 B^+\bar{B}^0$	not seen
$\Gamma_7 B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	not seen
$\Gamma_8 B^{*+}\bar{B}^{*0}$	(74^{+4}_{-6}) %

 $X(10650)^{\pm}$ BRANCHING RATIOS **$\Gamma(\gamma(1S)\pi^+)/\Gamma_{\text{total}}$** **$\Gamma_1/\Gamma$**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.7^{+0.7+0.3}_{-0.6-0.2}$	7 GARMASH	16 BELL	$e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15 BELL	$e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$
seen	BONDAR	12 BELL	$e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$

7 Assuming the $X(10650)^{\pm}$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUKI 16. **$\Gamma(\gamma(2S)\pi^+)/\Gamma_{\text{total}}$** **$\Gamma_2/\Gamma$**

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.39^{+0.48+0.34}_{-0.38-0.23}$	8 GARMASH	16	$e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15 BELL	$e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$
seen	BONDAR	12 BELL	$e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$

8 Assuming the $X(10650)^{\pm}$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUKI 16. **$\Gamma(\gamma(3S)\pi^+)/\Gamma_{\text{total}}$** **$\Gamma_3/\Gamma$**

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.63^{+0.53+0.39}_{-0.42-0.28}$	9 GARMASH	16 BELL	$e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15 BELL	$e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$
seen	BONDAR	12 BELL	$e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$

⁹ Assuming the $X(10650)^{\pm}$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^*+\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

$\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$8.41^{+2.43+1.49}_{-2.12-1.06}$	10 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^*+\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	11 MIZUK	16 BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$
seen	12 BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$

¹⁰ Assuming the $X(10650)^{\pm}$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^*+\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

¹¹ Using $e^+ e^-$ energies near the $\gamma(11020)$.

¹² Using $e^+ e^-$ energies near the $\gamma(10860)$.

$\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$14.7^{+3.2+2.8}_{-2.8-2.3}$	13 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^*+\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	14 MIZUK	16 BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$
seen	15 BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹³ Assuming the $X(10650)^{\pm}$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^*+\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

¹⁴ Using $e^+ e^-$ energies near the $\gamma(11020)$.

¹⁵ Using $e^+ e^-$ energies near the $\gamma(10860)$.

$\Gamma(B^+\bar{B}^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+\bar{B}^0$

$[\Gamma(B^+\bar{B}^{*0}) + \Gamma(B^*+\bar{B}^0)]/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+\bar{B}^{*0}, \pi^- \bar{B}^0 B^*$

$\Gamma(B^*+\bar{B}^{*0})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$73.7^{+3.4+2.7}_{-4.4-3.5}$	161	16 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^*+\bar{B}^{*0}$

¹⁶ Assuming the $X(10650)^{\pm}$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^*+\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

Using the mass and width of the $X(10650)^{\pm}$ from BONDAR 12.

$\Gamma(B^{*+}\bar{B}^{*0})/\left[\Gamma(\gamma(1S)\pi^+) + \Gamma(\gamma(2S)\pi^+) + \Gamma(\gamma(3S)\pi^+)\right]$	$\Gamma_8/(\Gamma_1+\Gamma_2+\Gamma_3+\Gamma_4+\Gamma_5)$			
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.80^{+0.69}_{-0.40}{}^{+0.54}_{-0.36}$	161	17 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<p>17 Combined with the results of BONDAR 12 and MIZUK 16. Not independent from $X(10650)^{\pm}$ branching fractions to $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$.</p>				

$X(10650)^{\pm}$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)