


**X(4360)** $I^G(J^{PC}) = ?^?(1^{--})$ 

Seen in radiative return from  $e^+ e^-$  collisions at  $\sqrt{s} = 9.54\text{--}10.58$  GeV by AUBERT 07S, WANG 07D, and LEES 14F. See also the review under the  $X(3872)$  particle listings. (See the index for the page number.)

**X(4360) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4341 ± 8 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
4320.0 ± 10.4 ± 7.0		1 ABLIKIM	17B BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
4347 ± 6 ± 3 279	279	2 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4340 ± 16 ± 9 37	37	3 LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4355 + 9 -10	± 9 74	4 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4324 ± 24		5 AUBERT	07S BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4361 ± 9 ± 9 47	47	3 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

<sup>1</sup> From a three-resonance fit.<sup>2</sup> From a two-resonance fit. Supersedes WANG 07D.<sup>3</sup> From a two-resonance fit.<sup>4</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.<sup>5</sup> From a single-resonance fit. Systematic errors not estimated.

**X(4360) WIDTH**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>102 ± 9 OUR AVERAGE</b>				
101.4 $^{+25.3}_{-19.7}$ $\pm 10.2$		1 ABLIKIM	17B BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
103 ± 9 ± 5 279	279	2 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
94 ± 32 ± 13 37	37	3 LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
103 $^{+17}_{-15}$ $\pm 11$ 74	74	4 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
172 ± 33		5 AUBERT	07S BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
74 ± 15 ± 10 47	47	3 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
1 From a three-resonance fit. 2 From a two-resonance fit. Supersedes WANG 07D. 3 From a two-resonance fit. 4 From a combined fit of AUBERT 07S and WANG 07D data with two resonances. 5 From a single-resonance fit. Systematic errors not estimated.				

**X(4360) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 e^+ e^-$	
$\Gamma_2 \psi(2S) \pi^+ \pi^-$	seen
$\Gamma_3 \psi(3823) \pi^+ \pi^-$	possibly seen
$\Gamma_4 J/\psi \eta$	
$\Gamma_5 D^0 D^{*-} \pi^+$	
$\Gamma_6 \chi_{c1} \gamma$	
$\Gamma_7 \chi_{c2} \gamma$	

**X(4360)  $\Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$** 

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u><math>\Gamma_2 \Gamma_1 / \Gamma</math></u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
9.2 ± 0.6 ± 0.6 279	279	1 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
10.9 ± 0.6 ± 0.7 279	279	2 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
6.0 ± 1.0 ± 0.5 37	37	3 LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
7.2 ± 1.0 ± 0.6 37	37	4 LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
11.1 $^{+1.3}_{-1.2}$	74	5 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
12.3 ± 1.2	74	6 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
10.4 ± 1.7 ± 1.5 47	47	3 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	
11.8 ± 1.8 ± 1.4 47	47	4 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	

<sup>1</sup> Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

<sup>2</sup> Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

<sup>3</sup> Solution I of two equivalent solutions in a fit using two interfering resonances.

<sup>4</sup> Solution II of two equivalent solutions in a fit using two interfering resonances.<sup>5</sup> Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.<sup>6</sup> Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances. $\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_4\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
<6.8	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_6\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.57	90	1 HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ . $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_7\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	1 HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .**X(4360) BRANCHING RATIOS** $\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S) \pi^+ \pi^-)$  $\Gamma_5/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	PAKHLOVA 09	BELL	$e^+e^- \rightarrow X(4360) \rightarrow D^0 D^{*-} \pi^+$

 $\Gamma(\psi(3823) \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
possibly seen	19	1 ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$

<sup>1</sup> From a fit of  $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ ,  $\psi(3823) \rightarrow \chi_{c1}\gamma$  cross sections taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the  $X(4360)$  line shape. $\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma \times \Gamma_1/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.72 \times 10^{-6}$	90	1 PAKHLOVA 09	BELL	$e^+e^- \rightarrow X(4360) \rightarrow D^0 D^{*-} \pi^+$

<sup>1</sup> Using  $4355^{+9}_{-10} \pm 9$  MeV for the mass of  $X(4360)$ .**X(4360) REFERENCES**

ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BES III Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)