

**X(4700)** $I^G(J^{PC}) = 0^+(0^{++})$ 

## OMMITTED FROM SUMMARY TABLE

Seen by AAIJ 17C in  $B^+ \rightarrow X K^+$ ,  $X \rightarrow J/\psi\phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  with a significance (accounting for systematic uncertainties) of  $5.6\sigma$ .

**X(4700) MASS**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4704 <math>\pm 10</math> <math>^{+14}_{-24}</math></b>	4289	<sup>1</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $5.6\sigma$ .

**X(4700) WIDTH**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>120 <math>\pm 31</math> <math>^{+42}_{-33}</math></b>	4289	<sup>2</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$

<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $5.6\sigma$ .

**X(4700) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad J/\psi\phi$	seen

**X(4700) BRANCHING RATIOS**

<u><math>\Gamma(J/\psi\phi)/\Gamma_{\text{total}}</math></u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u><math>\Gamma_1/\Gamma</math></u>
<b>seen</b>	4289	<sup>3</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$	

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $5.6\sigma$ .

**X(4700) REFERENCES**

AAIJ Also	17C PRL 118 022003 PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP (LHCb Collab.)
--------------	--	-----------------------	-------------------------------------