

# $\chi_{b0}(1P)$

$I^G(J^{PC}) = 0^+(0^{++})$   
 $J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(2S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

## $\chi_{b0}(1P)$ MASS

| VALUE (MeV)                                 | DOCUMENT ID  |
|---|--|
| <b>9859.44 ± 0.42 ± 0.31 OUR EVALUATION</b> | From average $\gamma$ energy below, using $\Upsilon(2S)$<br>mass = $10023.26 \pm 0.31$ MeV |

### $m_{\chi_{b1}(1P)} - m_{\chi_{b0}(1P)}$

| VALUE (MeV)         | DOCUMENT ID | TECN     | COMMENT   |
|---------------------|-------------|----------|---|
| <b>32.49 ± 0.93</b> | LEES        | 14M BABR | $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ |

## $\gamma$ ENERGY IN $\Upsilon(2S)$ DECAY

| VALUE (MeV)   | DOCUMENT ID   | TECN | COMMENT  |
|---|---------------|------|--|
| <b>162.5 ± 0.4 OUR AVERAGE</b>  |               |      |  |
| 162.56 ± 0.19 ± 0.42  | ARTUSO 05     | CLEO | $\Upsilon(2S) \rightarrow \gamma X$              |
| 162.0 ± 0.8 ± 1.2   | EDWARDS 99    | CLE2 | $\Upsilon(2S) \rightarrow \gamma\chi(1P)$        |
| 162.1 ± 0.5 ± 1.4   | ALBRECHT 85E  | ARG  | $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$ |
| 163.8 ± 1.6 ± 2.7   | NERNST 85     | CBAL | $\Upsilon(2S) \rightarrow \gamma X$              |
| 158.0 ± 7 ± 1   | HAAS 84       | CLEO | $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |               |      |  |
| 149.4 ± 0.7 ± 5.0   | KLOPFEN... 83 | CUSB | $\Upsilon(2S) \rightarrow \gamma X$              |

## $\chi_{b0}(1P)$ DECAY MODES

| Mode  | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level |
|---|--------------------------------|------------------|
| $\Gamma_1 \gamma \Upsilon(1S)$              | ( 1.94 ± 0.27 ) %              |                  |
| $\Gamma_2 D^0 X$                            | < 10.4 %                       | 90%              |
| $\Gamma_3 \pi^+ \pi^- K^+ K^- \pi^0$        | < 1.6 $\times 10^{-4}$         | 90%              |
| $\Gamma_4 2\pi^+ \pi^- K^- K_S^0$           | < 5 $\times 10^{-5}$           | 90%              |
| $\Gamma_5 2\pi^+ \pi^- K^- K_S^0 2\pi^0$    | < 5 $\times 10^{-4}$           | 90%              |
| $\Gamma_6 2\pi^+ 2\pi^- 2\pi^0$             | < 2.1 $\times 10^{-4}$         | 90%              |
| $\Gamma_7 2\pi^+ 2\pi^- K^+ K^-$            | ( 1.1 ± 0.6 ) $\times 10^{-4}$ |                  |
| $\Gamma_8 2\pi^+ 2\pi^- K^+ K^- \pi^0$      | < 2.7 $\times 10^{-4}$         | 90%              |
| $\Gamma_9 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$     | < 5 $\times 10^{-4}$           | 90%              |
| $\Gamma_{10} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$ | < 1.6 $\times 10^{-4}$         | 90%              |
| $\Gamma_{11} 3\pi^+ 3\pi^-$                 | < 8 $\times 10^{-5}$           | 90%              |
| $\Gamma_{12} 3\pi^+ 3\pi^- 2\pi^0$          | < 6 $\times 10^{-4}$           | 90%              |

|               |                               |                                |     |  |
|---------------|-------------------------------|--------------------------------|-----|--|
| $\Gamma_{13}$ | $3\pi^+ 3\pi^- K^+ K^-$       | $(2.4 \pm 1.2) \times 10^{-4}$ |     |  |
| $\Gamma_{14}$ | $3\pi^+ 3\pi^- K^+ K^- \pi^0$ | $< 1.0 \times 10^{-3}$         | 90% |  |
| $\Gamma_{15}$ | $4\pi^+ 4\pi^-$               | $< 8 \times 10^{-5}$           | 90% |  |
| $\Gamma_{16}$ | $4\pi^+ 4\pi^- 2\pi^0$        | $< 2.1 \times 10^{-3}$         | 90% |  |
| $\Gamma_{17}$ | $J/\psi J/\psi$               | $< 7 \times 10^{-5}$           | 90% |  |
| $\Gamma_{18}$ | $J/\psi \psi(2S)$             | $< 1.2 \times 10^{-4}$         | 90% |  |
| $\Gamma_{19}$ | $\psi(2S) \psi(2S)$           | $< 3.1 \times 10^{-5}$         | 90% |  |

 **$\chi_{b0}(1P)$  BRANCHING RATIOS**

$$\Gamma(\gamma \Upsilon(1S)) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

| VALUE (%)   | CL% | EVTS | DOCUMENT ID  | TECN     | COMMENT  |  |
|---|-----|------|--------------|----------|--|--|
| <b>1.94 ± 0.27 OUR AVERAGE</b>  |     |      |              |          |  |  |
| 2.07 ± 0.24 ± 0.21  |     |      | 1,2 LEES     | 14M BABR | $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$   |  |
| 1.76 ± 0.30 ± 0.18  |     | 87   | 3,4 KORNICER | 11 CLEO  | $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$      |  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |      |              |          |  |  |
| < 4.6   | 90  |      | 5 LEES       | 11J BABR | $\Upsilon(2S) \rightarrow X \gamma$                    |  |
| < 6   | 90  |      | WALK         | 86 CBAL  | $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |  |
| < 11  | 90  |      | PAUSS        | 83 CUSB  | $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |  |

<sup>1</sup> LEES 14M quotes  $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) / \Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) / \Gamma_{\text{total}}$  =  $(7.75 \pm 0.91) \times 10^{-4}$  combining the results from samples of  $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with and without converted photons. Assumes  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

<sup>2</sup> LEES 14M reports  $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) / \Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (7.75 \pm 0.91) \times 10^{-4}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Assuming  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$ .

<sup>4</sup> KORNICER 11 reports  $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) / \Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (6.59 \pm 0.96 \pm 0.60) \times 10^{-4}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup> LEES 11J quotes a central value of  $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) / \Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) / \Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$ .

$$\Gamma(D^0 X) / \Gamma_{\text{total}} \quad \Gamma_2 / \Gamma$$

| VALUE                   | CL% | DOCUMENT ID | TECN    | COMMENT                                 |  |
|-------------------------|-----|-------------|---------|---|--|
| $< 10.4 \times 10^{-2}$ | 90  | 6,7 BRIERE  | 08 CLEO | $\Upsilon(2S) \rightarrow \gamma D^0 X$ |  |

<sup>6</sup> For  $p_{D^0} > 2.5$  GeV/c.

<sup>7</sup> The authors also present their result as  $(5.6 \pm 3.6 \pm 0.5) \times 10^{-2}$ .

$$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_3 / \Gamma$$

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT   |  |
|--------------------------|-----|-------------|----------|---|--|
| < 1.6                    | 90  | 8 ASNER     | 08A CLEO | $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$ |  |

<sup>8</sup> ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0) / \Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

| <u>VALUE (units <math>10^{-4}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>   |
|---|------------|--------------------|-------------|--|
| <b>&lt;0.5</b>                            | 90         | 9 ASNER            | 08A CLEO    | $\Gamma(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$ |

9 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 2 \times 10^{-6}$  which we divide by our best value  $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

| <u>VALUE (units <math>10^{-4}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|---|------------|--------------------|-------------|---|
| <b>&lt;5</b>                              | 90         | 10 ASNER           | 08A CLEO    | $\Gamma(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$ |

10 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 18 \times 10^{-6}$  which we divide by our best value  $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

| <u>VALUE (units <math>10^{-4}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                       |
|---|------------|--------------------|-------------|--|
| <b>&lt;2.1</b>                            | 90         | 11 ASNER           | 08A CLEO    | $\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$ |

11 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 8 \times 10^{-6}$  which we divide by our best value  $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

| <u>VALUE (units <math>10^{-4}</math>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|---|-------------|--------------------|-------------|---|
| <b>1.1 ± 0.6 ± 0.1</b>                    | 7           | 12 ASNER           | 08A CLEO    | $\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$ |

12 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] = (4 \pm 2 \pm 1) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

| <u>VALUE (units <math>10^{-4}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|---|------------|--------------------|-------------|---|
| <b>&lt;2.7</b>                            | 90         | 13 ASNER           | 08A CLEO    | $\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$ |

13 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 10 \times 10^{-6}$  which we divide by our best value  $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

| <u>VALUE (units <math>10^{-4}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>   |
|---|------------|--------------------|-------------|--|
| <b>&lt;5</b>                              | 90         | 14 ASNER           | 08A CLEO    | $\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$ |

14 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 20 \times 10^{-6}$  which we divide by our best value  $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT   |
|--------------------------|-----|-------------|----------|---|
| <b>&lt;1.6</b>           | 90  | 15 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$ |

15 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|--------------------------|-----|-------------|----------|---|
| <b>&lt;0.8</b>           | 90  | 16 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$ |

16 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT  |
|--------------------------|-----|-------------|----------|--|
| <b>&lt;6</b>             | 90  | 17 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$ |

17 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 22 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

| VALUE (units $10^{-4}$ )                | EVTS | DOCUMENT ID | TECN     | COMMENT   |
|---|------|-------------|----------|---|
| <b><math>2.4 \pm 1.2 \pm 0.2</math></b> | 9    | 18 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$ |

18 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] = (9 \pm 4 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT   |
|--------------------------|-----|-------------|----------|---|
| <b>&lt;10</b>            | 90  | 19 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$ |

19 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 37 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|--------------------------|-----|-------------|----------|---|
| <b>&lt;0.8</b>           | 90  | 20 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$ |

20 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN     | COMMENT  |
|--------------------------|-----|-------------|----------|--|
| <b>&lt;21</b>            | 90  | 21 ASNER    | 08A CLEO | $\gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$ |

21 ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 77 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

$\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

| VALUE (units $10^{-5}$ )   | CL% | DOCUMENT ID | TECN    | COMMENT                               |
|--|-----|-------------|---------|---------------------------------------|
| <7   | 90  | 22 SHEN     | 12 BELL | $\Gamma(2S) \rightarrow \gamma\psi X$ |
| $^{22}_{-22}$ SHEN 12 reports $< 7.1 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P))] \text{ assuming } B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . |     |             |         |                                       |

 $\Gamma(J/\psi\psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

| VALUE (units $10^{-5}$ )   | CL% | DOCUMENT ID | TECN    | COMMENT                               |
|--|-----|-------------|---------|---------------------------------------|
| <12  | 90  | 23 SHEN     | 12 BELL | $\Gamma(2S) \rightarrow \gamma\psi X$ |
| $^{23}_{-23}$ SHEN 12 reports $< 12 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi\psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P))] \text{ assuming } B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . |     |             |         |                                       |

 $\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

| VALUE (units $10^{-5}$ )  | CL% | DOCUMENT ID | TECN    | COMMENT                               |
|---|-----|-------------|---------|---------------------------------------|
| <3.1  | 90  | 24 SHEN     | 12 BELL | $\Gamma(2S) \rightarrow \gamma\psi X$ |
| $^{24}_{-24}$ SHEN 12 reports $< 3.1 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P))] \text{ assuming } B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . |     |             |         |                                       |

 $\chi_{b0}(1P)$  CROSS-PARTICLE BRANCHING RATIOS $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma \times \Gamma_{49}^{\Gamma(2S)}/\Gamma^{\Gamma(2S)}$ 

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                          |
|---|-----|-------------|----------|----------------------------------|
| < $1.7 \times 10^{-3}$  | 90  | 25 LEES     | 11J BABR | $\Gamma(2S) \rightarrow X\gamma$ |
| $^{25}_{-25}$ LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6)^{+3.7}_{-2.6} \times 10^{-4}$ and derives a 90% CL upper limit of $\Gamma(\gamma \Gamma(1S))/\Gamma_{\text{total}} < 4.6\%$ using $B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4)\%$ . |     |             |          |                                  |

 $B(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) \times B(\Gamma(1S) \rightarrow \ell^+\ell^-)$ 

| VALUE (units $10^{-5}$ )  | EVTS | DOCUMENT ID | TECN     | COMMENT   |
|---|------|-------------|----------|---|
| <b>1.67 ± 0.28 OUR AVERAGE</b>  |      |             |          |   |
| 2.9 $^{+1.7}_{-1.4}$ $^{+0.1}_{-0.8}$   |      | 26 LEES     | 14M BABR | $\Gamma(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ |
| 1.63 $\pm 0.24 \pm 0.15$  | 87   | KORNICER    | 11 CLEO  | $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$   |
| $^{26}_{-26}$ From a sample of $\Gamma(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon. |      |             |          |   |

$$[B(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(2S) \rightarrow \gamma\chi_{b0}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(2S) \rightarrow \gamma\chi_{b1}(1P))]$$

| VALUE (%)   | DOCUMENT ID | TECN     | COMMENT   |
|---|-------------|----------|---|
| 3.28 $\pm 0.37$   | 27 LEES     | 14M BABR | $\Gamma(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ |
| $^{27}_{-27}$ From a sample of $\Gamma(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ without converted photons. |             |          |   |

## $\chi_{b0}(1P)$ REFERENCES

|            |     |               |                               |                          |
|------------|-----|---------------|-------------------------------|--------------------------|
| LEES       | 14M | PR D90 112010 | J.P. Lees <i>et al.</i>       | (BABAR Collab.)          |
| SHEN       | 12  | PR D85 071102 | C.P. Shen <i>et al.</i>       | (BELLE Collab.)          |
| KORNICER   | 11  | PR D83 054003 | M. Kornicer <i>et al.</i>     | (CLEO Collab.)           |
| LEES       | 11J | PR D84 072002 | J.P. Lees <i>et al.</i>       | (BABAR Collab.)          |
| ASNER      | 08A | PR D78 091103 | D.M. Asner <i>et al.</i>      | (CLEO Collab.)           |
| BRIERE     | 08  | PR D78 092007 | R.A. Briere <i>et al.</i>     | (CLEO Collab.)           |
| ARTUSO     | 05  | PRL 94 032001 | M. Artuso <i>et al.</i>       | (CLEO Collab.)           |
| EDWARDS    | 99  | PR D59 032003 | K.W. Edwards <i>et al.</i>    | (CLEO Collab.)           |
| WALK       | 86  | PR D34 2611   | W.S. Walk <i>et al.</i>       | (Crystal Ball Collab.)   |
| ALBRECHT   | 85E | PL 160B 331   | H. Albrecht <i>et al.</i>     | (ARGUS Collab.)          |
| NERNST     | 85  | PRL 54 2195   | R. Nernst <i>et al.</i>       | (Crystal Ball Collab.)   |
| HAAS       | 84  | PRL 52 799    | J. Haas <i>et al.</i>         | (CLEO Collab.)           |
| KLOPFEN... | 83  | PRL 51 160    | C. Klopfenstein <i>et al.</i> | (CUSB Collab.)           |
| PAUSS      | 83  | PL 130B 439   | F. Pauss <i>et al.</i>        | (MPIM, COLU, CORN, LSU+) |