

**$\chi_{b2}(1P)$** 
 $I^G(J^{PC}) = 0^+(2^{++})$   
 $J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(2S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .  $J = 2$  from SKWARNICKI 87.

 **$\chi_{b2}(1P)$  MASS**VALUE (MeV)DOCUMENT ID
**9912.21±0.26±0.31 OUR EVALUATION** From average  $\gamma$  energy below, using  $\Upsilon(2S)$  mass =  $10023.26 \pm 0.31$  MeV
 **$m_{\chi_{b2}(1P)} - m_{\chi_{b1}(1P)}$** VALUE (MeV)DOCUMENT IDTECNCOMMENT
**19.10±0.25 OUR AVERAGE** Error includes scale factor of 1.1.

$19.81 \pm 0.65 \pm 0.20$	<sup>1</sup> AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$
$19.01 \pm 0.24$	LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$

<sup>1</sup> From the  $\chi_{bj}(1P) \rightarrow \Upsilon(1S)\gamma$  transition.
 **$\gamma$  ENERGY IN  $\Upsilon(2S)$  DECAY**VALUE (MeV)DOCUMENT IDTECNCOMMENT
**110.44±0.29 OUR AVERAGE** Error includes scale factor of 1.1.

$110.58 \pm 0.08 \pm 0.30$	ARTUSO	05	CLEO	$\Upsilon(2S) \rightarrow \gamma X$
$110.8 \pm 0.3 \pm 0.6$	EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
$107.0 \pm 1.1 \pm 1.3$	WALK	86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$110.6 \pm 0.3 \pm 0.9$	ALBRECHT	85E	ARG	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
$110.4 \pm 0.8 \pm 2.2$	NERNST	85	CBAL	$\Upsilon(2S) \rightarrow \gamma X$
$109.5 \pm 0.7 \pm 1.0$	HAAS	84	CLEO	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
$108.2 \pm 0.3 \pm 2.0$	KLOPFEN...	83	CUSB	$\Upsilon(2S) \rightarrow \gamma X$
$108.8 \pm 4.0$	PAUSS	83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

 **$\chi_{b2}(1P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 \gamma \Upsilon(1S)$	$(18.8 \pm 1.1) \%$	
$\Gamma_2 D^0 X$	$< 7.9 \%$	90%
$\Gamma_3 \pi^+ \pi^- K^+ K^- \pi^0$	$(8 \pm 5) \times 10^{-5}$	
$\Gamma_4 2\pi^+ \pi^- K^- K_S^0$	$< 1.0 \times 10^{-4}$	90%
$\Gamma_5 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$(5.3 \pm 2.4) \times 10^{-4}$	
$\Gamma_6 2\pi^+ 2\pi^- 2\pi^0$	$(3.5 \pm 1.4) \times 10^{-4}$	
$\Gamma_7 2\pi^+ 2\pi^- K^+ K^-$	$(1.1 \pm 0.4) \times 10^{-4}$	
$\Gamma_8 2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.1 \pm 0.9) \times 10^{-4}$	
$\Gamma_9 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(3.9 \pm 1.8) \times 10^{-4}$	

$\Gamma_{10}$	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 5 \times 10^{-4}$	90%
$\Gamma_{11}$	$3\pi^+ 3\pi^-$	$(7.0 \pm 3.1) \times 10^{-5}$	
$\Gamma_{12}$	$3\pi^+ 3\pi^- 2\pi^0$	$(1.0 \pm 0.4) \times 10^{-3}$	
$\Gamma_{13}$	$3\pi^+ 3\pi^- K^+ K^-$	$< 8 \times 10^{-5}$	90%
$\Gamma_{14}$	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(3.6 \pm 1.5) \times 10^{-4}$	
$\Gamma_{15}$	$4\pi^+ 4\pi^-$	$(8 \pm 4) \times 10^{-5}$	
$\Gamma_{16}$	$4\pi^+ 4\pi^- 2\pi^0$	$(1.8 \pm 0.7) \times 10^{-3}$	
$\Gamma_{17}$	$J/\psi J/\psi$	$< 4 \times 10^{-5}$	90%
$\Gamma_{18}$	$J/\psi \psi(2S)$	$< 5 \times 10^{-5}$	90%
$\Gamma_{19}$	$\psi(2S) \psi(2S)$	$< 1.6 \times 10^{-5}$	90%

 **$\chi_{b2}(1P)$  BRANCHING RATIOS**

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<b><math>0.188 \pm 0.011</math> OUR AVERAGE</b>	

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.188 \pm 0.011</math> OUR AVERAGE</b>				
0.185 $\pm 0.008 \pm 0.009$	2,3,4 LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
0.186 $\pm 0.011 \pm 0.009$	1770 4,5 KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$	
$0.194^{+0.014}_{-0.017} \pm 0.009$	8k 6 LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$	
0.25 $\pm 0.06 \pm 0.01$	35 4,7,8 WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
0.20 $\pm 0.05$	KLOPFEN...	83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

<sup>2</sup> LEES 14M quotes  $\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}$  =  $(1.32 \pm 0.06)\%$  combining the results from samples of  $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with and without converted photons.

<sup>3</sup> LEES 14M reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.32 \pm 0.06) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Assuming  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

<sup>5</sup> KORNICER 11 reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.33 \pm 0.04 \pm 0.07) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>6</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (13.9 \pm 0.5^{+0.9}_{-1.1}) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>7</sup> WALK 86 quotes  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (4.4 \pm 0.9 \pm 0.5)\%$ .

<sup>8</sup> WALK 86 reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (17.7 \pm 3.6 \pm 2.0) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;7.9 \times 10^{-2}</math></b>	90	9,10 BRIERE	08	CLEO $\gamma(2S) \rightarrow \gamma D^0 X$

<sup>9</sup> For  $p_{D^0} > 2.5$  GeV/c.

<sup>10</sup> The authors also present their result as  $(5.4 \pm 1.9 \pm 0.5) \times 10^{-2}$ .

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$			
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.84 \pm 0.50 \pm 0.04</math></b>	8	11 ASNER	08A	CLEO $\gamma(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

<sup>11</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 3 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$	$\Gamma_4/\Gamma$			
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.0</math></b>	90	12 ASNER	08A	CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

<sup>12</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] < 7 \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma$			
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.3 \pm 2.4 \pm 0.3</math></b>	11	13 ASNER	08A	CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

<sup>13</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (38 \pm 14 \pm 10) \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma$			
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.5 \pm 1.4 \pm 0.2</math></b>	19	14 ASNER	08A	CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

<sup>14</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (25 \pm 8 \pm 6) \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma$			
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.1 \pm 0.4 \pm 0.1</math></b>	14	15 ASNER	08A	CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

<sup>15</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (8 \pm 2 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1±0.9±0.1</b>	13	16 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
<sup>16</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (15 \pm 5 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.9±1.8±0.2</b>	11	17 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
<sup>17</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (28 \pm 11 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	18 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
<sup>18</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] < 36 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .				

$\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.70±0.31±0.03</b>	9	19 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<sup>19</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (5 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.2±3.6±0.5</b>	34	20 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
<sup>20</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (73 \pm 16 \pm 20) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.8</b>	90	21 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<sup>21</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .				

$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.6±1.5±0.2</b>	14	22 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
22 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (26 \pm 8 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.84±0.40±0.04</b>	7	23 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
23 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18±7±1</b>	29	24 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
24 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (132 \pm 31 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	25 SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$
25 SHEN 12 reports $< 4.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ .				

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	26 SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$
26 SHEN 12 reports $< 4.9 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ .				

 $\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.6</b>	90	27 SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$
27 SHEN 12 reports $< 1.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ .				

### $\chi_{b2}(1P)$ Cross-Particle Branching Ratios

$$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) / \Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_1 / \Gamma \times \Gamma_{48}^{\Upsilon(2S)} / \Gamma^{\Upsilon(2S)}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>13.9 \pm 0.5^{+0.9}_{-1.1}</math></b>	8k	LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$

$$B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.38 \pm 0.16</math> OUR AVERAGE</b>				

$3.63^{+0.36+0.18}_{-0.34-0.19}$  28 LEES 14M BABR  $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$

$3.29 \pm 0.09 \pm 0.16$  1770 KORNICER 11 CLEO  $e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$

$4.4 \pm 0.9 \pm 0.5$  35 WALK 86 CBAL  $\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

28 From a sample of  $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$  with converted photons.

$$[B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b><math>55.6 \pm 1.6</math></b>	29 LEES 14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$	

29 From a sample of  $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$  events without converted photons.

$$B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.8 \pm 0.5</math> OUR AVERAGE</b>				

$4.68^{+0.99}_{-0.92} \pm 0.37$  30 LEES 14M BABR  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

$3.56 \pm 0.40 \pm 0.41$  126 KORNICER 11 CLEO  $e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$

30 From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with converted photons.

### $\chi_{b2}(1P)$ REFERENCES

AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFENSTEIN	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)