

$f_0(1370)$

$I^G(J^{PC}) = 0^+(0^{++})$

See also the mini-reviews on scalar mesons under $f_0(500)$ (see the index for the page number) and on non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).

$f_0(1370)$ T-MATRIX POLE POSITION

Note that $\Gamma \approx 2 \operatorname{Im}(\sqrt{s_{\text{pole}}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1200–1500)–i(150–250) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(1290 \pm 50) - i(170^{+20}_{-40})$	¹ ANISOVICH 09 RVUE $0.0 \bar{p}p, \pi N$		
$(1373 \pm 15) - i(137 \pm 10)$	² BARGIOTTI 03 OBLX $\bar{p}p$		
$(1302 \pm 17) - i(166 \pm 18)$	³ BARBERIS 00C $450 pp \rightarrow p_f 4\pi p_s$		
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS 99D OMEG $450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$		
$(1406 \pm 19) - i(80 \pm 6)$	⁴ KAMINSKI 99 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$		
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH 98B RVUE Compilation		
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS 97B OMEG $450 pp \rightarrow pp 2(\pi^+ \pi^-)$		
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN 97C OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$		
$(1380 \pm 40) - i(180 \pm 25)$	ABELE 96B CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$		
$(1300 \pm 15) - i(115 \pm 8)$	BUGG 96 RVUE		
$(1330 \pm 50) - i(150 \pm 40)$	⁵ AMSLER 95B CBAR $\bar{p}p \rightarrow 3\pi^0$		
$(1360 \pm 35) - i(150–300)$	⁵ AMSLER 95C CBAR $\bar{p}p \rightarrow \pi^0 \eta\eta$		
$(1390 \pm 30) - i(190 \pm 40)$	⁶ AMSLER 95D CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$		
$1346 - i249$	^{7,8} JANSEN 95 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$		
$1214 - i168$	^{8,9} TORNQVIST 95 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$		
$1364 - i139$	AMSLER 94D CBAR $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$		
$(1365^{+20}_{-55}) - i(134 \pm 35)$	ANISOVICH 94 CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$		
$(1340 \pm 40) - i(127^{+30}_{-20})$	¹⁰ BUGG 94 RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$		
$(1430 \pm 5) - i(73 \pm 13)$	¹¹ KAMINSKI 94 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$		
$1420 - i220$	¹² AU 87 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$		

¹ Another pole is found at $(1510 \pm 130) - i(800^{+100}_{-150})$ MeV.

² Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

³ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

⁴ T-matrix pole on sheet ——.

⁵ Supersedes ANISOVICH 94.

⁶ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0$, $\pi^0 \eta\eta$, and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁷ Analysis of data from FALVARD 88.

⁸ The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

- ⁹ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.
- ¹⁰ Reanalysis of ANISOVICH 94 data.
- ¹¹ T-matrix pole on sheet III.
- ¹² Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

f₀(1370) BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETER

VALUE (MeV)
1200 to 1500 OUR ESTIMATE

DOCUMENT ID

ππ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1400 ± 40		1 AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 ^{+ 6 + 72} _{- 7 - 255}		2 UEHARA	08A BELL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1259 ± 55	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
1309 ± 1 ± 15		3 BUGG	07A RVUE	$0.0 p\bar{p} \rightarrow 3\pi^0$
1449 ± 13	4.3k	4 GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
1350 ± 50		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
1265 ± 30 ^{+ 20} _{- 35}		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1434 ± 18 ± 9	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
1308 ± 10		BARBERIS	99B OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
1315 ± 50		BELLAZZINI	99 GAM4	$450 pp \rightarrow p p \pi^0 \pi^0$
1315 ± 30		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
1280 ± 55		BERTIN	98 OBLX	$0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1186	5,6 TORNQVIST	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472 ± 12		ARMSTRONG	91 OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275 ± 20		BREAKSTONE	90 SFM	$62 pp \rightarrow pp\pi^+ \pi^-$
1420 ± 20		AKESSON	86 SPEC	$63 pp \rightarrow pp\pi^+ \pi^-$
1256		FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

¹ Breit-Wigner mass.

² Breit-Wigner mass. May also be the f₀(1500).

³ Reanalysis of ABELE 96C data.

⁴ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.

⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁶ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1360 \pm 31 \pm 28	430	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
1350 \pm 48 \pm 15	168	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
1440 \pm 6		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1391 \pm 10		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440 \pm 50		BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1463 \pm 9		ETKIN 82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$
1425 \pm 15		WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
~ 1300		POLYCHRO...	STRC	$7 \pi^- p \rightarrow n2K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² From a fit to a Breit-Wigner line shape with fixed $\Gamma = 346$ MeV. **4π MODE $2(\pi\pi)_S + \rho\rho$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1395 \pm 40		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1374 \pm 38		AMSLER 94	CBAR	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
1345 \pm 12		ADAMO 93	OBLX	$\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
1386 \pm 30		GASPERO 93	DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 1410	5751	¹ BETTINI 66	DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant. **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1262^{+51+82}_{-78-103}$	¹ UEHARA 10A	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1430	AMSLER 92	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
1220 \pm 40	ALDE 86D	GAM4	$100 \pi^- p \rightarrow n2\eta$

¹ Breit-Wigner mass. May also be the $f_0(1500)$.**COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
1306 \pm 20	¹ ANISOVICH 03	RVUE
¹ K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.		

 $f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID
200 to 500 OUR ESTIMATE	

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
300 ± 80		¹ AUBERT	09L	BABR $B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
90 ^{+ 2} _{- 1} ^{+ 50} _{- 22}		² UEHARA	08A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
298 ± 21	2.6k	BONVICINI	07	CLEO $D^+ \rightarrow \pi^- \pi^+ \pi^+$
126 ± 25	4286	³ GARMASH	06	BELL $B^+ \rightarrow K^+ \pi^+ \pi^-$
265 ± 40		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$
350 ± 100 ^{+ 105} _{- 60}		ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
173 ± 32 ± 6	848	AITALA	01A	E791 $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
222 ± 20		BARBERIS	99B	OMEG 450 $p p \rightarrow p_S p_F \pi^+ \pi^-$
255 ± 60		BELLAZZINI	99	GAM4 450 $p p \rightarrow p p \pi^0 \pi^0$
190 ± 50		ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
323 ± 13		BERTIN	98	OBLX 0.05–0.405 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
350		^{4,5} TORNQVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195 ± 33		ARMSTRONG	91	OMEG 300 $p p \rightarrow p p \pi\pi, p p K\bar{K}$
285 ± 60		BREAKSTONE	90	SFM 62 $p p \rightarrow p p \pi^+ \pi^-$
460 ± 50		AKESSON	86	SPEC 63 $p p \rightarrow p p \pi^+ \pi^-$
~ 400		⁶ FROGGATT	77	RVUE $\pi^+ \pi^-$ channel

¹ The systematic errors are not reported.² Breit-Wigner width. May also be the $f_0(1500)$.³ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.⁴ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.⁵ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays⁶ Width defined as distance between 45 and 135° phase shift. **$K\bar{K}$ MODE**

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
121 ± 15		VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
55 ± 26		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
250 ± 80		BOLONKin	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
118 ^{+ 138} _{- 16}		ETKIN	82B	MPS 23 $\pi^- p \rightarrow n 2 K_S^0$
160 ± 30		WICKLUND	80	SPEC 6 $\pi N \rightarrow K^+ K^- N$
~ 150		POLYCHRO...	79	STRC 7 $\pi^- p \rightarrow n 2 K_S^0$

 4π MODE 2($\pi\pi$) $s+\rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
275 ± 55		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 ± 61		AMSLER	94	CBAR $0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
398 ± 26		ADAMO	93	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 ± 50		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 90	5751	¹ BETTINI	66	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant.

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
484 ⁺²⁴⁶ ₋₁₇₀ ⁺²⁴⁶ ₋₂₆₃	¹ UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
250	AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
320 \pm 40	ALDE	86D GAM4	$100 \pi^- p \rightarrow n2\eta$

¹ Breit-Wigner width. May also be the $f_0(1500)$.

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
147 ⁺³⁰ ₋₅₀	¹ ANISOVICH	03 RVUE
¹ K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.		

 $f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 4\pi$	seen
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 2\pi^+ 2\pi^-$	seen
$\Gamma_5 \pi^+ \pi^- 2\pi^0$	seen
$\Gamma_6 \rho\rho$	dominant
$\Gamma_7 2(\pi\pi)_S$ -wave	seen
$\Gamma_8 \pi(1300)\pi$	seen
$\Gamma_9 a_1(1260)\pi$	seen
$\Gamma_{10} \eta\eta$	seen
$\Gamma_{11} K\bar{K}$	seen
$\Gamma_{12} K\bar{K}n\pi$	not seen
$\Gamma_{13} 6\pi$	not seen
$\Gamma_{14} \omega\omega$	not seen
$\Gamma_{15} \gamma\gamma$	seen
$\Gamma_{16} e^+ e^-$	not seen

 $f_0(1370)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$**

See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90.

 Γ_{15} **$\Gamma(e^+ e^-)$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBIEV	88 ND	$e^+ e^- \rightarrow \pi^0 \pi^0$

 Γ_{16}

$f_0(1370) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{10}\Gamma_{15}/\Gamma$		
<u>VALUE</u> (eV)	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$121^{+133+169}_{-53-106}$	¹ UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
¹ Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.			

$f_0(1370)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_1/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.26 ± 0.09	BUGG	96	RVUE
<0.15	¹ AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
<0.06	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
¹ Using AMSLER 95B ($3\pi^0$).			

$\Gamma(4\pi)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
>0.72	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$

$\Gamma(4\pi^0)/\Gamma(4\pi)$	Γ_3/Γ_2		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
seen	ABELE	96	CBAR $0.0 \bar{p}p \rightarrow 5\pi^0$
0.068 ± 0.005	¹ GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
¹ Model-dependent evaluation.			

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(4\pi)$	$\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3 + \Gamma_4 + \Gamma_5)$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.420 ± 0.014	¹ GASPERO	93	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
¹ Model-dependent evaluation.			

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(4\pi)$	$\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3 + \Gamma_4 + \Gamma_5)$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.512 ± 0.019	¹ GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
¹ Model-dependent evaluation.			

$\Gamma(\rho\rho)/\Gamma(4\pi)$	Γ_6/Γ_2		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.26 ± 0.07	ABELE	01B CBAR	$0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(2(\pi\pi)s\text{-wave})/\Gamma(\pi\pi)$ Γ_7/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.6 ± 2.6	¹ ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$

¹ From the combined data of ABELE 96 and ABELE 96C.

$\Gamma(2(\pi\pi)s\text{-wave})/\Gamma(4\pi)$ Γ_7/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.51 ± 0.09	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)s\text{-wave})$ Γ_6/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
large	BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_s$
1.6 ± 0.2	AMSLER	94	$\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
~ 0.65	GASPERO	93	$0.0 \bar{p}n \rightarrow \text{hadrons}$

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_8/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.17 ± 0.06	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_9/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.06 ± 0.02	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(\eta\eta)/\Gamma(4\pi)$ $\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(28 \pm 11) \times 10^{-3}$	¹ ANISOVICH	02D	SPEC Combined fit
$(4.7 \pm 2.0) \times 10^{-3}$	BARBERIS	00E	$450 pp \rightarrow p_f \eta\eta p_s$

¹ From a combined K-matrix analysis of Crystal Barrel ($0.0 \bar{p}\bar{p} \rightarrow \pi^0 \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ± 0.13	BUGG	96	RVUE

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_{11}/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.08 ± 0.08	ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-, \phi K^+ K^-$
0.91 ± 0.20	¹ BARGIOTTI	03	OBLX $\bar{p}p$
0.12 ± 0.06	² ANISOVICH	02D	SPEC Combined fit
$0.46 \pm 0.15 \pm 0.11$	BARBERIS	99D	OMEG $450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$

¹ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta \eta n$, $\eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.03	GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons

$\Gamma(6\pi)/\Gamma_{\text{total}}$

Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.22	GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.13	GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons

$f_0(1370)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>	
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>	
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168 481.		
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)

ABELE	96	PL B380	453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385	425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609	562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	96	NP B471	59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342	433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353	571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355	425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
JANSSEN	95	PR D52	2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI) (HELS)
TORNQVIST	95	ZPHY C68	647	N.A. Tornqvist	
AMSLER	94	PL B322	431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC
AMSLER	94D	PL B333	277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323	233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC
BUGG	94	PR D50	4412	D.V. Bugg <i>et al.</i>	(LOQM)
KAMINSKI	94	PR D50	3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)
ADAMO	93	NP A558	13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC
GASPERO	93	NP A562	407	M. Gaspero	(ROMAI) JPC
AMSLER	92	PL B291	347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ARMSTRONG	91	ZPHY C51	351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ARMSTRONG	91B	ZPHY C52	389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BREAKSTONE	90	ZPHY C48	569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MORGAN	90	ZPHY C48	623	D. Morgan, M.R. Pennington	(RAL, DURH)
ASTON	88	NP B296	493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BOLONKIN	88	NP B309	426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38	2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAZO+)
VOROBYEV	88	SJNP 48	273	P.V. Vorobiev <i>et al.</i>	(NOVO)
Translated from YAF 48 436.					
AU	87	PR D35	1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
AKESSON	86	NP B264	154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
ALDE	86D	NP B269	485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
CASON	83	PR D28	1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25	1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
WICKLUND	80	PRL 45	1469	A.B. Wicklund <i>et al.</i>	(ANL)
BECKER	79	NP B151	46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
POLYCHRO...	79	PR D19	1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
FROGGATT	77	NP B129	89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ROSSELET	77	PR D15	574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
GRAYER	74	NP B75	189	G. Grayer <i>et al.</i>	(CERN, MPIM)
HYAMS	73	NP B64	134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis		W. Ochs	(MPIM, MUNI)
BEIER	72B	PRL 29	511	E.W. Beier <i>et al.</i>	(PENN)
BETTINI	66	NC 42A	695	A. Bettini <i>et al.</i>	(PADO, PISA)