

$f_0(1710)$ $I^G(J^{PC}) = 0^+(0^{++})$

See our mini-review in the 2004 edition of this *Review*, Physics Letters **B592** 1 (2004). See also the mini-review on scalar mesons under $f_0(500)$ (see the index for the page number).

 $f_0(1710)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1723^{+6}_{-5} OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
1759 ± 6	$+_{-25}^{14}$	5.5k	¹ ABLIKIM	$13N$ BES3 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1750 ± 6	$+_{-18}^{29}$		UEHARA	13 BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
1701 ± 5	$+_{-2}^{9}$	4k	² CHEKANOV	08 ZEUS $e p \rightarrow K_S^0 K_S^0 X$
1765 ± 4	± 13		ABLIKIM	$06V$ BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1760 ± 15	$+_{-10}^{15}$		³ ABLIKIM	$05Q$ BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^- K^+ K^-$
1738 ± 30			ABLIKIM	$04E$ BES2 $J/\psi \rightarrow \omega K^+ K^-$
1740 ± 4	$+_{-25}^{10}$		⁴ BAI	$03G$ BES $J/\psi \rightarrow \gamma K\bar{K}$
1740	$+_{-25}^{30}$		⁴ BAI	$00A$ BES $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1698 ± 18			⁵ BARBERIS	$00E$ $450 pp \rightarrow p_f \eta\eta p_s$
1710 ± 12	± 11		⁶ BARBERIS	$99D$ OMEG $450 pp \rightarrow K^+ K^-, \pi^+\pi^-$
1710 ± 25			⁷ FRENCH	99 $300 pp \rightarrow p_f (K^+ K^-) p_s$
1707 ± 10			⁸ AUGUSTIN	88 DM2 $J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0 X$
1698 ± 15			⁸ AUGUSTIN	87 DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
1720 ± 10	± 10		⁹ BALTRUSAIT..87	$MRK3$ $J/\psi \rightarrow \gamma K^+ K^-$
1742 ± 15			⁸ WILLIAMS	84 MPSF $200 \pi^- N \rightarrow 2K_S^0 X$
1670 ± 50			BLOOM	83 CBAL $J/\psi \rightarrow \gamma 2\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1744 ± 7	± 5	381 ^{10,11}	DOBBS	15 $J/\psi \rightarrow \gamma\pi^+\pi^-$
1705 ± 11	± 5	237 ^{10,11}	DOBBS	15 $\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1706 ± 4	± 5	1.0k ^{10,11}	DOBBS	15 $J/\psi \rightarrow \gamma K^+ K^-$
1690 ± 8	± 3	349 ^{10,11}	DOBBS	15 $\psi(2S) \rightarrow \gamma K^+ K^-$
1750 ± 13			AMSLER	06 CBAR $1.64 \bar{p}p \rightarrow K^+ K^- \pi^0$
1747 ± 5		80k ^{12,13}	UMAN	06 E835 $5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
1776 ± 15			VLADIMIRSK..06	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1790	$+_{-30}^{40}$		³ ABLIKIM	05 BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$
1670 ± 20			¹² BINON	05 GAMS $33 \pi^- p \rightarrow \eta\eta n$
1726 ± 7		74	¹³ CHEKANOV	04 ZEUS $e p \rightarrow K_S^0 K_S^0 X$
1732 ± 15			¹⁴ ANISOVICH	03 RVUE
1682 ± 16			TIKHOMIROV	03 SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1670 ± 26		3.6k ^{4,15}	NICHITU	02 OBLX
1770 ± 12		16,17	ANISOVICH	$99B$ SPEC $0.6-1.2 p\bar{p} \rightarrow \eta\eta\pi^0$

1730±15		⁴ BARBERIS	99	OMEG	450 $p p \rightarrow p_s p_f K^+ K^-$
1750±20		⁴ BARBERIS	99B	OMEG	450 $p p \rightarrow p_s p_f \pi^+ \pi^-$
1750±30		¹⁸ ANISOVICH	98B	RVUE	Compilation
1720±39		BAI	98H	BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
1775± 1.5	57	¹⁹ BARKOV	98		$\pi^- p \rightarrow K_S^0 K_S^0 n$
1690±11		20 ABREU	96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$
1696± 5	⁺⁹ -34	9 BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
1781± 8	⁺¹⁰ -31	4 BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
1768±14		BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
1750±15		²¹ BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1620±16		9 BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1748±10		⁸ ARMSTRONG	93C	E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
~1750		BREAKSTONE	93	SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
1744±15		22 ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta \eta n$
1713±10		23 ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K^+ K^-$
1706±10		23 ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K_S^0 K_S^0$
1700±15		9 BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1720±60		4 BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1638±10		24 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
1690± 4		25 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
1755± 8		26 ALDE	86C	GAM2	$38 \pi^- p \rightarrow n 2\eta$
1730 ⁺² -10		27 LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n 2 K_S^0$
1650±50		BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
1640±50	28,29	EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$
1730±10 ±20		30 ETKIN	82C	MPS	$23 \pi^- p \rightarrow n 2 K_S^0$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

³ This state may be different from $f_0(1710)$, see CLOSE 05.

⁴ $J^P = 0^+$.

⁵ T-matrix pole.

⁶ Supersedes BARBERIS 99 and BARBERIS 99B.

⁷ $J^P = 0^+$, supersedes by ARMSTRONG 89D.

⁸ No $J^P C$ determination.

⁹ $J^P = 2^+$.

¹⁰ Using CLEO-c data but not authored by the CLEO Collaboration.

¹¹ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 135$ MeV.

¹² Breit-Wigner mass.

¹³ Systematic errors not estimated.

¹⁴ K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.


¹⁵ Decaying to $f_0(1370) \pi \pi$.

¹⁶ $J^P = 0^+$.

¹⁷ Not seen by AMSLER 02.

¹⁸ T-matrix pole, assuming $J^P = 0^+$

- 19 No $J^P C$ determination.
- 20 No $J^P C$ determination, width not determined.
- 21 From a fit to the 0^+ partial wave.
- 22 ALDE 92D combines all the GAMS-2000 data.
- 23 $J^P = 2^+$, superseded by FRENCH 99.
- 24 From an analysis ignoring interference with $f'_2(1525)$.
- 25 From an analysis including interference with $f'_2(1525)$.
- 26 Superseded by ALDE 92D.
- 27 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
- 28 $J^P = 2^+$ preferred.
- 29 From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
- 30 Superseded by LONGACRE 86.



$f_0(1710)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
139 ± 8 OUR AVERAGE		Error includes scale factor of 1.1.		
172 ± 10	+32 -16	5.5k	1 ABLIKIM	13N BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
139	+ 11 - 12	+ 96 - 50	UEHARA	13 BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
100 ± 24	+ 7 - 22	4k	2 CHEKANOV	08 ZEUS $e p \rightarrow K_S^0 K_S^0 X$
145 ± 8	± 69		ABLIKIM	06V BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
125 ± 25	+ 10 - 15		3 ABLIKIM	05Q BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^- K^+ K^-$
125 ± 20			ABLIKIM	04E BES2 $J/\psi \rightarrow \omega K^+ K^-$

166	\pm	5	$+15$		4 BAI	03G	BES	$J/\psi \rightarrow \gamma K\bar{K}$
120	\pm	50			4 BAI	00A	BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
120	\pm	26			5 BARBERIS	00E		$450 pp \rightarrow p_f \eta \eta p_s$
126	\pm	16	± 18		6 BARBERIS	99D	OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$
105	\pm	34			7 FRENCH	99		$300 pp \rightarrow p_f(K^+ K^-)p_s$
166.4 \pm 33.2					8 AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0 X$
136	\pm	28			8 AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
130	\pm	20			9 BALTRUSAIT	..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
57	\pm	38			10 WILLIAMS	84	MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
160	\pm	80			BLOOM	83	CBAL	$J/\psi \rightarrow \gamma 2\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •								
148	\pm	40			AMSLER	06	CBAR	$1.64 \bar{p}p \rightarrow K^+ K^- \pi^0$
188	\pm	13		80k	3,11 UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
250	\pm	30			VLADIMIRSK	..06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
270	\pm	60			12 ABLIKIM	05	BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
260	\pm	50			3 BINON	05	GAMS	$33 \pi^- p \rightarrow \eta \eta n$
38	\pm	20		74	11 CHEKANOV	04	ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
144	\pm	30			13,14 ANISOVICH	03	RVUE	
320	\pm	50			14,15 ANISOVICH	03	RVUE	
102	\pm	26			TIKHOMIROV	03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
267	\pm	44		3651	4,16 NICHTIU	02	OBLX	
220	\pm	40			17,18 ANISOVICH	99B	SPEC	$0.6-1.2 p\bar{p} \rightarrow \eta \eta \pi^0$
100	\pm	25			4 BARBERIS	99	OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$
160	\pm	30			4 BARBERIS	99B	OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
250	\pm	140			19 ANISOVICH	98B	RVUE	Compilation
30	\pm	7		57	20 BARKOV	98		$\pi^- p \rightarrow K_S^0 K_S^0 n$
103	\pm	18	$+30$		9 BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
85	\pm	24	$+22$		4 BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
56	\pm	19			BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
160	\pm	40			21 BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
160	\pm	60			9 BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
264	\pm	25			8 ARMSTRONG	93C	E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
200	to	300			BREAKSTONE	93	SFM	$pp \rightarrow pp\pi^+ \pi^- \pi^+ \pi^-$
< 80	90% CL				22 ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta \eta N^*$
181	\pm	30			23 ARMSTRONG	89D	OMEG	$300 pp \rightarrow pp K^+ K^-$
104	\pm	30			23 ARMSTRONG	89D	OMEG	$300 pp \rightarrow pp K_S^0 K_S^0$
30	\pm	20			9 BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
350	\pm	150			4 BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
148	\pm	17			24 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
184	\pm	6			25 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
122	\pm	74			26 LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n2K_S^0$

200	± 100	BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
220	$+100$ -70	27,28 EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$
200	$+156$ -9	29 ETKIN	82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

³ Breit-Wigner width.

⁴ $J^P = 0^+$.

⁵ T-matrix pole.

⁶ Supersedes BARBERIS 99 and BARBERIS 99B.

⁷ $J^P = 0^+$, supersedes by ARMSTRONG 89D.

⁸ No $J^P C$ determination.

⁹ $J^P = 2^+$.

¹⁰ No $J^P C$ determination.

¹¹ Systematic errors not estimated.

¹² This state may be different from $f_0(1710)$, see CLOSE 05.

¹³ (Solution I)

¹⁴ K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

¹⁵ (Solution I)

¹⁶ Decaying to $f_0(1370) \pi \pi$.

¹⁷ $J^P = 0^+$.

¹⁸ Not seen by AMSLER 02.

¹⁹ T-matrix pole, assuming $J^P = 0^+$

²⁰ No $J^P C$ determination.

²¹ From a fit to the 0^+ partial wave.

²² ALDE 92D combines all the GAMS-2000 data.

²³ $J^P = 2^+$, (0^+ excluded).

²⁴ From an analysis ignoring interference with $f'_2(1525)$.

²⁵ From an analysis including interference with $f'_2(1525)$.

²⁶ Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

²⁷ $J^P = 2^+$ preferred.

²⁸ From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.

²⁹ From an amplitude analysis of the $K_S^0 K_S^0$ system, superseded by LONGACRE 86.

$f_0(1710)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}$	seen
Γ_2 $\eta \eta$	seen
Γ_3 $\pi \pi$	seen
Γ_4 $\gamma \gamma$	
Γ_5 $\omega \omega$	seen

$f_0(1710)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_4/\Gamma$

<u>VALUE</u> (eV)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12⁺³₋₂⁺²²⁷₋₈		UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<480	95	ALBRECHT	90G	ARG $\gamma\gamma \rightarrow K^+ K^-$
<110	95	¹ BEHREND	89C	CELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
<280	95	¹ ALTHOFF	85B	TASS $\gamma\gamma \rightarrow K K\pi$

¹ Assuming helicity 2.

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_4/\Gamma$

<u>VALUE</u> (keV)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.82	95	¹ BARATE	00E	ALEP $\gamma\gamma \rightarrow \pi^+ \pi^-$

¹ Assuming spin 0.

 $f_0(1710)$ BRANCHING RATIOS $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_1/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	1004	¹ DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
seen	349	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
0.36 ± 0.12		ALBALADEJO 08	RVUE	
$0.38^{+0.09}_{-0.19}$		² LONGACRE	86	MPS $22 \pi^- p \rightarrow n 2 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
0.22 ± 0.12	ALBALADEJO 08	RVUE
$0.18^{+0.03}_{-0.13}$	¹ LONGACRE	86 RVUE

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	381	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
seen	237	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
not seen		AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$
$0.039^{+0.002}_{-0.024}$		² LONGACRE	86	RVUE

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_1
0.41^{+0.11}_{-0.17}		ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.32 \pm 0.14		ALBALADEJO	08 RVUE		
< 0.11	95	¹ ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$	
5.8 ^{+9.1} _{-5.5}		² ANISOVICH	02D SPEC	Combined fit	
0.2 \pm 0.024 \pm 0.036		BARBERIS	99D OMEG	$450 pp \rightarrow K^+K^-, \pi^+\pi^-$	
0.39 \pm 0.14		ARMSTRONG	91 OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$	

¹ Using data from ABLIKIM 04A.² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data. $\Gamma(\eta\eta)/\Gamma(K\bar{K})$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ_1
0.48\pm0.15		BARBERIS	00E	$450 pp \rightarrow p_f\eta\eta p_s$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.46 ^{+0.70} _{-0.38}		¹ ANISOVICH	02D SPEC	Combined fit	
<0.02	90	² PROKOSHKIN	91 GA24	$300 \pi^-p \rightarrow \pi^-p\eta\eta$	
¹ From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.					
² Combining results of GAM4 with those of ARMSTRONG 89D.					

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ
seen	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$	

f₀(1710) REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
ALBALADEJO	08	PRL 101 252002	M. Albaladejo, J.A. Oller	
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSKY	06	PAN 69 493	V.V. Vladimirskey <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
CLOSE	05	PR D71 094022	F.E. Close, Q. Zhao	
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		

AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>
		Translated from YAF 65 1583.	
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>
FRENCH	99	PL B460 213	B. French <i>et al.</i>
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>
		Translated from UFN 168 481.	
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>
		Translated from YAF 58 50.	
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>
Also		SJNP 54 451	D.M. Alde <i>et al.</i>
		Translated from YAF 54 745.	
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin
		Translated from DANS 316 900.	
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>
