

**$f_0(980)$**  $I^G(J^{PC}) = 0^+(0^{++})$ 

See also the minireview on scalar mesons under  $f_0(500)$ . (See the index for the page number.)

### **$f_0(980)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>990 ± 20 OUR ESTIMATE</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
989.4 ± 1.3	424	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
989.9 ± 0.4	706	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma 3\pi$
1003 $\begin{array}{l} +5 \\ -27 \end{array}$	1,2 GARCIA-MAR..11	RVUE	Compilation	
996 $\pm 7$	1,3 GARCIA-MAR..11	RVUE	Compilation	
996 $\begin{array}{l} +4 \\ -14 \end{array}$	4 MOUSSALLAM11	RVUE	Compilation	
981 $\pm 43$	5 MENNESSIER 10	RVUE	Compilation	
1030 $\begin{array}{l} +30 \\ -10 \end{array}$	6 ANISOVICH 09	RVUE	0.0 $\bar{p}p, \pi N$	
977 $\begin{array}{l} +11 \\ -9 \end{array} \pm 1$	44	7 ECKLUND 09	CLEO	$4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + c.c.$
982.2 ± 1.0 $\begin{array}{l} +8.1 \\ -8.0 \end{array}$	8 UEHARA 08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
976.8 ± 0.3 $\begin{array}{l} +10.1 \\ -0.6 \end{array}$	64k	9 AMBROSINO 07	KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
984.7 ± 0.4 $\begin{array}{l} +2.4 \\ -3.7 \end{array}$	64k	10 AMBROSINO 07	KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
973 $\pm 3$	262 ± 30	11 AUBERT 07AKBABR	10.6 $e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$	
970 $\pm 7$	54 ± 9	11 AUBERT 07AKBABR	10.6 $e^+ e^- \rightarrow \phi \pi^0 \pi^0 \gamma$	
953 $\pm 20$	2.6k	12 BONVICINI 07	CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
985.6 $\begin{array}{l} +1.2 \\ -1.5 \end{array} \begin{array}{l} +1.1 \\ -1.6 \end{array}$	13 MORI 07	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
983.0 ± 0.6 $\begin{array}{l} +4.0 \\ -3.0 \end{array}$	14 AMBROSINO 06B	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
977.3 ± 0.9 $\begin{array}{l} +3.7 \\ -4.3 \end{array}$	15 AMBROSINO 06B	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
950 $\pm 9$	4286	16 GARMASH 06	BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
965 $\pm 10$		17 ABLIKIM 05	BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-, \phi K^+ K^-$
1031 $\pm 8$		18 ANISOVICH 03	RVUE	
1037 $\pm 31$		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
973 $\pm 1$	2438	19 ALOISIO 02D	KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
977 $\pm 3 \pm 2$	848	20 AITALA 01A	E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
969.8 ± 4.5	419	21 ACHASOV 00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
985 $\begin{array}{l} +16 \\ -12 \end{array}$	419	22,23 ACHASOV 00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
976 $\pm 5 \pm 6$		24 AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

977 $\pm$ 3 $\pm$ 6	268	24 AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
975 $\pm$ 4 $\pm$ 6		25 AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
975 $\pm$ 4 $\pm$ 6		26 AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma,$ $\pi^0 \pi^0 \gamma$
985 $\pm$ 10		BARBERIS	99 OMEG	450 $p p \rightarrow p_s p_f K^+ K^-$
982 $\pm$ 3		BARBERIS	99B OMEG	450 $p p \rightarrow p_s p_f \pi^+ \pi^-$
982 $\pm$ 3		BARBERIS	99C OMEG	450 $p p \rightarrow p_s p_f \pi^0 \pi^0$
987 $\pm$ 6 $\pm$ 6		27 BARBERIS	99D OMEG	450 $p p \rightarrow K^+ K^-,$ $\pi^+ \pi^-$
989 $\pm$ 15		BELLAZZINI	99 GAM4	450 $p p \rightarrow p p \pi^0 \pi^0$
991 $\pm$ 3		28 KAMINSKI	99 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}, \sigma \sigma$
$\sim$ 980		28 OLLER	99 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}$
$\sim$ 993.5		OLLER	99B RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}$
$\sim$ 987		28 OLLER	99C RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}, \eta \eta$
957 $\pm$ 6		29 ACKERSTAFF	98Q OPAL	$Z \rightarrow f_0 X$
960 $\pm$ 10		ALDE	98 GAM4	
1015 $\pm$ 15		28 ANISOVICH	98B RVUE	Compilation
1008		30 LOCHER	98 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}$
955 $\pm$ 10		29 ALDE	97 GAM2	450 $p p \rightarrow p p \pi^0 \pi^0$
994 $\pm$ 9		31 BERTIN	97C OBLX	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
993.2 $\pm$ 6.5 $\pm$ 6.9		32 ISHIDA	96 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}$
1006		TORNQVIST	96 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}, K\pi,$ $\eta \pi$
997 $\pm$ 5	3k	33 ALDE	95B GAM2	38 $\pi^- p \rightarrow \pi^0 \pi^0 n$
960 $\pm$ 10	10k	34 ALDE	95B GAM2	38 $\pi^- p \rightarrow \pi^0 \pi^0 n$
994 $\pm$ 5		AMSLER	95B CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$
$\sim$ 996		35 AMSLER	95D CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$
987 $\pm$ 6		36 ANISOVICH	95 RVUE	
1015		JANSSEN	95 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}$
983		37 BUGG	94 RVUE	$\bar{p} p \rightarrow \eta 2\pi^0$
973 $\pm$ 2		38 KAMINSKI	94 RVUE	$\pi \pi \rightarrow \pi \pi, K\bar{K}$
988		39 ZOU	94B RVUE	
988 $\pm$ 10		40 MORGAN	93 RVUE	$\pi \pi (K\bar{K}) \rightarrow \pi \pi (K\bar{K}),$ $J/\psi \rightarrow \phi \pi \pi (K\bar{K}),$ $D_s \rightarrow \pi (\pi \pi)$
971.1 $\pm$ 4.0		29 AGUILAR-...	91 EHS	400 $p p$
979 $\pm$ 4		41 ARMSTRONG	91 OMEG	300 $p p \rightarrow p p \pi \pi,$ $p p K\bar{K}$
956 $\pm$ 12		BREAKSTONE	90 SFM	$p p \rightarrow p p \pi^+ \pi^-$
959.4 $\pm$ 6.5		29 AUGUSTIN	89 DM2	$J/\psi \rightarrow \omega \pi^+ \pi^-$
978 $\pm$ 9		29 ABACHI	86B HRS	$e^+ e^- \rightarrow \pi^+ \pi^- X$
985.0 $\pm$ 9.0 $-39.0$		ETKIN	82B MPS	23 $\pi^- p \rightarrow n 2K_S^0$
974 $\pm$ 4		41 GIDAL	81 MRK2	$J/\psi \rightarrow \pi^+ \pi^- X$
975		42 ACHASOV	80 RVUE	
986 $\pm$ 10		41 AGUILAR-...	78 HBC	0.7 $\bar{p} p \rightarrow K_S^0 K_S^0$
969 $\pm$ 5		41 LEEPER	77 ASPK	2–2.4 $\pi^- p \rightarrow$ $\pi^+ \pi^- n, K^+ K^- n$
987 $\pm$ 7		41 BINNIE	73 CNTR	$\pi^- p \rightarrow n MM$

1012 $\pm$ 6	<sup>43</sup> GRAYER	73	ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1007 $\pm$ 20	<sup>43</sup> HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
997 $\pm$ 6	<sup>43</sup> PROTOPOP...	73	HBC	7 $\pi^+ p \rightarrow \pi^+ p \pi^+ \pi^-$

<sup>1</sup> Quoted number refers to real part of pole position.

<sup>2</sup> Analytic continuation using Roy equations. Uses the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.

<sup>3</sup> Analytic continuation using GKPY equations. Uses the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.

<sup>4</sup> Pole position. Used Roy equations.

<sup>5</sup> Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.

<sup>6</sup> On sheet II in a 2-pole solution. The other pole is found on sheet III at  $(850-100i)$  MeV

<sup>7</sup> Using a relativistic Breit-Wigner function and taking into account the finite  $D_S$  mass.

<sup>8</sup> Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K K / g_{f_0} \pi\pi = 0$ .

<sup>9</sup> In the kaon-loop fit.

<sup>10</sup> In the no-structure fit.

<sup>11</sup> Systematic errors not estimated.

<sup>12</sup> FLATTE 76 parameterization.  $g_{f_0} \pi\pi = 329 \pm 96$  MeV/c<sup>2</sup> assuming  $g_{f_0} K\bar{K} / g_{f_0} \pi\pi = 2$ .

<sup>13</sup> Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K K / g_{f_0} \pi\pi = 4.21 \pm 0.25 \pm 0.21$  from ABLIKIM 05.

<sup>14</sup> In the kaon-loop fit following formalism of ACHASOV 89.

<sup>15</sup> In the no-structure fit assuming a direct coupling of  $\phi$  to  $f_0 \gamma$ .

<sup>16</sup> FLATTE 76 parameterization. Supersedes GARMASH 05.

<sup>17</sup> FLATTE 76 parameterization,  $g_{f_0} K\bar{K} / g_{f_0} \pi\pi = 4.21 \pm 0.25 \pm 0.21$ .

<sup>18</sup> K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K\bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.

<sup>19</sup> From the negative interference with the  $f_0(500)$  meson of AITALA 01B using the ACHASOV 89 parameterization for the  $f_0(980)$ , a Breit-Wigner for the  $f_0(500)$ , and ACHASOV 01F for the  $\rho\pi$  contribution.

<sup>20</sup> Coupled-channel Breit-Wigner, couplings  $g_\pi = 0.09 \pm 0.01 \pm 0.01$ ,  $g_K = 0.02 \pm 0.04 \pm 0.03$ .

<sup>21</sup> Supersedes ACHASOV 98I. Using the model of ACHASOV 89.

<sup>22</sup> Supersedes ACHASOV 98I.

<sup>23</sup> In the “narrow resonance” approximation.

<sup>24</sup> Assuming  $\Gamma(f_0) = 40$  MeV.

<sup>25</sup> From a narrow pole fit taking into account  $f_0(980)$  and  $f_0(1200)$  intermediate mechanisms.

<sup>26</sup> From the combined fit of the photon spectra in the reactions  $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$ ,  $\pi^0 \pi^0 \gamma$ .

<sup>27</sup> Supersedes BARBERIS 99 and BARBERIS 99B

<sup>28</sup> T-matrix pole.

<sup>29</sup> From invariant mass fit.

<sup>30</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(1039-93i)$  MeV.

<sup>31</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(963-29i)$  MeV.

<sup>32</sup> Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

<sup>33</sup> At high  $|t|$ .

<sup>34</sup> At low  $|t|$ .

- 35 On sheet II in a 4-pole solution, the other poles are found on sheet III at  $(953 - 55i)$  MeV and on sheet IV at  $(938 - 35i)$  MeV.  
 36 Combined fit of ALDE 95B, ANISOVICH 94, AMSLER 94D.  
 37 On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(996 - 103i)$  MeV.  
 38 From sheet II pole position.  
 39 On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(797 - 185i)$  MeV and can be interpreted as a shadow pole.  
 40 On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(978 - 28i)$  MeV.  
 41 From coupled channel analysis.  
 42 Coupled channel analysis with finite width corrections.  
 43 Included in AGUILAR-BENITEZ 78 fit.
- 

## $f_0(980)$ WIDTH

Width determination very model dependent. Peak width in  $\pi\pi$  is about 50 MeV, but decay width can be much larger.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10 to 100 OUR ESTIMATE</b>				
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
15.3 $\pm$ 4.7	424	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
9.5 $\pm$ 1.1	706	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma 3\pi$
42 $\pm$ 20 — 16		1,2 GARCIA-MAR..11	RVUE	Compilation
50 $\pm$ 20 — 12		2,3 GARCIA-MAR..11	RVUE	Compilation
48 $\pm$ 22 — 6		4 MOUSSALLAM11	RVUE	Compilation
36 $\pm$ 22		5 MENNESSIER 10	RVUE	Compilation
70 $\pm$ 20 — 32		6 ANISOVICH 09	RVUE	0.0 $\bar{p}p$ , $\pi N$
91 $\pm$ 30 — 22 $\pm$ 3	44	7 ECKLUND	09 CLEO	$4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + c.c.$
66.9 $\pm$ 2.2 $\pm$ 17.6 — 12.5		8 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
65 $\pm$ 13	262 $\pm$ 30	9 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
81 $\pm$ 21	54 $\pm$ 9	9 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \pi^0 \gamma$
51.3 $\pm$ 20.8 $\pm$ 13.2 — 17.7 — 3.8		10 MORI	07 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
61 $\pm$ 9 $\pm$ 14 — 8	2584	11 GARMASH	05 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
64 $\pm$ 16		12 ANISOVICH	03 RVUE	
121 $\pm$ 23		TIKHOMIROV 03	SPEC	$40.0 \frac{\pi^-}{K_S^0} \frac{C}{K_S^0} \rightarrow \frac{K_L^0}{K_S^0} \frac{K^0}{K_S^0} X$
$\sim$ 70		13 BRAMON	02 RVUE	$1.02 \frac{e^+ e^-}{\pi^0 \pi^0 \gamma} \rightarrow \frac{\pi^0}{\pi^0} \frac{\pi^0}{\pi^0} \gamma$
44 $\pm$ 2 $\pm$ 2	848	14 AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
201 $\pm$ 28	419	15 ACHASOV	00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
122 $\pm$ 13	419	16,17 ACHASOV	00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
56 $\pm$ 20		18 AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

65 $\pm$ 20		BARBERIS	99	OMEG	450 $pp \rightarrow p_s p_f K^+ K^-$
80 $\pm$ 10		BARBERIS	99B	OMEG	450 $pp \rightarrow p_s p_f \pi^+ \pi^-$
80 $\pm$ 10		BARBERIS	99C	OMEG	450 $pp \rightarrow p_s p_f \pi^0 \pi^0$
48 $\pm$ 12 $\pm$ 8		19 BARBERIS	99D	OMEG	450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$
65 $\pm$ 25		BELLAZZINI	99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
71 $\pm$ 14		20 KAMINSKI	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$\sim$ 28		20 OLLER	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
$\sim$ 25		OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
$\sim$ 14		20 OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
70 $\pm$ 20		ALDE	98	GAM4	
86 $\pm$ 16		20 ANISOVICH	98B	RVUE	Compilation
54		21 LOCHER	98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
69 $\pm$ 15		22 ALDE	97	GAM2	450 $pp \rightarrow pp\pi^0\pi^0$
38 $\pm$ 20		23 BERTIN	97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
$\sim$ 100		24 ISHIDA	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
34		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
48 $\pm$ 10	3k	25 ALDE	95B	GAM2	38 $\pi^- p \rightarrow \pi^0\pi^0 n$
95 $\pm$ 20	10k	26 ALDE	95B	GAM2	38 $\pi^- p \rightarrow \pi^0\pi^0 n$
26 $\pm$ 10		AMSLER	95B	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$
$\sim$ 112		27 AMSLER	95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$
80 $\pm$ 12		28 ANISOVICH	95	RVUE	
30		JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
74		29 BUGG	94	RVUE	$\bar{p}p \rightarrow \eta 2\pi^0$
29 $\pm$ 2		30 KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
46		31 ZOU	94B	RVUE	
48 $\pm$ 12		32 MORGAN	93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_s \rightarrow \pi(\pi\pi)$
37.4 $\pm$ 10.6		22 AGUILAR-...	91	EHS	400 $pp$
72 $\pm$ 8		33 ARMSTRONG	91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$
110 $\pm$ 30		BREAKSTONE	90	SFM	$pp \rightarrow pp\pi^+\pi^-$
29 $\pm$ 13		22 ABACHI	86B	HRS	$e^+ e^- \rightarrow \pi^+\pi^- X$
120 $\pm$ 281 $\pm$ 20		ETKIN	82B	MPS	$23 \pi^- p \rightarrow n 2 K_S^0$
28 $\pm$ 10		33 GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+\pi^- X$
70 to 300		34 ACHASOV	80	RVUE	
100 $\pm$ 80		35 AGUILAR-...	78	HBC	$0.7 \bar{p}p \rightarrow K_S^0 K_S^0$
30 $\pm$ 8		33 LEEPER	77	ASPK	$2-2.4 \pi^- p \rightarrow \pi^+\pi^- n, K^+ K^- n$
48 $\pm$ 14		33 BINNIE	73	CNTR	$\pi^- p \rightarrow n MM$
32 $\pm$ 10		36 GRAYER	73	ASPK	$17 \pi^- p \rightarrow \pi^+\pi^- n$
30 $\pm$ 10		36 HYAMS	73	ASPK	$17 \pi^- p \rightarrow \pi^+\pi^- n$
54 $\pm$ 16		36 PROTOPOP...	73	HBC	$7 \pi^+ p \rightarrow \pi^+\pi^+\pi^-$

- <sup>1</sup> Analytic continuation using Roy equations. Uses the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.
- <sup>2</sup> Quoted number refers to twice imaginary part of pole position.
- <sup>3</sup> Analytic continuation using GKPY equations. Uses the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.
- <sup>4</sup> Pole position. Used Roy equations.
- <sup>5</sup> Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.
- <sup>6</sup> On sheet II in a 2-pole solution. The other pole is found on sheet III at  $(850-100i)$  MeV
- <sup>7</sup> Using a relativistic Breit-Wigner function and taking into account the finite  $D_S$  mass.
- <sup>8</sup> Breit-Wigner  $\pi\pi$  width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K K / g_{f_0} \pi\pi = 0$ .
- <sup>9</sup> Systematic errors not estimated.
- <sup>10</sup> Breit-Wigner  $\pi\pi$  width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K K / g_{f_0} \pi\pi = 4.21 \pm 0.25 \pm 0.21$  from ABLIKIM 05.
- <sup>11</sup> Breit-Wigner, solution 1, PWA ambiguous.
- <sup>12</sup> K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.
- <sup>13</sup> Using the data of AKHMETSHIN 99C, ACHASOV 00H, and ALOISIO 02D.
- <sup>14</sup> Breit-Wigner width.
- <sup>15</sup> Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
- <sup>16</sup> Supersedes ACHASOV 98I.
- <sup>17</sup> In the “narrow resonance” approximation.
- <sup>18</sup> From the combined fit of the photon spectra in the reactions  $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$ ,  $\pi^0 \pi^0 \gamma$ .
- <sup>19</sup> Supersedes BARBERIS 99 and BARBERIS 99B
- <sup>20</sup> T-matrix pole.
- <sup>21</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(1039-93i)$  MeV.
- <sup>22</sup> From invariant mass fit.
- <sup>23</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(963-29i)$  MeV.
- <sup>24</sup> Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- <sup>25</sup> At high  $|t|$ .
- <sup>26</sup> At low  $|t|$ .
- <sup>27</sup> On sheet II in a 4-pole solution, the other poles are found on sheet III at  $(953-55i)$  MeV and on sheet IV at  $(938-35i)$  MeV.
- <sup>28</sup> Combined fit of ALDE 95B, ANISOVICH 94,
- <sup>29</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(996-103i)$  MeV.
- <sup>30</sup> From sheet II pole position.
- <sup>31</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(797-185i)$  MeV and can be interpreted as a shadow pole.
- <sup>32</sup> On sheet II in a 2 pole solution. The other pole is found on sheet III at  $(978-28i)$  MeV.
- <sup>33</sup> From coupled channel analysis.
- <sup>34</sup> Coupled channel analysis with finite width corrections.
- <sup>35</sup> From coupled channel fit to the HYAMS 73 and PROTOPOPESCU 73 data. With a simultaneous fit to the  $\pi\pi$  phase-shifts, inelasticity and to the  $K_S^0 K_S^0$  invariant mass.
- <sup>36</sup> Included in AGUILAR-BENITEZ 78 fit.

**$f_0(980)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \pi\pi$	dominant
$\Gamma_2 K\bar{K}$	seen
$\Gamma_3 \gamma\gamma$	seen
$\Gamma_4 e^+e^-$	

 **$f_0(980)$  PARTIAL WIDTHS**

$\Gamma(\gamma\gamma)$			$\Gamma_3$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.31 <math>^{+0.05}_{-0.04}</math> OUR AVERAGE</b>			
0.32 $\pm 0.05$	<sup>1</sup> DAI	14A RVUE	Compilation
$0.286 \pm 0.017$ $^{+0.211}_{-0.070}$	<sup>2</sup> UEHARA	08A BELL	$10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
$0.205 \pm 0.095$ $^{+0.147}_{-0.083}$ $^{+0.147}_{-0.117}$	<sup>3</sup> MORI	07 BELL	$10.6 e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
0.42 $\pm 0.06$ $\pm 0.18$	<sup>4</sup> OEST	90 JADE	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.16 $\pm 0.01$	<sup>5</sup> MENNESSIER	11 RVUE	
0.29 $\pm 0.21$ $^{+0.02}_{-0.07}$	<sup>6</sup> MOUSSALLAM	11 RVUE	Compilation
0.42	<sup>7,8</sup> PENNINGTON	08 RVUE	Compilation
0.10	<sup>8,9</sup> PENNINGTON	08 RVUE	Compilation
0.28 $^{+0.09}_{-0.13}$	<sup>10</sup> BOGLIONE	99 RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$
0.29 $\pm 0.07$ $\pm 0.12$	<sup>11,12</sup> BOYER	90 MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
0.31 $\pm 0.14$ $\pm 0.09$	<sup>11,12</sup> MARSISKE	90 CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
0.63 $\pm 0.14$	<sup>13</sup> MORGAN	90 RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$

<sup>1</sup> Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.

<sup>2</sup> Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K\bar{K}/g_{f_0}\pi\pi = 0$ .

<sup>3</sup> Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K\bar{K}/g_{f_0}\pi\pi = 4.21 \pm 0.25 \pm 0.21$  from ABLIKIM 05.

<sup>4</sup> OEST 90 quote systematic errors  $^{+0.08}_{-0.18}$ . We use  $\pm 0.18$ . Observed 60 events.

<sup>5</sup> Uses an analytic K-matrix model. Compilation.

<sup>6</sup> Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.

<sup>7</sup> Solution A (preferred solution based on  $\chi^2$ -analysis).

<sup>8</sup> Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

<sup>9</sup> Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

<sup>10</sup> Supersedes MORGAN 90.

<sup>11</sup> From analysis allowing arbitrary background unconstrained by unitarity.

<sup>12</sup> Data included in MORGAN 90, BOGLIONE 99 analyses.

<sup>13</sup> From amplitude analysis of BOYER 90 and MARSISKE 90, data corresponds to resonance parameters  $m = 989$  MeV,  $\Gamma = 61$  MeV.

$\Gamma(e^+ e^-)$	$\Gamma_4$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<8.4	90	VOROBIEV	88	$e^+ e^- \rightarrow \pi^0 \pi^0$

## $f_0(980)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/[\Gamma(\pi\pi) + \Gamma(K\bar{K})]$	$\Gamma_1/(\Gamma_1 + \Gamma_2)$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
0.52 ± 0.12	9.9k	<sup>1</sup> AUBERT	060	BABR $B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$
0.75 <sup>+0.11</sup> <sub>-0.13</sub>		<sup>2</sup> ABLIKIM	05Q	BES2 $\chi_{c0} \rightarrow 2\pi^+ 2\pi^-$ , $\pi^+ \pi^- K^+ K^-$
0.84 ± 0.02		<sup>3</sup> ANISOVICH	02D	SPEC Combined fit
~0.68		OLLER	99B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
0.67 ± 0.09		<sup>4</sup> LOVERRE	80	HBC $4\pi^- p \rightarrow n2K_S^0$
0.81 <sup>+0.09</sup> <sub>-0.04</sub>		<sup>4</sup> CASON	78	STRC $7\pi^- p \rightarrow n2K_S^0$
0.78 ± 0.03		<sup>4</sup> WETZEL	76	OSPK $8.9\pi^- p \rightarrow n2K_S^0$

<sup>1</sup> Recalculated by us using  $\Gamma(K^+ K^-) / \Gamma(\pi^+ \pi^-) = 0.69 \pm 0.32$  from AUBERT 060 and isospin relations.

<sup>2</sup> Using data from ABLIKIM 04G.

<sup>3</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta\eta$ ,  $\pi^0 \pi^0 \eta$ ), GAMS ( $\pi p \rightarrow \pi^0 \pi^0 n$ ,  $\eta\eta n$ ,  $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K}n$ ) data.

<sup>4</sup> Measure  $\pi\pi$  elasticity assuming two resonances coupled to the  $\pi\pi$  and  $K\bar{K}$  channels only.

## $f_0(980)$ REFERENCES

ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)
GARCIA-MAR... 11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	
GARCIA-MAR... 11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	
MENNESSIER 11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang		
MOUSSALLAM 11	EPJ C71 1814	B. Moussallam		
BATLEY 10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	
MENNESSIER 10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang		
ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		
ECKLUND 09	PR D80 052009	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	
BATLEY 08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	
PENNINGTON 08	EPJ C56 1	M.R. Pennington <i>et al.</i>		
UEHARA 08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	
AMBROSINO 07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	
AUBERT 07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	
BONVICINI 07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	
MORI 07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	
AMBROSINO 06B	PL B634 148	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	
AUBERT 06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)	
GARMASH 06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)	
ABLIKIM 05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	
ABLIKIM 05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	
ACHASOV 05	PR D72 013006	N.N. Achasov, G.N. Shestakov		
GARMASH 05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	
ABLIKIM 04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	
BUETTIKER 04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam		
ANISOVICH 03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		
TIKHOMIROV 03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		

Translated from YAF 66 860.

ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65	1583.	
BRAMON	02	EPJ C26 253	A. Bramon <i>et al.</i>	
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99C	PL B453 325	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>	
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington	
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>	
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset	
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset	
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>	
ACKERSTAFF	98Q	EPJ C4 19	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62	446.	
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168	481.	
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ISHIDA	96	PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)
MORGAN	93	PR D48 1185	D. Morgan, M.R. Pennington	(RAL, DURH)
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko	
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
		Translated from YAF 48	436.	
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
ACHASOV	80	SJNP 32 566	N.N. Achasov, S.A. Devyanin, G.N. Shestakov	(NOVM)
		Translated from YAF 32	1098.	
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+) IJP
AGUILAR-...	78	NP B140 73	M. Aguilar-Benitez <i>et al.</i>	(MADR, BOMB+)
CASON	78	PRL 41 271	N.M. Cason <i>et al.</i>	(NDAM, ANL)
LEEPER	77	PR D16 2054	R.J. Leeper <i>et al.</i>	(ISU)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
BINNIE	73	PRL 31 1534	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)

GRAYER 73 Tallahassee  
HYAMS 73 NP B64 134  
PROTOPOP... 73 PR D7 1279

G. Grayer *et al.*  
B.D. Hyams *et al.*  
S.D. Protopopescu *et al.*

(CERN, MPIM)  
(CERN, MPIM)  
(LBL)

---