

**$\omega(1650)$**  $I^G(J^{PC}) = 0^-(1^{--})$  **$\omega(1650)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1670 \pm 30</math> OUR ESTIMATE</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1660 $\pm$ 10	898	1 ACHASOV	16B SND	1.34–2.00 $e^+e^- \rightarrow \omega\eta$
1680 $\pm$ 10	13.1k	2 AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1667 $\pm$ 13 $\pm$ 6		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1645 $\pm$ 8	13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
1660 $\pm$ 10 $\pm$ 2		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1770 $\pm$ 50 $\pm$ 60	1.2M	3 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1619 $\pm$ 5		4 HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1700 $\pm$ 20		EUGENIO	01 SPEC	18 $\pi^-p \rightarrow \omega\eta n$
1705 $\pm$ 26	612	5 AKHMETSHIN	00D CMD2	1.05–1.80 $e^+e^- \rightarrow \omega\pi^+\pi^-$
1820 $^{+190}_{-150}$		6 ACHASOV	98H RVUE	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1840 $^{+100}_{-70}$		7 ACHASOV	98H RVUE	1.05–1.80 $e^+e^- \rightarrow \omega\pi^+\pi^-$
1780 $^{+170}_{-300}$		8 ACHASOV	98H RVUE	1.05–1.80 $e^+e^- \rightarrow K^+K^-$
$\sim 2100$		9 ACHASOV	98H RVUE	1.05–1.80 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606 $\pm$ 9		10 CLEGG	94 RVUE	
1662 $\pm$ 13	750	11 ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1670 $\pm$ 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
1657 $\pm$ 13		CORDIER	81 DM1	1.05–1.80 $e^+e^- \rightarrow \omega 2\pi$
1679 $\pm$ 34	21	ESPOSITO	80 FRAM	1.05–1.80 $e^+e^- \rightarrow 3\pi$
1652 $\pm$ 17		COSME	79 OSPK	1.05–1.80 $e^+e^- \rightarrow 3\pi$

<sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ .<sup>2</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .<sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.<sup>4</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.<sup>5</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.<sup>6</sup> Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.<sup>7</sup> Using the data from ANTONELLI 92.<sup>8</sup> Using the data from IVANOV 81 and BISELLO 88B.<sup>9</sup> Using the data from BISELLO 91C.<sup>10</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.<sup>11</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.

## $\omega(1650)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>315 \pm 35</math> OUR ESTIMATE</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
110 $\pm$ 20	898	<sup>1</sup> ACHASOV	16B SND	1.34–2.00 $e^+ e^- \rightarrow \omega\eta$
310 $\pm$ 30	13.1k	<sup>2</sup> AULCHENKO	15A SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+\pi^-\pi^0$
222 $\pm$ 25 $\pm$ 20		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega\pi^+\pi^-\gamma$
114 $\pm$ 14	13	AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \omega\eta\gamma$
230 $\pm$ 30 $\pm$ 20		AUBERT,B	04N BABR	10.6 $e^+ e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
$490^{+200}_{-150} \pm 130$	1.2M	<sup>3</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+ e^- \rightarrow \pi^+\pi^-\pi^0$
250 $\pm$ 14		<sup>4</sup> HENNER	02 RVUE	1.2–2.0 $e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
250 $\pm$ 50		EUGENIO	01 SPEC	$18\pi^- p \rightarrow \omega\eta n$
370 $\pm$ 25	612	<sup>5</sup> AKHMETSHIN	00D CMD2	$e^+ e^- \rightarrow \omega\pi^+\pi^-$
113 $\pm$ 20		<sup>6</sup> CLEGG	94 RVUE	
280 $\pm$ 24	750	<sup>7</sup> ANTONELLI	92 DM2	1.34–2.4 $e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
160 $\pm$ 20		ATKINSON	83B OMEG	$20\text{--}70\gamma p \rightarrow 3\pi X$
136 $\pm$ 46		CORDIER	81 DM1	$e^+ e^- \rightarrow \omega 2\pi$
99 $\pm$ 49	21	ESPOSITO	80 FRAM	$e^+ e^- \rightarrow 3\pi$
42 $\pm$ 17		COSME	79 OSPK	$e^+ e^- \rightarrow 3\pi$

<sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ .

<sup>2</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .

<sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

<sup>4</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

<sup>5</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

<sup>6</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

<sup>7</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.

## $\omega(1650)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \rho\pi$	seen
$\Gamma_2 \omega\pi\pi$	seen
$\Gamma_3 \omega\eta$	seen
$\Gamma_4 e^+ e^-$	seen

$\omega(1650) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$ 
 $\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ 
 $\Gamma_1/\Gamma \times \Gamma_4/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
1.56 $\pm 0.23$	13.1k	<sup>1</sup> AULCHENKO	15A SND	$1.05\text{--}1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1.3 $\pm 0.1$	$\pm 0.1$	AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1.2 $\pm 0.4$	$\pm 0.8$	1.2M <sup>2,3</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.921 $\pm 0.230$		4,5 CLEGG	94 RVUE	
0.479 $\pm 0.050$	750	6,7 ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ 
 $\Gamma_2/\Gamma \times \Gamma_4/\Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
7.0 $\pm 0.5$		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
4.1 $\pm 0.9$	$\pm 1.3$	1.2M <sup>2,3</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 $\pm 0.95$		<sup>8</sup> AKHMETSHIN	00D CMD2	$1.2\text{--}1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 $\pm 0.80$		4,5 CLEGG	94 RVUE	
6.07 $\pm 0.61$	750	6,7 ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ 
 $\Gamma_3/\Gamma \times \Gamma_4/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.44 $\pm 0.05$	898	<sup>9</sup> ACHASOV	16B SND	$1.34\text{--}2.00 e^+e^- \rightarrow \omega\eta$	I
0.57 $\pm 0.06$	13	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$	
<6	90	<sup>10</sup> AKHMETSHIN	03B CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$	

<sup>1</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .

<sup>2</sup> Calculated by us from the cross section at the peak.

<sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

<sup>4</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

<sup>5</sup> From the partial and leptonic width given by the authors.

<sup>6</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.

<sup>7</sup> From the product of the leptonic width and partial branching ratio given by the authors.

<sup>8</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

<sup>9</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ .

<sup>10</sup>  $\omega(1650)$  mass and width fixed at 1700 MeV and 250 MeV, respectively.

**$\omega(1650)$  BRANCHING RATIOS** **$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$**  **$\Gamma_2/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
~ 0.35	1.2M	<sup>1</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 \pi^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0$
$0.620 \pm 0.014$		<sup>2</sup> HENNER	02 RVUE	$1.2\text{--}2.0 \pi^+ \pi^- \rightarrow \rho\pi, \omega\pi\pi$

 **$\Gamma(\rho\pi)/\Gamma_{\text{total}}$**  **$\Gamma_1/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
~ 0.65	1.2M	<sup>1</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 \pi^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0$
$0.380 \pm 0.014$		<sup>2</sup> HENNER	02 RVUE	$1.2\text{--}2.0 \pi^+ \pi^- \rightarrow \rho\pi, \omega\pi\pi$

 **$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$**  **$\Gamma_4/\Gamma$** 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
~ 18	1.2M	<sup>2,3</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 \pi^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0$
$32 \pm 1$		<sup>2</sup> HENNER	02 RVUE	$1.2\text{--}2.0 \pi^+ \pi^- \rightarrow \rho\pi, \omega\pi\pi$

<sup>1</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+ \pi^- \pi^0$  and ANTONELLI 92 on the  $\omega \pi^+ \pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

<sup>2</sup> Assuming that the  $\omega(1650)$  decays into  $\rho\pi$  and  $\omega\pi\pi$  only.

<sup>3</sup> Calculated by us from the cross section at the peak.

 **$\omega(1650)$  REFERENCES**

ACHASOV AULCHENKO	16B 15A	PR D94 092002 JETP 121 27	M.N. Achasov <i>et al.</i> V.M. Aulchenko <i>et al.</i> Translated from ZETF 148 34.	(SND Collab.) (SND Collab.)
AUBERT AUBERT AUBERT,B	07AU 06D 04N	PR D76 092005 PR D73 052003 PR D70 072004	B. Aubert <i>et al.</i> B. Aubert <i>et al.</i> B. Aubert <i>et al.</i>	(BABAR Collab.) (BABAR Collab.) (BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>	
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)