

$\phi(2170)$

$I^G(J^{PC}) = 0^-(1^{--})$

Observed by AUBERT,BE 06D in the initial-state radiation process
 $e^+ e^- \rightarrow \phi f_0(980) \gamma$.

$\phi(2170)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2188±10 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.			
2200± 6± 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta \phi \pi^+ \pi^-$	
2180± 8± 8	1,2 LEES		12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$	
2186±10± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$	
2125±22±10	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2079±13 ⁺⁷⁹ ₋₂₈	4.8k	3 SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$	
2192±14	116	4 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$	
2169±20	149	4 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$	
2175±10±15	201	2,5 AUBERT,BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$	


¹ Fit includes interference with the $\phi(1680)$.

² From the $\phi f_0(980)$ component.

³ From a fit with two incoherent Breit-Wigners.

⁴ From the $K^+ K^- f_0(980)$ component.

⁵ Superseded by LEES 12F.



$\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83±12 OUR AVERAGE				
104±15±15	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
77±15±10	6,7 LEES		12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
65±23±17	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
61±50±13	483	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
192±23 ⁺²⁵ ₋₆₁	4.8k	⁸ SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
71±21	116	⁹ AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
102±27	149	⁹ AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
58±16±20	201 ^{7,10}	AUBERT,BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$
6 Fit includes interference with the $\phi(1680)$.				
7 From the $\phi f_0(980)$ component.				
8 From a fit with two incoherent Breit-Wigners.				
9 From the $K^+K^-\pi^0\pi^0$ component.				
10 Superseded by LEES 12F.				

$\phi(2170)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+e^-$	seen
$\Gamma_2 \phi\eta$	
$\Gamma_3 \phi\pi\pi$	
$\Gamma_4 \phi f_0(980)$	seen
$\Gamma_5 K^+K^-\pi^+\pi^-$	
$\Gamma_6 K^+K^-\pi^0\pi^0 \rightarrow K^+K^-\pi^+\pi^-$	seen
$\Gamma_7 K^+K^-\pi^0\pi^0$	
$\Gamma_8 K^+K^-\pi^0\pi^0 \rightarrow K^+K^-\pi^0\pi^0$	seen
$\Gamma_9 K^{*0}K^\pm\pi^\mp$	not seen
$\Gamma_{10} K^*(892)^0\bar{K}^*(892)^0$	not seen

$\phi(2170) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_1/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.7±0.7±1.3	483	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$

$\Gamma(\phi f_0(980)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_1/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.3±0.3±0.3	11,12 LEES		12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.5±0.8±0.4	201 12,13 AUBERT,BE		06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

¹¹ From a fit with constructive interference with the $\phi(1680)$. In a fit with destructive interference, the value is larger by a factor of 12.

¹² From the $\phi f_0(980)$ component.

¹³Superseded by I FES 12E

$$\phi(2170) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$$

$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_1/\Gamma$			
VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.65 \pm 0.15 \pm 0.18	4.8k	¹⁴ SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
¹⁴ Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.				

$\phi(2170)$ BRANCHING RATIOS

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_6/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AUBERT	07AK BABR	$10.6 \text{ e}^+ \text{e}^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$	Γ_8/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AUBERT	07AK BABR	$10.6 \text{ e}^+ \text{e}^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

$\Gamma(K^{*0}K^\pm\pi^\mp)/\Gamma_{\text{total}}$	Γ_9/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	AUBERT	07akBAPP	10.6 GeV, e^+e^-

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}$	Γ_{10}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	10C	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

ϕ(2170) REFERENCES

ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)