

$\psi(2S)$

$I^G(J^{PC}) = 0^-(1^{--})$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.097±0.025 OUR FIT		Error includes scale factor of 2.6.		
3686.097±0.010 OUR AVERAGE				
3686.099±0.004±0.009		¹ ANASHIN	15	KEDR $e^+ e^- \rightarrow$ hadrons
3686.12 ± 0.06 ± 0.10	4k	AAIJ	12H	LHCb $p p \rightarrow J/\psi \pi^+ \pi^- X$
3685.95 ± 0.10	413	² ARTAMONOV 00	OLYA	$e^+ e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		³ ARMSTRONG 93B	E760	$\bar{p} p \rightarrow e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3686.114±0.007 ^{+0.011} _{-0.016}		⁴ ANASHIN	12	KEDR $e^+ e^- \rightarrow$ hadrons
3686.111±0.025±0.009		AULCHENKO 03	03	KEDR $e^+ e^- \rightarrow$ hadrons
3686.00 ± 0.10	413	⁵ ZHOLENTZ 80	OLYA	$e^+ e^-$

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁵ Superseded by ARTAMONOV 00.

$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.188±0.028 OUR AVERAGE			
589.194±0.027±0.011	¹ AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
589.7 ± 1.2	LEMOIGNE 82	GOLI 185	$\pi^- Be \rightarrow \gamma \mu^+ \mu^- A$
589.07 ± 0.13	¹ ZHOLENTZ 80	OLYA	$e^+ e^-$
588.7 ± 0.8	LUTH 75	MRK1	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
588 ± 1	² BAI 98E	BES	$e^+ e^-$

¹ Redundant with data in mass above.

² Systematic errors not evaluated.

$\psi(2S)$ WIDTH

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
296± 8 OUR FIT				
286±16 OUR AVERAGE				
358±88± 4		ABLIKIM	08B BES2	$e^+ e^- \rightarrow$ hadrons
290±25± 4	2.7k	ANDREOTTI	07 E835	$p\bar{p} \rightarrow e^+ e^-, J/\psi X$
331±58± 2		ABLIKIM	06L BES2	$e^+ e^- \rightarrow$ hadrons
264±27		¹ BAI	02B BES2	$e^+ e^-$
287±37±16		2 ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+ e^-$
¹ From a simultaneous fit to the hadronic and $\mu^+ \mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.				
² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].				

$\psi(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(97.85 ± 0.13) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(1.73 ± 0.14) %	S=1.5
Γ_3 ggg	(10.6 ± 1.6) %	
Γ_4 γgg	(1.03 ± 0.29) %	
Γ_5 light hadrons	(15.4 ± 1.5) %	
Γ_6 $e^+ e^-$	(7.89 ± 0.17) × 10 ⁻³	
Γ_7 $\mu^+ \mu^-$	(7.9 ± 0.9) × 10 ⁻³	
Γ_8 $\tau^+ \tau^-$	(3.1 ± 0.4) × 10 ⁻³	
Decays into $J/\psi(1S)$ and anything		
Γ_9 $J/\psi(1S)$ anything	(61.0 ± 0.6) %	
Γ_{10} $J/\psi(1S)$ neutrals	(25.14 ± 0.33) %	
Γ_{11} $J/\psi(1S)\pi^+\pi^-$	(34.49 ± 0.30) %	
Γ_{12} $J/\psi(1S)\pi^0\pi^0$	(18.17 ± 0.31) %	
Γ_{13} $J/\psi(1S)\eta$	(3.36 ± 0.05) %	
Γ_{14} $J/\psi(1S)\pi^0$	(1.268 ± 0.032) × 10 ⁻³	
Hadronic decays		
Γ_{15} $\pi^0 h_c(1P)$	(8.6 ± 1.3) × 10 ⁻⁴	
Γ_{16} $3(\pi^+\pi^-)\pi^0$	(3.5 ± 1.6) × 10 ⁻³	
Γ_{17} $2(\pi^+\pi^-)\pi^0$	(2.9 ± 1.0) × 10 ⁻³	S=4.7
Γ_{18} $\rho a_2(1320)$	(2.6 ± 0.9) × 10 ⁻⁴	
Γ_{19} $p\bar{p}$	(2.88 ± 0.10) × 10 ⁻⁴	
Γ_{20} $\Delta^{++}\bar{\Delta}^{--}$	(1.28 ± 0.35) × 10 ⁻⁴	
Γ_{21} $\Lambda\bar{\Lambda}\pi^0$	< 2.9 × 10 ⁻⁶	CL=90%
Γ_{22} $\Lambda\bar{\Lambda}\eta$	(2.5 ± 0.4) × 10 ⁻⁵	
Γ_{23} $\Lambda\bar{p}K^+$	(1.00 ± 0.14) × 10 ⁻⁴	
Γ_{24} $\Lambda\bar{p}K^+\pi^+\pi^-$	(1.8 ± 0.4) × 10 ⁻⁴	
Γ_{25} $\Lambda\bar{\Lambda}\pi^+\pi^-$	(2.8 ± 0.6) × 10 ⁻⁴	

Γ_{26}	$\Lambda\bar{\Lambda}$	$(3.57 \pm 0.18) \times 10^{-4}$
Γ_{27}	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.40 \pm 0.13) \times 10^{-4}$
Γ_{28}	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	$(1.54 \pm 0.14) \times 10^{-4}$
Γ_{29}	$\Sigma^0\bar{p}K^+ + \text{c.c.}$	$(1.67 \pm 0.18) \times 10^{-5}$
Γ_{30}	$\Sigma^+\bar{\Sigma}^-$	$(2.51 \pm 0.21) \times 10^{-4}$
Γ_{31}	$\Sigma^0\bar{\Sigma}^0$	$(2.32 \pm 0.16) \times 10^{-4}$
Γ_{32}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(8.5 \pm 0.7) \times 10^{-5}$
Γ_{33}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$(8.5 \pm 0.8) \times 10^{-5}$
Γ_{34}	$\Xi^-\bar{\Xi}^+$	$(2.72 \pm 0.12) \times 10^{-4}$
Γ_{35}	$\Xi^0\bar{\Xi}^0$	$(2.07 \pm 0.23) \times 10^{-4}$
Γ_{36}	$\Xi(1530)^0\bar{\Xi}(1530)^0$	$(5.2 \begin{array}{l} +3.2 \\ -1.2 \end{array}) \times 10^{-5}$
Γ_{37}	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(3.9 \pm 0.4) \times 10^{-5}$
Γ_{38}	$\Xi(1690)^-\bar{\Xi}^+ \rightarrow K^-\Lambda\bar{\Xi}^+ +$	$(5.2 \pm 1.6) \times 10^{-6}$
Γ_{39}	$\Xi(1820)^-\bar{\Xi}^+ \rightarrow K^-\Lambda\bar{\Xi}^+ +$ <small>c.c.</small>	$(1.20 \pm 0.32) \times 10^{-5}$
Γ_{40}	$K^-\Sigma^0\bar{\Xi}^+ + \text{c.c.}$	$(3.7 \pm 0.4) \times 10^{-5}$
Γ_{41}	$\Omega^-\bar{\Omega}^+$	$(4.7 \pm 1.0) \times 10^{-5}$
Γ_{42}	$\pi^0 p\bar{p}$	$(1.53 \pm 0.07) \times 10^{-4}$
Γ_{43}	$N(940)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(6.4 \begin{array}{l} +1.8 \\ -1.3 \end{array}) \times 10^{-5}$
Γ_{44}	$N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(7.3 \begin{array}{l} +1.7 \\ -1.5 \end{array}) \times 10^{-5}$
Γ_{45}	$N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(6.4 \begin{array}{l} +2.3 \\ -1.8 \end{array}) \times 10^{-6}$
Γ_{46}	$N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(2.5 \pm 1.0) \times 10^{-5}$
Γ_{47}	$N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(3.8 \begin{array}{l} +1.4 \\ -1.7 \end{array}) \times 10^{-5}$
Γ_{48}	$N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(1.79 \begin{array}{l} +0.26 \\ -0.70 \end{array}) \times 10^{-5}$
Γ_{49}	$N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(2.6 \begin{array}{l} +1.2 \\ -0.7 \end{array}) \times 10^{-5}$
Γ_{50}	$N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	$(2.13 \begin{array}{l} +0.40 \\ -0.31 \end{array}) \times 10^{-5}$
Γ_{51}	$\pi^0 f_0(2100) \rightarrow \pi^0 p\bar{p}$	$(1.1 \pm 0.4) \times 10^{-5}$
Γ_{52}	$\eta p\bar{p}$	$(6.0 \pm 0.4) \times 10^{-5}$
Γ_{53}	$\eta f_0(2100) \rightarrow \eta p\bar{p}$	$(1.2 \pm 0.4) \times 10^{-5}$
Γ_{54}	$N(1535)\bar{p} \rightarrow \eta p\bar{p}$	$(4.4 \pm 0.7) \times 10^{-5}$
Γ_{55}	$\omega p\bar{p}$	$(6.9 \pm 2.1) \times 10^{-5}$
Γ_{56}	$\phi p\bar{p}$	$< 2.4 \times 10^{-5}$ CL=90%
Γ_{57}	$\pi^+\pi^- p\bar{p}$	$(6.0 \pm 0.4) \times 10^{-4}$
Γ_{58}	$p\bar{n}\pi^- \text{ or c.c.}$	$(2.48 \pm 0.17) \times 10^{-4}$
Γ_{59}	$p\bar{n}\pi^-\pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$
Γ_{60}	$2(\pi^+\pi^-\pi^0)$	$(4.8 \pm 1.5) \times 10^{-3}$
Γ_{61}	$\eta\pi^+\pi^-$	$< 1.6 \times 10^{-4}$ CL=90%
Γ_{62}	$\eta\pi^+\pi^-\pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$
Γ_{63}	$2(\pi^+\pi^-)\eta$	$(1.2 \pm 0.6) \times 10^{-3}$

Γ_{64}	$\eta' \pi^+ \pi^- \pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$	
Γ_{65}	$\omega \pi^+ \pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{66}	$b_1^\pm \pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1
Γ_{67}	$b_1^0 \pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$	
Γ_{68}	$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{69}	$\pi^0 \pi^0 K^+ K^-$	$(2.6 \pm 1.3) \times 10^{-4}$	
Γ_{70}	$\pi^+ \pi^- K^+ K^-$	$(7.3 \pm 0.5) \times 10^{-4}$	
Γ_{71}	$\rho^0 K^+ K^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{72}	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{73}	$K^+ K^- \pi^+ \pi^- \eta$	$(1.3 \pm 0.7) \times 10^{-3}$	
Γ_{74}	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$	
Γ_{75}	$K^+ K^- 2(\pi^+ \pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$	
Γ_{76}	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$	
Γ_{77}	$K_S^0 K_S^0 \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{78}	$\rho^0 p \bar{p}$	$(5.0 \pm 2.2) \times 10^{-5}$	
Γ_{79}	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$	
Γ_{80}	$2(\pi^+ \pi^-)$	$(2.4 \pm 0.6) \times 10^{-4}$	S=2.2
Γ_{81}	$\rho^0 \pi^+ \pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	S=1.4
Γ_{82}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$	
Γ_{83}	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$(5.9 \pm 2.2) \times 10^{-5}$	
Γ_{84}	$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$	
Γ_{85}	$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$	
Γ_{86}	$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$	
Γ_{87}	$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$	
Γ_{88}	$\eta K^+ K^-$, no $\eta \phi$	$(3.1 \pm 0.4) \times 10^{-5}$	
Γ_{89}	$\omega K^+ K^-$	$(1.62 \pm 0.11) \times 10^{-4}$	S=1.1
Γ_{90}	$\omega K^*(892)^+ K^- + \text{c.c.}$	$(2.07 \pm 0.26) \times 10^{-4}$	
Γ_{91}	$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$(6.1 \pm 1.2) \times 10^{-5}$	
Γ_{92}	$\omega \bar{K}^*(892)^0 K^0$	$(1.68 \pm 0.30) \times 10^{-4}$	
Γ_{93}	$\omega \bar{K}_2^*(1430)^0 K^0$	$(5.8 \pm 2.2) \times 10^{-5}$	
Γ_{94}	$\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$	
Γ_{95}	$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	$(1.09 \pm 0.26) \times 10^{-5}$	
Γ_{96}	$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(3.0 \pm 1.0) \times 10^{-6}$	
Γ_{97}	$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	$(1.2 \pm 0.7) \times 10^{-6}$	
Γ_{98}	$3(\pi^+ \pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8
Γ_{99}	$p \bar{p} \pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$	
Γ_{100}	$K^+ K^-$	$(7.5 \pm 0.5) \times 10^{-5}$	
Γ_{101}	$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$	
Γ_{102}	$\pi^+ \pi^- \pi^0$	$(2.01 \pm 0.17) \times 10^{-4}$	S=1.7
Γ_{103}	$\rho(2150) \pi \rightarrow \pi^+ \pi^- \pi^0$	$(1.9 \pm 1.2) \times 10^{-4}$	
Γ_{104}	$\rho(770) \pi \rightarrow \pi^+ \pi^- \pi^0$	$(3.2 \pm 1.2) \times 10^{-5}$	S=1.8

Γ_{105}	$\pi^+ \pi^-$	$(7.8 \pm 2.6) \times 10^{-6}$	
Γ_{106}	$K_1(1400)^{\pm} K^{\mp}$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{107}	$K_2^*(1430)^{\pm} K^{\mp}$	$(7.1 \pm 1.3) \times 10^{-5}$	
Γ_{108}	$K^+ K^- \pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$	
Γ_{109}	$K^+ K^*(892)^- + \text{c.c.}$	$(2.9 \pm 0.4) \times 10^{-5}$	S=1.2
Γ_{110}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$	
Γ_{111}	$\phi \pi^+ \pi^-$	$(1.18 \pm 0.26) \times 10^{-4}$	S=1.5
Γ_{112}	$\phi f_0(980) \rightarrow \pi^+ \pi^-$	$(7.5 \pm 3.3) \times 10^{-5}$	S=1.6
Γ_{113}	$2(K^+ K^-)$	$(6.3 \pm 1.3) \times 10^{-5}$	
Γ_{114}	$\phi K^+ K^-$	$(7.0 \pm 1.6) \times 10^{-5}$	
Γ_{115}	$2(K^+ K^-) \pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$	
Γ_{116}	$\phi \eta$	$(3.10 \pm 0.31) \times 10^{-5}$	
Γ_{117}	$\phi \eta'$	$(3.1 \pm 1.6) \times 10^{-5}$	
Γ_{118}	$\omega \eta'$	$(3.2 \pm 2.5) \times 10^{-5}$	
Γ_{119}	$\omega \pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$	
Γ_{120}	$\rho \eta'$	$(1.9 \pm 1.7) \times 10^{-5}$	
Γ_{121}	$\rho \eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1
Γ_{122}	$\omega \eta$	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{123}	$\phi \pi^0$	$< 4 \times 10^{-7}$	CL=90%
Γ_{124}	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%
Γ_{125}	$p \bar{p} K^+ K^-$	$(2.7 \pm 0.7) \times 10^{-5}$	
Γ_{126}	$\Lambda n K_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$	
Γ_{127}	$\phi f'_2(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$	
Γ_{128}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	$< 8.8 \times 10^{-6}$	CL=90%
Γ_{129}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 1.0 \times 10^{-5}$	CL=90%
Γ_{130}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 7.0 \times 10^{-6}$	CL=90%
Γ_{131}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 2.6 \times 10^{-5}$	CL=90%
Γ_{132}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 6.0 \times 10^{-6}$	CL=90%
Γ_{133}	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$	

Radiative decays

Γ_{134}	$\gamma \chi_{c0}(1P)$	$(9.99 \pm 0.27) \%$	
Γ_{135}	$\gamma \chi_{c1}(1P)$	$(9.55 \pm 0.31) \%$	
Γ_{136}	$\gamma \chi_{c2}(1P)$	$(9.11 \pm 0.31) \%$	
Γ_{137}	$\gamma \eta_c(1S)$	$(3.4 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{138}	$\gamma \eta_c(2S)$	$(7 \pm 5) \times 10^{-4}$	
Γ_{139}	$\gamma \pi^0$	$(1.6 \pm 0.4) \times 10^{-6}$	
Γ_{140}	$\gamma \eta'(958)$	$(1.23 \pm 0.06) \times 10^{-4}$	
Γ_{141}	$\gamma f_2(1270)$	$(2.73 \pm 0.29) \times 10^{-4}$	S=1.8
Γ_{142}	$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(3.1 \pm 1.7) \times 10^{-5}$	
Γ_{143}	$\gamma f_0(1500)$	$(9.2 \pm 1.9) \times 10^{-5}$	

Γ_{144}	$\gamma f'_2(1525)$	$(3.3 \pm 0.8) \times 10^{-5}$	
Γ_{145}	$\gamma f_0(1710)$		
Γ_{146}	$\gamma f_0(1710) \rightarrow \gamma \pi\pi$	$(3.5 \pm 0.6) \times 10^{-5}$	
Γ_{147}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(6.6 \pm 0.7) \times 10^{-5}$	
Γ_{148}	$\gamma f_0(2100) \rightarrow \gamma \pi\pi$	$(4.8 \pm 1.0) \times 10^{-6}$	
Γ_{149}	$\gamma f_0(2200) \rightarrow \gamma K\bar{K}$	$(3.2 \pm 1.0) \times 10^{-6}$	
Γ_{150}	$\gamma f_J(2220) \rightarrow \gamma \pi\pi$	$< 5.8 \times 10^{-6}$	CL=90%
Γ_{151}	$\gamma f_J(2220) \rightarrow \gamma K\bar{K}$	$< 9.5 \times 10^{-6}$	CL=90%
Γ_{152}	$\gamma\gamma$	$< 1.5 \times 10^{-4}$	CL=90%
Γ_{153}	$\gamma\eta$	$(1.4 \pm 0.5) \times 10^{-6}$	
Γ_{154}	$\gamma\eta\pi^+\pi^-$	$(8.7 \pm 2.1) \times 10^{-4}$	
Γ_{155}	$\gamma\eta(1405)$		
Γ_{156}	$\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi$	$< 9 \times 10^{-5}$	CL=90%
Γ_{157}	$\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$	
Γ_{158}	$\gamma\eta(1475)$		
Γ_{159}	$\gamma\eta(1475) \rightarrow K\bar{K}\pi$	$< 1.4 \times 10^{-4}$	CL=90%
Γ_{160}	$\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-$	$< 8.8 \times 10^{-5}$	CL=90%
Γ_{161}	$\gamma 2(\pi^+\pi^-)$	$(4.0 \pm 0.6) \times 10^{-4}$	
Γ_{162}	$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	$(3.7 \pm 0.9) \times 10^{-4}$	
Γ_{163}	$\gamma K^{*0} \bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$	
Γ_{164}	$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{165}	$\gamma K^+ K^- \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{166}	$\gamma p\bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0
Γ_{167}	$\gamma f_2(1950) \rightarrow \gamma p\bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$	
Γ_{168}	$\gamma f_2(2150) \rightarrow \gamma p\bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$	
Γ_{169}	$\gamma X(1835) \rightarrow \gamma p\bar{p}$	$(4.6 \pm 1.8) \times 10^{-6}$	
Γ_{170}	$\gamma X \rightarrow \gamma p\bar{p}$	[a] $< 2 \times 10^{-6}$	CL=90%
Γ_{171}	$\gamma\pi^+\pi^- p\bar{p}$	$(2.8 \pm 1.4) \times 10^{-5}$	
Γ_{172}	$\gamma 2(\pi^+\pi^-) K^+ K^-$	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{173}	$\gamma 3(\pi^+\pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{174}	$\gamma K^+ K^- K^+ K^-$	$< 4 \times 10^{-5}$	CL=90%
Γ_{175}	$\gamma\gamma J/\psi$	$(3.1 \pm 1.0) \times 10^{-4}$	

Other decays

Γ_{176}	invisible	$< 1.6 \%$	CL=90%
----------------	-----------	------------	--------

[a] For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 239 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 342.4$ for 190 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_7	3										
x_8	1	0									
x_{11}	30	8	2								
x_{12}	29	5	1	49							
x_{13}	13	3	1	36	16						
x_{19}	0	0	0	5	3	2					
x_{134}	1	0	0	3	1	1	0				
x_{135}	2	0	0	4	1	1	0	0			
x_{136}	1	0	0	4	1	1	0	0	0		
Γ	-81	-3	-1	-39	-35	-17	-8	-1	-2	-2	
	x_6	x_7	x_8	x_{11}	x_{12}	x_{13}	x_{19}	x_{134}	x_{135}	x_{136}	

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
258 ± 26	BAI	02B BES2	$e^+ e^-$
224 ± 56	LUTH	75 MRK1	$e^+ e^-$

$\Gamma(e^+ e^-)$

Γ_6

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

2.34 ± 0.04 OUR FIT

2.30 ± 0.06 OUR AVERAGE

2.24 ± 0.10 ± 0.02	¹ ABLIKIM	15V BES3	$4.0\text{--}4.4\text{ }e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
2.338 ± 0.037 ± 0.096	ABLIKIM	08B BES2	$e^+ e^- \rightarrow \text{hadrons}$
2.330 ± 0.036 ± 0.110	ABLIKIM	06L BES2	$e^+ e^- \rightarrow \text{hadrons}$
2.44 ± 0.21	² BAI	02B BES2	$e^+ e^-$
2.14 ± 0.21	ALEXANDER	89 RVUE	See γ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.0 ± 0.3	BRANDELIK	79C DASP	$e^+ e^-$
2.1 ± 0.3	³ LUTH	75 MRK1	$e^+ e^-$

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau / 0.38847$.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

$\Gamma(\gamma\gamma)$	Γ_{152}			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<43	90	BRANDELIK	79C DASP	e^+e^-

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(*i*) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.


$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_6/\Gamma$		
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.233±0.015±0.042	¹ ANASHIN	12 KEDR	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.2 ± 0.4	ABRAMS	75 MRK1	e^+e^-
¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.			

$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_8\Gamma_6/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.0 ± 2.6	79	¹ ANASHIN	07 KEDR	$e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{11}\Gamma_6/\Gamma$			
<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.807±0.013 OUR FIT				
0.837±0.025 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.837 $\pm 0.028 \pm 0.005$	1 LEES	12E BABR	10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$	
0.852 $\pm 0.010 \pm 0.026$	19.5k ADAM	06 CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$	
0.68 ± 0.09	2 BAI	98E BES	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.88 $\pm 0.08 \pm 0.03$	256 ³ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$	
0.755 $\pm 0.048 \pm 0.004$	544 ⁴ AUBERT	05D BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$	

- ¹ LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ³ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.11 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.



$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{12}\Gamma_6/\Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.425±0.009 OUR FIT				
0.411±0.008±0.018	$3.6k \pm 96$	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.6 ± 1.6 OUR FIT				
87 ± 9 OUR AVERAGE				
83 ± 25 ± 5	14	¹ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ± 7	291 ± 24	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$
${}^1\text{AUBERT } 07\text{AU quotes } \Gamma_{ee}^{\psi(2S)} \cdot \mathcal{B}(\psi(2S) \rightarrow J/\psi\eta) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+\mu^-) \cdot \mathcal{B}(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07 \text{ eV.}$				

 $\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_6/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8	90	<37	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$

 $\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.673 ± 0.023 OUR FIT				

0.63 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.2.
0.67 ± 0.12 ± 0.02	¹ LEES 130 BABR $e^+e^- \rightarrow p\bar{p}\gamma$
0.74 ± 0.07 ± 0.04	² LEES 13Y BABR $e^+e^- \rightarrow p\bar{p}\gamma$
$0.579 \pm 0.038 \pm 0.036$	ANDREOTTI 07 E835 $p\bar{p} \rightarrow e^+e^-$, $J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.70 ± 0.17 ± 0.03	³ AUBERT 06B BABR $e^+e^- \rightarrow p\bar{p}\gamma$

¹ISR photon reconstructed in the detector²ISR photon undetected³Superseded by LEES 130 $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.5 ± 0.4 ± 0.1		AUBERT	07BD	BABR $10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

 $\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{60}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.2 ± 3.3 ± 1.3	43	AUBERT	06D	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

 $\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{69}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.60 ± 0.31 ± 0.03	17	LEES	12F	BABR $10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$

 $\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{75}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4 ± 2.1 ± 0.3	26	AUBERT	06D	BABR $10.6 e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$

$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{70}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.92±0.30±0.06	133	LEES	12F	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

2.56±0.42±0.16 85 ¹AUBERT 07AK BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹ Superseded by LEES 12F.

 $\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{112}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.347±0.129±0.003	12	¹ LEES	12F	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.347±0.168±0.003 6 ± 3 ²AUBERT 07AK BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹ LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.17 \pm 0.06 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.17 \pm 0.08 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{113}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.22±0.10±0.02	13	LEES	12F	BABR $10.6 e^+e^- \rightarrow K^+K^-K^+K^-\gamma$

 $\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{111}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.55±0.19±0.01	19	¹ LEES	12F	BABR $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.57±0.23±0.01 10 ²AUBERT,BE 06D BABR $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

¹ LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.27 \pm 0.09 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.28 \pm 0.11 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2(\pi^+\pi^-)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.7±2.2±1.8	410	AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{65}\Gamma_6/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.01±0.84±0.02	37	1 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.69 \pm 0.73 \pm 0.16$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{63}\Gamma_6/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.87±1.41±0.01	16	1 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{82}\Gamma_6/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±1.3±0.3	32	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$

$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{73}\Gamma_6/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.04±1.79±0.02	7	1 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{100}\Gamma_6/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.147±0.035±0.005	66	1 LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
0.197±0.035±0.005	66	2 LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
0.35 ± 0.14 ± 0.03	11	3 LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant K^+K^- production not taken into account.

$\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.9785±0.0013 OUR AVERAGE			
0.9779±0.0015	1 BAI	02B BES2	e^+e^-
0.981 ± 0.003	1 LUTH	75 MRK1	e^+e^-

¹ Includes cascade decay into $J/\psi(1S)$.

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0173±0.0014 OUR AVERAGE	Error includes scale factor of 1.5.		
0.0166±0.0010	1, ² SETH	04	RVUE $e^+ e^-$
0.0199±0.0019	¹ BAI	02B	BES2 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.029 ± 0.004	¹ LUTH	75	MRK1 $e^+ e^-$

¹ Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

² Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(gg\bar{g})/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
10.58±1.62	2.9 M	¹ LIBBY	09	CLEO $\psi(2S) \rightarrow \text{hadrons}$
¹ Calculated using $\Gamma(\gamma gg)/\Gamma(gg\bar{g}) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow X J/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma \eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+ \ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.				

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.025±0.288	200 k	¹ LIBBY	09	CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$
¹ Calculated using $\Gamma(\gamma gg)/\Gamma(gg\bar{g}) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(gg\bar{g})/\Gamma_{\text{total}}$ LIBBY 09 measurement.				

$\Gamma(\gamma gg)/\Gamma(gg\bar{g})$

Γ_4/Γ_3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.7±2.6±1.6	2.9 M	LIBBY	09	CLEO $\psi(2S) \rightarrow (\gamma +)$ hadrons

$\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.154±0.015	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.169±0.026	² ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S)$
¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07. ² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ} \gamma)$, $B(\eta_c \gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04. Superseded by MENDEZ 08.			

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

Γ_6/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
78.9± 1.7 OUR FIT			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
88 ± 13	¹ FELDMAN	77	RVUE $e^+ e^-$
¹ From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.			

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
79 ± 9 OUR FIT	

$\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_7/Γ_6

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.00 ± 0.11 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 ± 0.16 BOYARSKI 75C MRK1 e^+e^-

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
31 ± 4 OUR FIT			

$30.8 \pm 2.1 \pm 3.8$ ¹ ABLIKIM 06W BES $e^+e^- \rightarrow \psi(2S)$

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

— DECAYS INTO $J/\psi(1S)$ AND ANYTHING —

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.610 ± 0.006 OUR FIT

0.55 ± 0.07 OUR AVERAGE

0.51 ± 0.12 BRANDELIK 79C DASP $e^+e^- \rightarrow \mu^+\mu^-X$

0.57 ± 0.08 ABRAMS 75B MRK1 $e^+e^- \rightarrow \mu^+\mu^-X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.6254 \pm 0.0016 \pm 0.0155$ 1.1M ¹ MENDEZ 08 CLEO $\psi(2S) \rightarrow \ell^+\ell^-X$

$0.5950 \pm 0.0015 \pm 0.0190$ 151k ADAM 05A CLEO Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_6/\Gamma_9 = \Gamma_6 / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.339\Gamma_{135} + 0.192\Gamma_{136})$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

1.294 ± 0.026 OUR FIT


1.28 ± 0.04 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

$1.22 \pm 0.02 \pm 0.05$ 5097 ± 73 ¹ ANDREOTTI 05 E835 $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$

$1.28 \pm 0.03 \pm 0.02$ ¹ AMBROGIANI 00A E835 $p\bar{p} \rightarrow \psi(2S)$

$1.44 \pm 0.08 \pm 0.02$ ¹ ARMSTRONG 97 E760 $\bar{p}p \rightarrow \psi(2S)$

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.



$$\Gamma(\mu^+ \mu^-)/\Gamma(J/\psi(1S) \text{anything})$$

$$\Gamma_7/\Gamma_9 = \Gamma_7/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.339\Gamma_{135} + 0.192\Gamma_{136})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0130±0.0014 OUR FIT			
0.014 ± 0.003	HILGER	75	SPEC $e^+ e^-$

$$\Gamma(J/\psi(1S) \text{ neutrals})/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma$$


VALUE	DOCUMENT ID
0.2514±0.0033 OUR FIT	

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_{11}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3449±0.0030 OUR FIT				
0.348 ± 0.005 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
0.3498±0.0002±0.0045	20M	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
0.3504±0.0007±0.0077	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.323 ± 0.014		BAI	02B	BES2 $e^+ e^-$
0.32 ± 0.04		ABRAMS	75B	MRK1 $e^+ e^- \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.3354±0.0014±0.0110	60k	¹ ADAM	05A	CLEO Repl. by MENDEZ 08

¹ Not independent from other values reported by ADAM 05A.



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_{11}
0.0229±0.0005 OUR FIT				
0.0252±0.0028±0.0011	¹ AUBERT	02B	BABR e^+e^-	

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_{11}
0.0229±0.0025 OUR FIT				
0.0224±0.0029 OUR AVERAGE				
0.0216±0.0026±0.0014	¹ AUBERT	02B	BABR e^+e^-	
0.0327±0.0077±0.0072	¹ GRIBUSHIN	96	FMPS 515 π^- Be → $2\mu X$	

¹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

$\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_{11}
8.9 ±1.1 OUR FIT				
8.73±1.39±1.57	BAI	02	BES e^+e^-	


$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{11}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.5653±0.0026 OUR FIT				
0.554 ±0.008 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.5604±0.0009±0.0062	565k	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.525 ±0.009 ±0.022	4k	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.536 ±0.007 ±0.016	20k	1,2 ABLIKIM 04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.496 ±0.037		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.5637±0.0027±0.0046	60k	ADAM 05A	CLEO	Repl. by MENDEZ 08

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$.



$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$$

Γ_{11}/Γ_9

$\Gamma(J/\psi(1S)\text{ neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$$\Gamma_{10}/\Gamma_{11} = (0.9761\Gamma_{12} + 0.719\Gamma_{13} + 0.339\Gamma_{135} + 0.192\Gamma_{136})/\Gamma_{11}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.729±0.008 OUR FIT			
0.73 ±0.09	TANENBAUM 76	MRK1	$e^+ e^-$

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1817±0.0031 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.1769±0.0008±0.0053	61k	1 MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.1652±0.0014±0.0058	13.4k	2 ADAM 05A	CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{12}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2977 \pm 0.0031 OUR FIT				
0.320 \pm 0.012 OUR AVERAGE				
0.300 \pm 0.008	\pm 0.022	1655 \pm 44	ANDREOTTI 05 E835	$\psi(2S) \rightarrow J/\psi X$
0.328 \pm 0.013	\pm 0.008		AMBROGIANI 00A E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 \pm 0.033			ARMSTRONG 97 E760	$\bar{p}p \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.2829 \pm 0.0012 \pm 0.0056	61k	MENDEZ 08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$	
0.2776 \pm 0.0025 \pm 0.0043	13.4k	ADAM 05A CLEO	Repl. by MENDEZ 08	

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{12}/Γ_{11}


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.527 \pm 0.008 OUR FIT				
0.513 \pm 0.022 OUR AVERAGE Error includes scale factor of 2.2.				
0.5047 \pm 0.0022 \pm 0.0102	61k	MENDEZ 08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$	
0.570 \pm 0.009 \pm 0.026	14k	¹ ABLIKIM 04B BES	$\psi(2S) \rightarrow J/\psi X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.4924 \pm 0.0047 \pm 0.0086	73k	^{2,3} ADAM 05A CLEO	Repl. by MENDEZ 08	
0.571 \pm 0.018 \pm 0.044		⁴ ANDREOTTI 05 E835	$\psi(2S) \rightarrow J/\psi X$	
0.53 \pm 0.06		TANENBAUM 76 MRK1	$e^+ e^-$	
0.64 \pm 0.15		⁵ HILGER 75 SPEC	$e^+ e^-$	

¹ From a fit to the J/ψ recoil mass spectra.² Not independent from other values reported by ADAM 05A.³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.⁴ Not independent from other values reported by ANDREOTTI 05.⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
33.6 \pm 0.5 OUR FIT				
32.9 \pm 1.7 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.				
33.75 \pm 0.17 \pm 0.86	68.2k	ABLIKIM 12M BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$	
29.8 \pm 0.9 \pm 2.3	5.7k	BAI 04I BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$	
25.5 \pm 2.9	386	¹ OREGLIA 80 CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$	
45 \pm 12	17	² BRANDELIK 79B DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$	
42 \pm 6	164	² BARTEL 78B CNTR	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
34.3 \pm 0.4 \pm 0.9	18.4k	³ MENDEZ 08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$	
32.5 \pm 0.6 \pm 1.1	2.8k	⁴ ADAM 05A CLEO	Repl. by MENDEZ 08	
43 \pm 8	44	TANENBAUM 76 MRK1	$e^+ e^-$	

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.³ Not independent from other measurements of MENDEZ 08.⁴ Not independent from other values reported by ADAM 05A.




$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{13}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0551 ± 0.0008 OUR FIT				
0.058 ± 0.007 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.050 ± 0.006	± 0.003	298 ± 20	ANDREOTTI 05 E835	$\psi(2S) \rightarrow J/\psi X$
0.072 ± 0.009			AMBROGIANI 00A E835	$p\bar{p} \rightarrow \psi(2S)$
0.061 ± 0.015			ARMSTRONG 97 E760	$\bar{p}p \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.0549 \pm 0.0006 \pm 0.0009$	18.4k	¹ MENDEZ 08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$	
$0.0546 \pm 0.0010 \pm 0.0007$	2.8k	ADAM 05A CLEO	Repl. by MENDEZ 08	

¹ Not independent from other measurements of MENDEZ 08.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{13}/Γ_{11}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0974±0.0014 OUR FIT				
0.0979±0.0018 OUR AVERAGE				
0.0979±0.0010±0.0015	18.4k	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.098 ± 0.005 ± 0.010	2k	¹ ABLIKIM 04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ± 0.021		² HIMEL 80	MRK2	$e^+ e^- \rightarrow \psi(2S) X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0968±0.0019±0.0013	2.8k	³ ADAM 05A	CLEO	Repl. by MENDEZ 08
0.095 ± 0.007 ± 0.007		⁴ ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S)) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.68±0.32 OUR AVERAGE				
12.6 ± 0.2 ± 0.3	4.1k	ABLIKIM 12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
13.3 ± 0.8 ± 0.3	530	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
14.3 ± 1.4 ± 1.2	280	BAI 04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
14 ± 6	7	HIMEL 80	MRK2	$e^+ e^-$
9 ± 2 ± 1	23	¹ OREGLIA 80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13 ± 1 ± 1	88	ADAM 05A	CLEO	Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.339\Gamma_{135} + 0.192\Gamma_{136})$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.213 $\pm 0.012 \pm 0.003$	527	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow J/\psi \gamma\gamma$
0.22 $\pm 0.02 \pm 0.01$		² ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma\gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$$\Gamma_{14}/\Gamma_{11}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.380 $\pm 0.022 \pm 0.005$	527	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow J/\psi \gamma\gamma$
0.39 $\pm 0.04 \pm 0.01$		² ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma\gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

— HADRONIC DECAYS —

$\Gamma(\pi^0 h_c(1P))/\Gamma_{\text{total}}$

$$\Gamma_{15}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 ± 1.3 OUR AVERAGE				
9.0 $\pm 1.5 \pm 1.3$	3k	¹ GE	11	CLEO $\psi(2S) \rightarrow \pi^0$ anything
8.4 $\pm 1.3 \pm 1.0$	11k	ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 h_c$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	92^{+23}_{-22}	ADAMS	09	CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$
seen	1282	DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
seen	168 ± 40	ROSNER	05	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86$ MeV $\equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

$$\Gamma_{16}/\Gamma$$


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
35 ± 16	6	FRANKLIN	83	MRK2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

$$\Gamma_{17}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
29 ± 10 OUR AVERAGE				
				Error includes scale factor of 4.7. See the ideogram below.
24.9 $\pm 0.7 \pm 3.6$	2173	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$
127 $\pm 12 \pm 2$	410	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)\pi^0 \gamma$
26.1 $\pm 0.7 \pm 3.0$	1703	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
30 ± 8	42	FRANKLIN	83	MRK2 $e^+ e^-$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.55±0.73±0.47		112 ± 31	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3		90	BAI	98J BES	$e^+ e^-$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.10 OUR FIT				
3.00±0.13 OUR AVERAGE				Error includes scale factor of 1.1.
3.08±0.05±0.18	4.5k	¹ DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.36±0.09±0.25	1.6k	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.87±0.12±0.15	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
1.4 ± 0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.3 ± 0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{19}/Γ_{11} VALUE (units 10^{-4}) **8.34 ± 0.28 OUR FIT** **$6.98 \pm 0.49 \pm 0.97$** DOCUMENT IDTECNCOMMENT

BAI

01

BES

 $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ $\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$ Γ_{20}/Γ VALUE (units 10^{-5})EVTSDOCUMENT IDTECNCOMMENT **$12.8 \pm 1.0 \pm 3.4$**

157

1 BAI

01

BES

 $e^+e^- \rightarrow \psi(2S) \rightarrow$
hadrons¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$. $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ VALUE (units 10^{-5})CL%DOCUMENT IDTECNCOMMENT**< 0.29**

90

1 ABLIKIM

13F

BES3

 $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12

90

2 ABLIKIM

07H

BES2

 $e^+e^- \rightarrow \psi(2S)$ ¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$. $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{22}/Γ VALUE (units 10^{-5})CL%EVTSDOCUMENT IDTECNCOMMENT **$2.48 \pm 0.34 \pm 0.19$**

60

1 ABLIKIM

13F

BES3

 $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.9

90

2 ABLIKIM

07H

BES2

 $e^+e^- \rightarrow \psi(2S)$ ¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$. $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{23}/Γ VALUE (units 10^{-4})EVTSDOCUMENT IDTECNCOMMENT **$1.0 \pm 0.1 \pm 0.1$**

74.0

BRIERE

05

CLEO

 $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^-$ $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ VALUE (units 10^{-4})EVTSDOCUMENT IDTECNCOMMENT **$1.8 \pm 0.3 \pm 0.3$**

45.8

BRIERE

05

CLEO

 $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^+\pi^-\pi^-$ $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ VALUE (units 10^{-4})EVTSDOCUMENT IDTECNCOMMENT **$2.8 \pm 0.4 \pm 0.5$**

73.4

BRIERE

05

CLEO

 $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}2(\pi^+\pi^-)$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.57 ± 0.18 OUR AVERAGE					
3.75 $\pm 0.09 \pm 0.23$	1.9k	1	DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
3.39 $\pm 0.20 \pm 0.32$	337	ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
6.4 $\pm 1.8 \pm 0.1$		2	AUBERT	07BD BABR	$10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
3.28 $\pm 0.23 \pm 0.25$	208	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.81 $\pm 0.20 \pm 0.27$	80	3 BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
< 4	90	FELDMAN	77 MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm$

$1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

 $\Gamma(\Lambda\bar{\Sigma}^+ + c.c.)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.40 \pm 0.03 \pm 0.13$	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

 $\Gamma(\Lambda\bar{\Sigma}^- + c.c.)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.54 \pm 0.04 \pm 0.13$	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

 $\Gamma(\Sigma^0\bar{p}K^+ + c.c.)/\Gamma_{\text{total}}$ Γ_{29}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.67 \pm 0.13 \pm 0.12$	276	1 ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$

¹ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$.

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.51 ± 0.21 OUR AVERAGE				
2.51 $\pm 0.15 \pm 0.16$	281	1 DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.57 $\pm 0.44 \pm 0.68$	35	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.32 ± 0.16 OUR AVERAGE				
2.25 $\pm 0.11 \pm 0.16$	439	1 DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.35 $\pm 0.36 \pm 0.32$	59	ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.63 $\pm 0.35 \pm 0.21$	58	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 $\pm 0.4 \pm 0.4$ 8 ² BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
8.5±0.7 OUR AVERAGE					
8.4±0.5±0.5	1.5k	ABLIKIM	16L	BES3	$\psi(2S) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
11 ± 3	± 3	14	1 BAI	01	BES

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
8.5±0.6±0.6					

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
2.72±0.12 OUR AVERAGE						
2.78±0.05±0.14	5k	ABLIKIM	16L	BES3	$\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$	
2.66±0.12±0.20	548	¹ DOBBS	14		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
3.03±0.40±0.32	67	ABLIKIM	07C	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
2.38±0.30±0.21	63	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.94±0.27±0.15	12	² BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<2	90	FEELDMAN	77	MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.07±0.23 OUR AVERAGE					
2.02±0.19±0.15	112	¹ DOBBS	14		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.75±0.64±0.61	19	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration. $\Gamma(\Xi(1530)^0 \bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
5.2±0.3^{+3.2}_{-1.2}		527	¹ ABLIKIM	13S	BES3	$\psi(2S) \rightarrow \eta p \bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<32	90	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
< 8.1	90	² BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

¹ With $N(1535)$ decaying to $p\eta$.² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.86±0.27±0.32					

$\Gamma(\Xi(1690)^-\Xi^+ \rightarrow K^-\Lambda\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{38}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.21±1.48±0.57	74	ABLIKIM	15I	$e^+e^- \rightarrow \psi(2S) \rightarrow K^-\Lambda\Xi^+ + \text{c.c.}$

 $\Gamma(\Xi(1820)^-\Xi^+ \rightarrow K^-\Lambda\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.03±2.94±1.22	136	ABLIKIM	15I	$e^+e^- \rightarrow \psi(2S) \rightarrow K^-\Lambda\Xi^+ + \text{c.c.}$

 $\Gamma(K^-\Sigma^0\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{40}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.67±0.33±0.28	142	ABLIKIM	15I	$e^+e^- \rightarrow \psi(2S) \rightarrow K^-\Sigma^0\Xi^+ + \text{c.c.}$

 $\Gamma(\Omega^-\bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{41}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.47±0.09±0.05		27	¹ DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	90	ABLIKIM	12Q	BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<1.6	90	PEDLAR	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<0.73	90	² BAI	01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

 $\Gamma(\pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{42}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.53±0.07 OUR AVERAGE				
1.65±0.03±0.15	4.5k	ABLIKIM	13A	$\psi(2S) \rightarrow p\bar{p}\pi^0$
1.54±0.06±0.06	948	ALEXANDER	10	$\psi(2S) \rightarrow \pi^0 p\bar{p}$
1.32±0.10±0.15	256	¹ ABLIKIM	05E	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
1.4 ± 0.5	9	FRANKLIN	83	MRK2 e^+e^-

¹ Computed using $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$.

 $\Gamma(N(940)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{43}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.42±0.20^{+1.78}_{-1.28}	1.9k	¹ ABLIKIM	13A	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{44}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

7.3 ± 1.7 OUR AVERAGE Error includes scale factor of 2.5.

3.58 ± 0.25 ± 1.59 ± 0.84	1.1k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
8.1 ± 0.7 ± 0.3	474	² ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.² From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances. $\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{45}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.64 ± 0.05 ± 0.22 ± 0.17	0.2k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states. $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{46}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.47 ± 0.28 ± 0.99 ± 0.97	0.7k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states. $\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{47}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.76 ± 0.28 ± 1.37 ± 1.66	1.1k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states. $\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{48}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.79 ± 0.10 ± 0.24 ± 0.71	0.5k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states. $\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{49}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.62 ± 0.28 ± 1.12 ± 0.64	0.9k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states. $\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{50}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.13 ± 0.08 ± 0.40 ± 0.30	0.8k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.4 \pm 0.1$	76	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \pi^0 p\bar{p}$

¹ From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N_1^*(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

 $\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.4 OUR AVERAGE				
6.4 $\pm 0.2 \pm 0.6$	679	¹ ABLIKIM	13S	BES3 $\psi(2S) \rightarrow \eta p\bar{p}$
5.6 $\pm 0.6 \pm 0.3$	154	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \eta p\bar{p}$
5.8 $\pm 1.1 \pm 0.7$	44.8 ± 8.5	² ABLIKIM	05E	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
8 ± 3 ± 3	9.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+ \pi^- \pi^0$

¹ With $N(1535)$ decaying to $p\eta$.

² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.

 $\Gamma(\eta f_0(2100) \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.2 \pm 0.4 \pm 0.1$	31	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \eta p\bar{p}$

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

 $\Gamma(N(1535)\bar{p} \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 0.6 \pm 0.3$	123	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \eta p\bar{p}$

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

 $\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.69 ± 0.21 OUR AVERAGE				
0.6 ± 0.2 ± 0.2	21.2	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+ \pi^- \pi^0$
0.8 ± 0.3 ± 0.1	14.9 ± 0.1	¹ BAI	03B	BES $\psi(2S) \rightarrow p\bar{p}\pi^+ \pi^- \pi^0$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26	90	¹ BAI	03B	BES $\psi(2S) \rightarrow K^+ K^- p\bar{p}$
-------	----	------------------	-----	---

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+\pi^- p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.0 ± 0.4 OUR AVERAGE				
$5.9 \pm 0.2 \pm 0.4$	904.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$
8 ± 2		¹ TANENBAUM	78	MRK1 $e^+ e^-$

¹ Assuming entirely strong decay. $\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.48 ± 0.17 OUR AVERAGE				
$2.45 \pm 0.11 \pm 0.21$	851	ABLIKIM	06I	BES2 $e^+ e^- \rightarrow p\pi^- X$
$2.52 \pm 0.12 \pm 0.22$	849	ABLIKIM	06I	BES2 $e^+ e^- \rightarrow \bar{p}\pi^+ X$

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.18 \pm 0.50 \pm 0.50$	135 ± 21	ABLIKIM	06I	BES2 $e^+ e^- \rightarrow p\pi^-\pi^0 X$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.6	90	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.5 \pm 0.7 \pm 1.5$		¹ BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.3 \pm 0.8 \pm 1.4 201.7 2 BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi (\eta \rightarrow \gamma\gamma)$ 8.1 \pm 1.4 \pm 1.6 50.0 2 BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi (\eta \rightarrow 3\pi)$ ¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.² Not independent from other values reported by BRIERE 05. $\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.2 \pm 0.6 \pm 0.1$	16	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$


¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$. $\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.5 \pm 1.6 \pm 1.3$	12.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±1.2 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.				
8.4±0.5±1.2	386	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
12.2±2.2±0.7	37	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
8.2±0.5±0.7	391	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
4.8±0.6±0.7	100 ± 22	² BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16 \text{ eV}$.				
² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

Γ_{66}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ±0.6 OUR AVERAGE Error includes scale factor of 1.1.				
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
4.18 ^{+0.43} _{-0.42} ± 0.92	170	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	^{1,2} BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ± 0.8 ± 1.0		¹ BAI	99C BES	Repl. by BAI 03B

¹ Assuming $B(b_1 \rightarrow \omega\pi)=1$.

² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{67}/Γ
$2.35^{+0.47}_{-0.42} \pm 0.40$	45	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$	

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{68}/Γ
2.2 ± 0.4 OUR AVERAGE						
$2.3 \pm 0.5 \pm 0.4$		57	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$	
$2.05 \pm 0.41 \pm 0.38$		62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
<1.5	90	¹ BAI	03B BES		$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
<1.7	90	BAI	98J BES		Repl. by BAI 03B	

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{70}/Γ
7.3 ± 0.5 OUR AVERAGE					
$8.1 \pm 1.3 \pm 0.3$	133	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	
$7.1 \pm 0.3 \pm 0.4$	817.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	
16 ± 4	1 TANENBAUM 78	MRK1	¹ TANENBAUM 78 MRK1	$e^+ e^-$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$10.9 \pm 1.9 \pm 0.2$	85	² AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	

¹ Assuming entirely strong decay.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{71}/Γ
$2.2 \pm 0.2 \pm 0.4$	223.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{72}/Γ
1.86 $\pm 0.32 \pm 0.43$	93 ± 16	BAI	04C		$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
<1.2	90	BAI	98J BES		$e^+ e^-$	

$\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{73}/Γ
1.3 $\pm 0.7 \pm 0.1$	7	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$	
¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$ eV.					

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.0±2.5±1.8	65	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.0±1.8±2.1	¹ BAI	99C	BES	$e^+ e^-$

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.20±0.25±0.37	83 ± 9	ABLIKIM	050 BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\rho^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.5±0.1±0.2	61.1	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{79}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.7±2.5		TANENBAUM 78	MRK1	$e^+ e^-$

 $\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{80}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.4±0.6 OUR AVERAGE		Error includes scale factor of 2.2.		
$2.2 \pm 0.2 \pm 0.2$	308	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.5 ± 1.0		TANENBAUM 78	MRK1	$e^+ e^-$

 $\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{81}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.2±0.6 OUR AVERAGE		Error includes scale factor of 1.4.		
$2.0 \pm 0.2 \pm 0.4$	285.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.2 ± 1.5		TANENBAUM 78	MRK1	$e^+ e^-$

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.6±0.9 OUR AVERAGE				
$18.8 \pm 5.7 \pm 0.3$	32	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$
$11.7 \pm 1.0 \pm 1.5$	597	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$12.7 \pm 0.5 \pm 1.0$	711.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{83}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.9±2.0±0.9	19	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{84}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.6±1.3±1.8	238	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{85}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.6±2.2±1.7	133	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{86}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.3±2.2±1.4	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{87}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1±1.3±1.2	125	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(\eta K^+ K^-, \text{no } \eta\phi)/\Gamma_{\text{total}}$ Γ_{88}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.08±0.29±0.25	0.3k	1	ABLIKIM	12L BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13	90	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-----	----	--------	----	------	--

1 Excluding $\eta\phi$.

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{89}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.62±0.11 OUR AVERAGE				Error includes scale factor of 1.1.
1.56±0.04±0.11	2.8k	ABLIKIM	14G BES3	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
2.38±0.37±0.29	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.9 ± 0.3 ± 0.3	76.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.5 ± 0.3 ± 0.2	23	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

1 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\omega K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{90}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.7±2.6 OUR AVERAGE				
18.9±2.9±2.2	396	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
22.6±3.0±2.4	535	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

$\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{91}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1 ± 1.2 OUR AVERAGE				
6.39 ± 1.50 ± 0.78	128	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
5.86 ± 1.61 ± 0.83	143	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

$\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$ Γ_{92}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.8 ± 2.5 ± 1.6	356	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

$\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$ Γ_{93}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.82 ± 2.08 ± 0.72	116	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

$\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{94}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.60 ± 0.27 ± 0.24	109	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ $X(1440)$ compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{95}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.09 ± 0.20 ± 0.16	82	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ $X(1440)$ compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{96}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.302 ± 0.098 ± 0.027	22	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ Statistical significance 4.5 σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.

$\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{97}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.125 ± 0.070 ± 0.013	10	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ Statistical significance 3.2 σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{98}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.5 ± 2.0 OUR AVERAGE				Error includes scale factor of 2.8.

5.45 ± 0.42 ± 0.87 671 ABLIKIM 05H BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$

1.5 ± 1.0 ¹ TANENBAUM 78 MRK1 $e^+ e^-$

¹ Assuming entirely strong decay.

$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_{99}/Γ
$7.3 \pm 0.4 \pm 0.6$	434.9

DOCUMENT ID *TECN* *COMMENT*

BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$	Γ_{100}/Γ
$7.48 \pm 0.23 \pm 0.39$	$1.3k$

DOCUMENT ID *TECN* *COMMENT*

¹ METREVELI 12 $\psi(2S) \rightarrow K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.2 \pm 1.5 \pm 0.2$	66	^{2,3} LEES	$15J$	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$8.3 \pm 1.5 \pm 0.2$	66	^{3,4} LEES	$15J$	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$6.3 \pm 0.6 \pm 0.3$		⁵ DOBBS	$06A$	CLEO	e^+e^-
10 ± 7		⁵ BRANDELIK	$79C$	DASP	e^+e^-
< 5	90	FELDMAN	77	MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+e^-) = (2.37 \pm 0.04)$ keV.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant K^+K^- production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$	Γ_{101}/Γ
5.34 ± 0.33 OUR AVERAGE	

DOCUMENT ID *TECN* *COMMENT*

¹ METREVELI 12 $\psi(2S) \rightarrow K_S^0 K_L^0$

DOBBS 06A CLEO e^+e^-

² BAI 04B BES2 $\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_{102}/Γ
2.01 ± 0.17 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.

DOCUMENT ID *TECN* *COMMENT*

¹ ABLIKIM 12H BES3 $e^+e^- \rightarrow \psi(2S)$


² ABLIKIM 05J BES2 $e^+e^- \rightarrow \psi(2S)$

ADAM 05 CLEO $e^+e^- \rightarrow \psi(2S)$

FRANKLIN 83 MRK2 $e^+e^- \rightarrow \text{hadrons}$

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.

² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.



$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{103}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.94 \pm 0.25^{+1.15}_{-0.34}$	¹ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{104}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.12 OUR AVERAGE					Error includes scale factor of 1.8.
$0.51 \pm 0.07 \pm 0.11$			¹ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
$0.24^{+0.08}_{-0.07} \pm 0.02$		22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83	MRK2	e^+e^-
<10	90		BARTEL	76	CNTR	e^+e^-
<10	90		² ABRAMS	75	MRK1	e^+e^-

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

² Final state $\rho^0\pi^0$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{105}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.26 OUR AVERAGE					
$0.76 \pm 0.25 \pm 0.06$		30	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-$
8 ± 5			BRANDELIK	79C DASP	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	DOBBS	06A	CLEO	$e^+ e^- \rightarrow \psi(2S)$
<5	90	FELDMAN	77	MRK1	$e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+ \pi^-$ for continuum subtraction.

$\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	1 BAI	99c	$e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.12 \pm 0.62^{+1.13}_{-0.61}$	251 ± 22	ABLIKIM	12L	$BES3$ $e^+ e^- \rightarrow \psi(2S)$

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.07 \pm 0.16 \pm 0.26$	0.9k	ABLIKIM	12L	$BES3$	$e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.9	90	1	FRANKLIN	83	MRK2	$e^+ e^- \rightarrow$ hadrons
------	----	---	----------	----	------	-------------------------------

$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR AVERAGE			Error includes scale factor of 1.2.		
$3.18 \pm 0.30^{+0.26}_{-0.31}$	0.2k	ABLIKIM	12L	$BES3$	$e^+ e^- \rightarrow \psi(2S)$
$2.9^{+1.3}_{-1.7} \pm 0.4$	9.6 ± 4.2	ABLIKIM	05I	$BES2$	$e^+ e^- \rightarrow \psi(2S)$
$1.3^{+1.0}_{-0.7} \pm 0.3$	7	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.4	90	FRANKLIN	83	MRK2	$e^+ e^- \rightarrow$ hadrons
------	----	----------	----	------	-------------------------------

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ± 2.0 OUR AVERAGE				
$13.3^{+2.4}_{-2.8} \pm 1.7$	65.6 ± 9.0	ABLIKIM	05I	$BES2$ $e^+ e^- \rightarrow \psi(2S)$
$9.2^{+2.7}_{-2.2} \pm 0.9$	25	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$

$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$ $\Gamma_{109}/\Gamma_{110}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.06 OUR AVERAGE			
$0.22^{+0.10}_{-0.14}$	ABLIKIM	05I	$BES2$ $e^+ e^- \rightarrow \psi(2S)$
$0.14^{+0.08}_{-0.06}$	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{111}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.18±0.26 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
2.3 ± 0.8 ± 0.1	19 ± 6	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.9 ± 0.2 ± 0.1	47.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
1.5 ± 0.2 ± 0.2	51.5 ± 8.3	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.44±0.96±0.04	10 ± 4	^{2,3} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+K^-) = (49.3 \pm 0.6)\%$.

WEIGHTED AVERAGE
1.18±0.26 (Error scaled by 1.5)



$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{112}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75±0.33 OUR AVERAGE				Error includes scale factor of 1.6.
1.5 ± 0.5 ± 0.1	12 ± 4	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.6 ± 0.2 ± 0.1	18.4 ± 6.4	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.45±0.70±0.02	6 ± 3	^{2,3} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.34 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+K^-) = (49.3 \pm 0.6)\%$.

$\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{113}/Γ
$0.63 \pm 0.13 \text{ OUR AVERAGE}$					
0.9 ± 0.4 ± 0.1	13	LEES	12F	BABR $e^+e^- \rightarrow 2(K^+K^-)\gamma$	
0.6 ± 0.1 ± 0.1	59.2	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(K^+K^-)$	

$\Gamma(\phi K^+K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{114}/Γ
$0.70 \pm 0.16 \text{ OUR AVERAGE}$					
0.8 ± 0.2 ± 0.1	36.8	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(K^+K^-)$	
0.6 ± 0.2 ± 0.1	16.1 \pm 5.0	¹ BAI	03B	BES $\psi(2S) \rightarrow 2(K^+K^-)$	

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(2(K^+K^-)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{115}/Γ
$1.1 \pm 0.2 \pm 0.2$					
1.1 ± 0.2 ± 0.2	44.7	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(K^+K^-)\pi^0$	

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{116}/Γ
$3.10 \pm 0.31 \text{ OUR AVERAGE}$					
3.14 ± 0.23 ± 0.23	0.2k	ABLIKIM	12L	BES3 $e^+e^- \rightarrow \psi(2S)$	
2.0 $^{+1.5}_{-1.1}$ ± 0.4	6	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$	
3.3 ± 1.1 ± 0.5	17	ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$	

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{117}/Γ
$3.1 \pm 1.4 \pm 0.7$					
3.1 ± 1.4 ± 0.7	8	¹ ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$	

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

$\Gamma(\omega\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{118}/Γ
$3.2^{+2.4}_{-2.0} \pm 0.7$					
3.2 $^{+2.4}_{-2.0}$ ± 0.7	4	¹ ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$	

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS
--------------------------	------

 2.1 ± 0.6 OUR AVERAGE

$2.5 \begin{array}{l} +1.2 \\ -1.0 \end{array}$	± 0.2	14
---	-----------	----

$1.87 \begin{array}{l} +0.68 \\ -0.62 \end{array}$	± 0.28	14
--	------------	----

 Γ_{119}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$
------	----	-------------------------------------

ABLIKIM	04L	BES $e^+ e^- \rightarrow \psi(2S)$
---------	-----	------------------------------------

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS
--------------------------	------

$1.87 \begin{array}{l} +1.64 \\ -1.11 \end{array}$	± 0.33
--	------------

 Γ_{120}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

ABLIKIM	04L	BES $e^+ e^- \rightarrow \psi(2S)$
---------	-----	------------------------------------

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS
--------------------------	------

 2.2 ± 0.6 OUR AVERAGE

Error includes scale factor of 1.1.

$3.0 \begin{array}{l} +1.1 \\ -0.9 \end{array}$	± 0.2	18
---	-----------	----

$1.78 \begin{array}{l} +0.67 \\ -0.62 \end{array}$	± 0.17	13
--	------------	----

 Γ_{121}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$
------	----	-------------------------------------

ABLIKIM	04L	BES $e^+ e^- \rightarrow \psi(2S)$
---------	-----	------------------------------------

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%
--------------------------	-----

<1.1	90
------	----

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1	90
------	----

 Γ_{122}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$
------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1	90
------	----

 $\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%
--------------------------	-----

<0.04	90
-------	----

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90
------	----

<0.4	90
------	----

 Γ_{123}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

ABLIKIM	12L	BES3 $e^+ e^- \rightarrow \psi(2S)$
---------	-----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90
------	----

<0.4	90
------	----

 $\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%
--------------------------	-----

<1.0	90
------	----

 Γ_{124}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

PEDLAR	07	CLEO $e^+ e^- \rightarrow \psi(2S)$
--------	----	-------------------------------------

 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS
--------------------------	------

$2.7 \pm 0.6 \pm 0.4$	30.1
-----------------------	------

 Γ_{125}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
--------	----	--

 $\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
--------------------------	------

$0.81 \pm 0.11 \pm 0.14$	50
--------------------------	----

 Γ_{126}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

¹ ABLIKIM	08C	BES2 $e^+ e^- \rightarrow J/\psi$
----------------------	-----	-----------------------------------

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$				Γ_{127}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.44 ± 0.12 ± 0.11		20 ± 6	BAI	04C	$\psi(2S) \rightarrow 2(K^+ K^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.45	90	BAI	98J	BES	$e^+ e^- \rightarrow 2(K^+ K^-)$

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{128}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.88	90	BAI	04G	BES2

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$				Γ_{129}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.0	90	BAI	04G	BES2

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$				Γ_{130}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.70	90	BAI	04G	BES2

$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$				Γ_{131}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6	90	BAI	04G	BES2

$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$				Γ_{132}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.60	90	BAI	04G	BES2

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{133}/Γ
<u>VALUE (units 10^{-4})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.046		¹ BAI	04D	BES

¹ Forbidden by CP.

———— RADIATIVE DECAYS ——

$\Gamma(\gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$				Γ_{134}/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.99 ± 0.27 OUR FIT				
9.2 ± 0.4 OUR AVERAGE				
9.22 ± 0.11 ± 0.46	72600	ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
9.9 ± 0.5 ± 0.8		¹ GAISER	86	CBAL $e^+ e^- \rightarrow \gamma X$
7.2 ± 2.3		¹ BIDDICK	77	CNTR $e^+ e^- \rightarrow \gamma X$
7.5 ± 2.6		¹ WHITAKER	76	MRK1 $e^+ e^-$

¹ Angular distribution ($1+\cos^2\theta$) assumed.

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{135}/Γ

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.55±0.31 OUR FIT				
8.9 ±0.5 OUR AVERAGE				
9.07±0.11±0.54	76700	ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
9.0 ±0.5 ±0.7		¹ GAISER	86	CBAL $e^+ e^- \rightarrow \gamma X$
7.1 ±1.9		² BIDDICK	77	CNTR $e^+ e^- \rightarrow \gamma X$

¹ Angular distribution ($1 - 0.189 \cos^2\theta$) assumed.² Valid for isotropic distribution of the photon. $\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_{136}/Γ

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.11±0.31 OUR FIT				
8.8 ±0.5 OUR AVERAGE				Error includes scale factor of 1.1.
9.33±0.14±0.61	79300	ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
8.0 ±0.5 ±0.7		¹ GAISER	86	CBAL $e^+ e^- \rightarrow \gamma X$
7.0 ±2.0		² BIDDICK	77	CNTR $e^+ e^- \rightarrow \gamma X$

¹ Angular distribution ($1 - 0.052 \cos^2\theta$) assumed.² Valid for isotropic distribution of the photon.

$$[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))] / \Gamma_{\text{total}} \quad (\Gamma_{134} + \Gamma_{135} + \Gamma_{136}) / \Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

27.6±0.3±2.0	¹ ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
--------------	--------------------	----	-------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$. $\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{134}/\Gamma_{135}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.02±0.01±0.07	¹ ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
----------------	--------------------	----	-------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$. $\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{136}/\Gamma_{135}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.03±0.02±0.03	¹ ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
----------------	--------------------	----	-------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$. $\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$ $\Gamma_{134}/\Gamma_{136}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.99±0.02±0.08	¹ ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
----------------	--------------------	----	-------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$


Γ_{137}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.432 ± 0.016 ± 0.060		MITCHELL 09	CLEO	$e^+ e^- \rightarrow \gamma X$
0.32 ± 0.04 ± 0.06	2560	1 ATHAR 04	CLEO	$e^+ e^- \rightarrow \gamma X$
0.28 ± 0.06		2 GAISER 86	CBAL	$e^+ e^- \rightarrow \gamma X$

¹ ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

² GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.

WEIGHTED AVERAGE
0.34±0.05 (Error scaled by 1.3)



$$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}} (\text{units } 10^{-2})$$

$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$

Γ_{138}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
7±2±4		1 ABLIKIM 12G	BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, KK\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8	90	2 CRONIN-HENNESSY 10	CLEO	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$
< 20	90	ATHAR 04	CLEO	$e^+ e^- \rightarrow \gamma X$
20–130	95	EDWARDS 82C	CBAL	$e^+ e^- \rightarrow \gamma X$

¹ ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$

Γ_{139}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.58 \pm 0.40 \pm 0.13$		37	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 5	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 5400	95	1 LIBERMAN	75	SPEC	e^+e^-
$< 1 \times 10^4$	90	WIIK	75	DASP	e^+e^-

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.0077$.

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

Γ_{140}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.23 ± 0.06 OUR AVERAGE					
1.26 $\pm 0.03 \pm 0.08$		2226	¹ ABLIKIM	10F BES3	$\psi(2S) \rightarrow 3\gamma\pi^+\pi^-$, $2\gamma\pi^+\pi^-$
1.19 $\pm 0.08 \pm 0.03$			PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
1.24 $\pm 0.27 \pm 0.15$		23	ABLIKIM	06R BES2	$e^+e^- \rightarrow \psi(2S)$
1.54 $\pm 0.31 \pm 0.20$		~ 43	BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^-2\gamma$, $\pi^+\pi^-3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60	90	² BRAUNSCH...	77	DASP	e^+e^-
< 11	90	³ BARTEL	76	CNTR	e^+e^-

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

Γ_{141}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.73^{+0.29}_{-0.25}$ OUR AVERAGE				Error includes scale factor of 1.8.
2.84 $\pm 0.15^{+0.03}_{-0.10}$	1.9k	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
2.12 $\pm 0.19 \pm 0.32$		^{3,4} BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.08 $\pm 0.19 \pm 0.33$	200.6 ± 18.8	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
2.90 $\pm 1.08 \pm 1.07$	29.9 ± 11.1	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{142}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$3.1 \pm 1.0 \pm 1.4$	175	¹ DOBBS	15 $\psi(2S) \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
$9.2 \pm 1.8 \pm 0.6$	274	1,2 DOBBS	15 $\psi(2S) \rightarrow \gamma\pi\pi$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi\pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi\pi) = (34.9 \pm 2.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
$3.3 \pm 0.8 \pm 0.1$	136	1,2 DOBBS	15 $\psi(2S) \rightarrow \gamma K\bar{K}$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ± 0.6 OUR AVERAGE				
3.6 ± 0.4 ± 0.5	290	1 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
$3.01 \pm 0.41 \pm 1.24$	35.6 ± 4.8	2 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.6 ± 0.7 OUR AVERAGE					
6.7 ± 0.6 ± 0.6		375	1 DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$
$6.04 \pm 0.90 \pm 1.32$		39.6 ± 5.9	2,3 BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 15.6	90	6.8 ± 3.1	2,3 BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Includes unknown branching fractions to K^+K^- or $K_S^0 K_S^0$. We have multiplied the K^+K^- result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K\bar{K}$ result.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	COMMENT
$4.8 \pm 0.5 \pm 0.9$	373	1 DOBBS	15 $\psi(2S) \rightarrow \gamma\pi\pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	COMMENT
$3.2 \pm 0.6 \pm 0.8$	207	1 DOBBS	15 $\psi(2S) \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<5.8 \times 10^{-6}$	90	1,2 DOBBS	$\psi(2S) \rightarrow \gamma\pi\pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<9.5 \times 10^{-6}$	90	1,2 DOBBS	$\psi(2S) \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for K^+K^- and $K_S^0K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.38 \pm 0.48 \pm 0.09$	13	1	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$, $\gamma 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 90	90	BAI	98F	BES	$\psi(2S) \rightarrow \pi^+\pi^-3\gamma$
<200	90	YAMADA	77	DASP	$e^+e^- \rightarrow 3\gamma$

¹ Combining the results from $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

 $\Gamma(\gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.71 \pm 1.25 \pm 1.64$	418	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
<1.2	90	¹ SCHARRE	80 MRK1	$e^+e^- \rightarrow K\bar{K}\pi$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

 $\Gamma(\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.36 \pm 0.25 \pm 0.05$	10	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma\eta(1475) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$
------	----	---------	----------	---

$\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{160}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.88	90	ABLIKIM	06R	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{161}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
39.6±2.8±5.0	583	ABLIKIM	07D	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^{*0}K^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{162}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
37.0±6.1±7.2	237	ABLIKIM	07D	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^{*0}\bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{163}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
24.0±4.5±5.0	41	ABLIKIM	07D	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K_S^0K^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{164}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.6±3.6±3.6	115	ABLIKIM	07D	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{165}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
19.1±2.7±4.3	132	ABLIKIM	07D	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{166}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.9 ± 0.5 OUR AVERAGE				Error includes scale factor of 2.0.
4.18±0.26±0.18	348	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$
2.9 ± 0.4 ± 0.4	142	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p} < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{167}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.2±0.2±0.1	111	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p} < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{168}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.72±0.18±0.03	73	¹ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p} < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

Γ_{169}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.57 \pm 0.36^{+1.77}_{-4.26}$		ABLIKIM	12D	$BES3$ $J/\psi \rightarrow \gamma p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
<5.4	90	ABLIKIM	07D	BES	$\psi(2S) \rightarrow \gamma p\bar{p}$

$\Gamma(\gamma X \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

Γ_{170}/Γ

For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2	90	ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

$\Gamma(\gamma\pi^+\pi^-p\bar{p})/\Gamma_{\text{total}}$

Γ_{171}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.8 \pm 1.2 \pm 0.7$	17	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma 2(\pi^+\pi^-)K^+K^-)/\Gamma_{\text{total}}$

Γ_{172}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<22	90	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{173}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<17	90	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^+K^-K^+K^-)/\Gamma_{\text{total}}$

Γ_{174}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma\gamma J/\psi)/\Gamma_{\text{total}}$

Γ_{175}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.1 \pm 0.6^{+0.8}_{-1.0}$	1.1k	ABLIKIM	120	BES3 $e^+e^- \rightarrow \psi(2S)$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+e^-)$

Γ_{176}/Γ_6

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.0	90	LEES	13I	BABR $B \rightarrow K^{(*)}\psi(2S)$

$\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$
see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

$a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
67⁺¹⁹₋₁₃	59k	¹ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
37⁺⁵³₋₄₇	59k	¹ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

$\psi(2S)$ REFERENCES

ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14G	PR D89 112006	M. Ablikim <i>et al.</i>	(BES III Collab.)
DOBBS	14	PL B739 90	S. Dobbs <i>et al.</i>	(NWES, WAYN)
ABLIKIM	13A	PRL 110 022001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13S	PR D88 032010	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13W	PR D88 112007	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12L	PR D86 072011	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12M	PR D86 092008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12Q	CP C36 1040	M. Ablikim <i>et al.</i>	(BES II Collab.)
ANASHIN	12	PL B711 280	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BES III Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... ¹⁰	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)

PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
		Translated from ZETFP 85 429.		
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Fermilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
GРИBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)

COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC
WIJK	75	Stanford Symp. 69	B.H. Wiik	(DESY)
