

$\psi(4040)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(4040)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4039 ± 1 OUR ESTIMATE			
4039.6 ± 4.3	¹ ABLIKIM 08D BES2 $e^+ e^- \rightarrow$ hadrons		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4034 ± 6	² MO 10 RVUE $e^+ e^- \rightarrow$ hadrons		
4037 ± 2	³ SETH 05A RVUE $e^+ e^- \rightarrow$ hadrons		
4040 ± 1	⁴ SETH 05A RVUE $e^+ e^- \rightarrow$ hadrons		
4040 ± 10	BRANDELIK 78C DASP $e^+ e^-$		
¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
³ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁴ From a fit to BES (BAI 02C) data.			

$\psi(4040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 ± 10 OUR ESTIMATE			
84.5 ± 12.3	⁵ ABLIKIM 08D BES2 $e^+ e^- \rightarrow$ hadrons		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87 ± 11	⁶ MO 10 RVUE $e^+ e^- \rightarrow$ hadrons		
85 ± 10	⁷ SETH 05A RVUE $e^+ e^- \rightarrow$ hadrons		
89 ± 6	⁸ SETH 05A RVUE $e^+ e^- \rightarrow$ hadrons		
52 ± 10	BRANDELIK 78C DASP $e^+ e^-$		
⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
⁶ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁷ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁸ From a fit to BES (BAI 02C) data.			

$\psi(4040)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+ e^-$	$(1.07 \pm 0.16) \times 10^{-5}$	
$\Gamma_2 D\bar{D}$	seen	
$\Gamma_3 D^0\bar{D}^0$	seen	
$\Gamma_4 D^+D^-$	seen	
$\Gamma_5 D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_6 D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_7 D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_8 D^*\bar{D}^*$	seen	
$\Gamma_9 D^*(2007)^0\bar{D}^*(2007)^0$	seen	
$\Gamma_{10} D^*(2010)^+D^*(2010)^-$	seen	
$\Gamma_{11} D\bar{D}\pi (\text{excl. } D^*\bar{D})$		
$\Gamma_{12} D^0 D^- \pi^+ + \text{c.c.} (\text{excl. } D^*(2007)^0\bar{D}^0 + \text{c.c.}, D^*(2010)^+D^- + \text{c.c.})$	not seen	
$\Gamma_{13} D\bar{D}^*\pi (\text{excl. } D^*\bar{D}^*)$	not seen	
$\Gamma_{14} D^0\bar{D}^{*-}\pi^+ + \text{c.c.} (\text{excl. } D^*(2010)^+D^*(2010)^-)$	seen	
$\Gamma_{15} D_s^+ D_s^-$	seen	
$\Gamma_{16} J/\psi(1S)\text{hadrons}$		
$\Gamma_{17} J/\psi\pi^+\pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{18} J/\psi\pi^0\pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{19} J/\psi\eta$	$(5.2 \pm 0.7) \times 10^{-3}$	
$\Gamma_{20} J/\psi\pi^0$	$< 2.8 \times 10^{-4}$	90%
$\Gamma_{21} J/\psi\pi^+\pi^-\pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{22} \chi_{c1}\gamma$	$< 3.4 \times 10^{-3}$	90%
$\Gamma_{23} \chi_{c2}\gamma$	$< 5 \times 10^{-3}$	90%
$\Gamma_{24} \chi_{c1}\pi^+\pi^-\pi^0$	$< 1.1 \%$	90%
$\Gamma_{25} \chi_{c2}\pi^+\pi^-\pi^0$	$< 3.2 \%$	90%
$\Gamma_{26} h_c(1P)\pi^+\pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{27} \phi\pi^+\pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{28} \Lambda\bar{\Lambda}\pi^+\pi^-$	$< 2.9 \times 10^{-4}$	90%
$\Gamma_{29} \Lambda\bar{\Lambda}\pi^0$	$< 9 \times 10^{-5}$	90%
$\Gamma_{30} \Lambda\bar{\Lambda}\eta$	$< 3.0 \times 10^{-4}$	90%
$\Gamma_{31} \Sigma^+\bar{\Sigma}^-$	$< 1.3 \times 10^{-4}$	90%
$\Gamma_{32} \Sigma^0\bar{\Sigma}^0$	$< 7 \times 10^{-5}$	90%

Γ_{33}	$\Xi^+\Xi^-$	< 1.6	$\times 10^{-4}$	90%
Γ_{34}	$\Xi^0\Xi^0$	< 1.8	$\times 10^{-4}$	90%
Γ_{35}	$\mu^+ \mu^-$			

$\psi(4040)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$	Γ_1		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.86±0.07 OUR ESTIMATE			
0.83±0.20	9 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.6 to 1.4	10 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
0.88 ± 0.11	11 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.91 ± 0.13	12 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.75 ± 0.15	BRANDELIK	78C DASP	$e^+ e^-$
9 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
10 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.			
11 From a fit to Crystal Ball (OSTERHELD 86) data.			
12 From a fit to BES (BAI 02C) data.			

$\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{22}\Gamma_1/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN
<2.9	90	13 HAN	15 BELL
13 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.			
$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{23}\Gamma_1/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN
<4.6	90	14 HAN	15 BELL
14 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.			

$\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{19}/\Gamma \times \Gamma_1/\Gamma$		
VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.1±1.4±1.5	15 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$
12.8±2.1±1.9	16 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$
15 Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.			
16 Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.			

$\psi(4040)$ BRANCHING RATIOS **$\Gamma(e^+e^-)/\Gamma_{\text{total}}$**

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_1/Γ</u>
---------------------------------	--------------------	-------------	----------------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0

FELDMAN 77 MRK1 e^+e^-

 $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_3/Γ</u>
seen	AUBERT 09M	BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$	
seen	PAKHLOVA 08	BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$	

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_4/Γ</u>
seen	AUBERT 09M	BABR	$e^+e^- \rightarrow D^+D^-\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$	
seen	PAKHLOVA 08	BELL	$e^+e^- \rightarrow D^+D^-\gamma$	

 $\Gamma(D\bar{D})/\Gamma(D^*\bar{D} + \text{c.c.})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_2/Γ_5</u>
0.24±0.05±0.12	AUBERT 09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}$	

 $\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_3/Γ_6</u>
0.05±0.03	17 GOLDHABER 77	MRK1	e^+e^-	

17 Phase-space factor (p^3) explicitly removed.

 $\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_6/Γ</u>
seen	AUBERT 09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$	

 $\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_7/Γ</u>
seen	AUBERT 09M	BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^-$	
seen	PAKHLOVA 07	BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$	

 $\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_7/Γ_6</u>
0.95±0.09±0.10	AUBERT 09M	BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}$	

 $\Gamma(D^*\bar{D}^*)/\Gamma(D^*\bar{D} + \text{c.c.})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_8/Γ_5</u>
0.18±0.14±0.03	AUBERT 09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	

$$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$

$$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.}) \quad \Gamma_9/\Gamma_6$$

VALUE	DOCUMENT ID	TECN	COMMENT
32.0 ± 12.0	18 GOLDHABER 77	MRK1	$e^+ e^-$

18 Phase-space factor (p^3) explicitly removed.

$$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

$$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.})) / \Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

$$\Gamma(D \bar{D}^* \pi (\text{excl. } D^* \bar{D}^*))/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

$$\Gamma(D^0 \bar{D}^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 \bar{D}^{*-} \pi^+ \gamma$

$$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

$$\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$

Γ_{19}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
5.2±0.5±0.5		¹⁹ ABLIKIM	12K BES3	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

¹⁹ ABLIKIM 12K measure $\sigma(e^+e^- \rightarrow J/\psi\eta) = 32.1 \pm 2.8 \pm 1.3$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$

Γ_{20}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.28	90	²⁰ ABLIKIM	12K BES3	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

²⁰ ABLIKIM 12K measure $\sigma(e^+e^- \rightarrow J/\psi\pi^0) < 1.6$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.

$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{21}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$

Γ_{22}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

<11 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$

Γ_{23}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

<17 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{24}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<32	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	²¹ PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

²¹ From several values of \sqrt{s} near the peak of the $\psi(4040)$, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 1.0 \pm 8.0 \pm 5.4 \pm 0.2$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{27}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{28}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.9	90	22 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

22 Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$	Γ_{29}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.9	90	23 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

23 Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$	Γ_{30}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.0	90	24 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

24 Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$	Γ_{31}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	25 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

25 Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$	Γ_{32}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.7	90	26 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

26 Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$	Γ_{33}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.6	90	27 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

27 Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$	Γ_{34}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.8	90	28 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

28 Assuming that interference effects between resonance and continuum can be neglected.

$\psi(4040)$ REFERENCES

HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
ABLIKIM	13Q	PR D87 112011	Ablikim M. <i>et al.</i>	(BES III Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12K	PR D86 071101	M. Ablikim <i>et al.</i>	(BES III Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
Also		ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)