

**$\Xi(1820)$**  $I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$  Status: \*\*\*

The clearest evidence is an 8-standard-deviation peak in  $\Lambda K^-$  seen by GAY 76C. TEODORO 78 favors  $J = 3/2$ , but cannot make a parity discrimination. BIAGI 87C is consistent with  $J = 3/2$  and favors negative parity for this  $J$  value.

 **$\Xi(1820)$  MASS**

We only average the measurements that appear to us to be most significant and best determined.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1823 ± 5 OUR ESTIMATE</b>					
<b>1823.4 ± 1.4 OUR AVERAGE</b>					
1819.4 ± 3.1 ± 2.0	280	<sup>1</sup> BIAGI	87	SPEC 0	$\Xi^-$ Be → $(\Lambda K^-)$ X
1826 ± 3 ± 1	54	BIAGI	87C	SPEC 0	$\Xi^-$ Be → $(\Lambda \bar{K}^0)$ X
1822 ± 6		JENKINS	83	MPS —	$K^- p \rightarrow K^+$ (MM)
1830 ± 6	300	BIAGI	81	SPEC —	SPS hyperon beam
1823 ± 2	130	GAY	76C	HBC —	$K^- p$ 4.2 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1817 ± 3		ADAMOVICH	99B	WA89	$\Sigma^-$ nucleus, 345 GeV
1797 ± 19	74	BRIEFEL	77	HBC 0	$K^- p$ 2.87 GeV/c
1829 ± 9	68	BRIEFEL	77	HBC —0	$\Xi(1530)\pi$
1860 ± 14	39	BRIEFEL	77	HBC —	$\Sigma^- \bar{K}^0$
1870 ± 9	44	BRIEFEL	77	HBC 0	$\Lambda \bar{K}^0$
1813 ± 4	57	BRIEFEL	77	HBC —	$\Lambda K^-$
1807 ± 27		DIBIANCA	75	DBC —0	$\Xi \pi \pi$ , $\Xi^* \pi$
1762 ± 8	28	<sup>2</sup> BADIER	72	HBC —0	$\Xi \pi$ , $\Xi \pi \pi$ , YK
1838 ± 5	38	<sup>2</sup> BADIER	72	HBC —0	$\Xi \pi$ , $\Xi \pi \pi$ , YK
1830 ± 10	25	<sup>3</sup> CRENNELL	70B	DBC —0	3.6, 3.9 GeV/c
1826 ± 12		<sup>4</sup> CRENNELL	70B	DBC —0	3.6, 3.9 GeV/c
1830 ± 10	40	ALITTI	69	HBC —	$\Lambda$ , $\Sigma \bar{K}$
1814 ± 4	30	BADIER	65	HBC 0	$\Lambda \bar{K}^0$
1817 ± 7	29	SMITH	65C	HBC —0	$\Lambda \bar{K}^0$ , $\Lambda K^-$
1770		HALSTEINSLID63	FBC	—0	$K^-$ freon 3.5 GeV/c


 **$\Xi(1820)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>24 ± 15 OUR ESTIMATE</b>					
<b>24 ± 6 OUR AVERAGE</b>					
Error includes scale factor of 1.5. See the ideogram below.					
24.6 ± 5.3	280	<sup>1</sup> BIAGI	87	SPEC 0	$\Xi^-$ Be → $(\Lambda K^-)$ X
12 ± 14 ± 1.7	54	BIAGI	87C	SPEC 0	$\Xi^-$ Be → $(\Lambda \bar{K}^0)$ X
72 ± 20	300	BIAGI	81	SPEC —	SPS hyperon beam
21 ± 7	130	GAY	76C	HBC —	$K^- p$ 4.2 GeV/c

• • • We do not use the following data for averages, fits, limits, etc. • • •

23	$\pm 13$	ADAMOVICH	99B	WA89	$\Sigma^-$ nucleus, 345 GeV
99	$\pm 57$	74	BRIEFEL	77	HBC 0 $K^- p$ 2.87 GeV/c
52	$\pm 34$	68	BRIEFEL	77	HBC $-0$ $\Xi(1530)\pi$
72	$\pm 17$	39	BRIEFEL	77	HBC $-$ $\Sigma^-\bar{K}^0$
44	$\pm 11$	44	BRIEFEL	77	HBC 0 $\Lambda\bar{K}^0$
26	$\pm 11$	57	BRIEFEL	77	HBC $-$ $\Lambda K^-$
85	$\pm 58$	DIBIANCA	75	DBC	$-0$ $\Xi\pi\pi, \Xi^*\pi$
51	$\pm 13$	2	BADIER	72	HBC $-0$ Lower mass
58	$\pm 13$	2	BADIER	72	HBC $-0$ Higher mass
103	$^{+38}_{-24}$	3	CRENNELL	70B	DBC $-0$ 3.6, 3.9 GeV/c
48	$^{+36}_{-19}$	4	CRENNELL	70B	DBC $-0$ 3.6, 3.9 GeV/c
55	$^{+40}_{-20}$	ALITTI	69	HBC	$-$ $\Lambda, \Sigma\bar{K}$
12	$\pm 4$	BADIER	65	HBC	0 $\Lambda\bar{K}^0$
30	$\pm 7$	SMITH	65B	HBC	$-0$ $\Lambda\bar{K}$
<80		HALSTEINSLID63	FBC	$-0$	$K^-$ freon 3.5 GeV/c

WEIGHTED AVERAGE  
24 $\pm$ 6 (Error scaled by 1.5)



## $\Xi(1820)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \Lambda\bar{K}$	large
$\Gamma_2 \Sigma\bar{K}$	small
$\Gamma_3 \Xi\pi$	small
$\Gamma_4 \Xi(1530)\pi$	small
$\Gamma_5 \Xi\pi\pi(\text{not } \Xi(1530)\pi)$	

## $\Xi(1820)$ BRANCHING RATIOS

The dominant modes seem to be  $\Lambda\bar{K}$  and (perhaps)  $\Xi(1530)\pi$ , but the branching fractions are very poorly determined.

### $\Gamma(\Lambda\bar{K})/\Gamma_{\text{total}}$

VALUE  
**0.25±0.05 OUR AVERAGE**

$0.24 \pm 0.05$   
 $0.30 \pm 0.15$

DOCUMENT ID	TECN	CHG	COMMENT
ANISOVICH	12A	DPWA	Multichannel
ALITTI	69	HBC	$K^- p$ 3.9–5 GeV/c

$\Gamma_1/\Gamma$

### $\Gamma(\Xi\pi)/\Gamma_{\text{total}}$

VALUE  
**0.10±0.10**

DOCUMENT ID	TECN	CHG	COMMENT
ALITTI	69	HBC	$K^- p$ 3.9–5 GeV/c

$\Gamma_3/\Gamma$

### $\Gamma(\Xi\pi)/\Gamma(\Lambda\bar{K})$

VALUE  
**<0.36** 95  
**0.20±0.20**

DOCUMENT ID	TECN	CHG	COMMENT
GAY	76C	HBC	$K^- p$ 4.2 GeV/c
BADIER	65	HBC	$K^- p$ 3 GeV/c

$\Gamma_3/\Gamma_1$

### $\Gamma(\Xi\pi)/\Gamma(\Xi(1530)\pi)$

VALUE  
**1.5<sup>+0.6</sup><sub>-0.4</sub>**

DOCUMENT ID	TECN	CHG	COMMENT
APSELL	70	HBC	$K^- p$ 2.87 GeV/c

$\Gamma_3/\Gamma_4$

### $\Gamma(\Sigma\bar{K})/\Gamma_{\text{total}}$

VALUE  
**0.30±0.15**

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.02$  TRIPP 67 RVUE Use SMITH 65C

$\Gamma_2/\Gamma$

### $\Gamma(\Sigma\bar{K})/\Gamma(\Lambda\bar{K})$

VALUE  
**0.24±0.10**

DOCUMENT ID	TECN	CHG	COMMENT
GAY	76C	HBC	$K^- p$ 4.2 GeV/c

$\Gamma_2/\Gamma_1$

$\Gamma(\Xi(1530)\pi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.30±0.15</b>	ALITTI	69	HBC	—	$K^- p$ 3.9–5 GeV/c
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
seen	ASTON	85B	LASS		$K^- p$ 11 GeV/c
not seen	<sup>5</sup> HASSALL	81	HBC		$K^- p$ 6.5 GeV/c
<0.25	<sup>6</sup> DAUBER	69	HBC		$K^- p$ 2.7 GeV/c

$\Gamma(\Xi(1530)\pi)/\Gamma(\Lambda\bar{K})$					$\Gamma_4/\Gamma_1$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.38±0.27 OUR AVERAGE</b>	Error includes scale factor of 2.3.				
1.0 ± 0.3	GAY	76C	HBC	—	$K^- p$ 4.2 GeV/c
0.26±0.13	SMITH	65C	HBC	—0	$K^- p$ 2.45–2.7 GeV/c

$\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Lambda\bar{K})$					$\Gamma_5/\Gamma_1$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.30±0.20</b>	BIAGI	87	SPEC	—	$\Xi^- Be$ 116 GeV
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
<0.14	<sup>7</sup> BADIER	65	HBC	0	1 st. dev. limit
>0.1	SMITH	65C	HBC	—0	$K^- p$ 2.45–2.7 GeV/c

$\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Xi(1530)\pi)$					$\Gamma_5/\Gamma_4$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
consistent with zero	GAY	76C	HBC	—	$K^- p$ 4.2 GeV/c
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
0.3±0.5	<sup>8</sup> APSELL	70	HBC	0	$K^- p$ 2.87 GeV/c

### $\Xi(1820)$ FOOTNOTES

<sup>1</sup> BIAGI 87 also sees weak signals in the  $\Xi^-\pi^+\pi^-$  channel at  $1782.6 \pm 1.4$  MeV ( $\Gamma = 6.0 \pm 1.5$  MeV) and  $1831.9 \pm 2.8$  MeV ( $\Gamma = 9.6 \pm 9.9$  MeV).

<sup>2</sup> BADIER 72 adds all channels and divides the peak into lower and higher mass regions. The data can also be fitted with a single Breit-Wigner of mass 1800 MeV and width 150 MeV.

<sup>3</sup> From a fit to inclusive  $\Xi\pi$ ,  $\Xi\pi\pi$ , and  $\Lambda K^-$  spectra.

<sup>4</sup> From a fit to inclusive  $\Xi\pi$  and  $\Xi\pi\pi$  spectra only.

<sup>5</sup> Including  $\Xi\pi\pi$ .

<sup>6</sup> DAUBER 69 uses in part the same data as SMITH 65C.

<sup>7</sup> For the decay mode  $\Xi^-\pi^+\pi^0$  only. This limit includes  $\Xi(1530)\pi$ .

<sup>8</sup> Or less. Upper limit for the 3-body decay.

## $\Xi(1820)$ REFERENCES

ANISOVICH	12A	EPJ A48 15	A.V. Anisovich <i>et al.</i>	(BONN, PNPI)
ADAMOVICH	99B	EPJ C11 271	M.I. Adamovich <i>et al.</i>	(CERN WA89 Collab.)
BIAGI	87	ZPHY C34 15	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+)
BIAGI	87C	ZPHY C34 175	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+) JP
ASTON	85B	PR D32 2270	D. Aston <i>et al.</i>	(SLAC, CARL, CNRC, CINC)
JENKINS	83	PRL 51 951	C.M. Jenkins <i>et al.</i>	(FSU, BRAN, LBL+)
BIAGI	81	ZPHY C9 305	S.F. Biagi <i>et al.</i>	(BRIS, CAVE, GEVA+)
HASSALL	81	NP B189 397	J.K. Hassall <i>et al.</i>	(CAVE, MSU)
TEODORO	78	PL 77B 451	D. Teodoro <i>et al.</i>	(AMST, CERN, NIJM+) JP
BRIEFEL	77	PR D16 2706	E. Briefel <i>et al.</i>	(BRAN, UMD, SYRA+)
Also		PRL 23 884	S.P. Apsell <i>et al.</i>	(BRAN, UMD, SYRA+)
GAY	76C	PL 62B 477	J.B. Gay <i>et al.</i>	(AMST, CERN, NIJM) IJ
DIBIANCA	75	NP B98 137	F.A. Dibianca, R.J. Endorf	(CMU)
BADIER	72	NP B37 429	J. Badier <i>et al.</i>	(EPOL)
APSELL	70	PRL 24 777	S.P. Apsell <i>et al.</i>	(BRAN, UMD, SYRA+) I
CRENNELL	70B	PR D1 847	D.J. Crennell <i>et al.</i>	(BNL)
ALITTI	69	PRL 22 79	J. Alitti <i>et al.</i>	(BNL, SYRA) I
DAUBER	69	PR 179 1262	P.M. Dauber <i>et al.</i>	(LRL)
TRIPP	67	NP B3 10	R.D. Tripp <i>et al.</i>	(LRL, SLAC, CERN+)
BADIER	65	PL 16 171	J. Badier <i>et al.</i>	(EPOL, SACL, AMST) I
SMITH	65B	Athens Conf. 251	G.A. Smith, J.S. Lindsey	(LRL)
SMITH	65C	PRL 14 25	G.A. Smith <i>et al.</i>	(LRL) IJP
HALSTEINSLID	63	Siena Conf. 1 73	A. Halsteinslid <i>et al.</i>	(BERG, CERN, EPOL+) I

## OTHER RELATED PAPERS

TEODORO	78	PL 77B 451	D. Teodoro <i>et al.</i>	(AMST, CERN, NIJM+) JP
BRIEFEL	75	PR D12 1859	E. Briefel <i>et al.</i>	(BRAN, UMD, SYRA+)
SCHMIDT	73	Purdue Conf. 363	P.E. Schmidt	(BRAN)
MERRILL	68	PR 167 1202	D.W. Merrill, J. Button-Shafer	(LRL)
SMITH	64	PRL 13 61	G.A. Smith <i>et al.</i>	(LRL) IJP