

**$\Xi_c(2815)$**  $I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$  Status: \*\*\*

Seen in both  $\Xi_c\pi$  and  $\Xi_c\pi\pi$  decays. The simplest assignment is that this belongs to the same SU(4) multiplet as the  $\Lambda(1520)$  and the  $\Lambda_c(2625)$ , but the spin and parity have not been measured.

 **$\Xi_c(2815)$  MASSES**

The masses are obtained from the mass-difference measurements that follow.

 **$\Xi_c(2815)^+$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2816.67 \pm 0.31</math> OUR FIT</b>		Error includes scale factor of 1.1.		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2817.0 $\pm 1.2$ $^{+0.7}_{-0.8}$	73 $\pm 10$	LESIAK	08 BELL	$e^+e^- \approx \gamma(4S)$

 **$\Xi_c(2815)^0$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2820.22 \pm 0.32</math> OUR FIT</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2820.4 $\pm 1.4$ $^{+0.9}_{-1.0}$	48 $\pm 8$	LESIAK	08 BELL	$e^+e^- \approx \gamma(4S)$

 **$\Xi_c(2815) - \Xi_c$  MASS DIFFERENCES** **$m_{\Xi_c(2815)^+} - m_{\Xi_c^+}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>348.80 \pm 0.10</math> OUR FIT</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
348.6 $\pm 0.6$ $\pm 1.0$	20	ALEXANDER	99B CLE2	$e^+e^- \approx \gamma(4S)$

 **$m_{\Xi_c(2815)^0} - m_{\Xi_c^0}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>349.35 \pm 0.11</math> OUR FIT</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
347.2 $\pm 0.7$ $\pm 2.0$	9	ALEXANDER	99B CLE2	$e^+e^- \approx \gamma(4S)$

 **$\Xi_c(2815)^+ - \Xi_c(2815)^0$  MASS DIFFERENCE** **$m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-3.55 \pm 0.28</math> OUR FIT</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-3.47 $\pm 0.12 \pm 0.48$		YELTON	16 BELL	941 and 1258 evts
-3.4 $\pm 1.9 \pm 0.9$		LESIAK	08 BELL	73 & 48 events

## $\Xi_c(2815)$ WIDTHS

### $\Xi_c(2815)^+$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.43±0.20±0.17</b>	941		YELTON 16	BELL	$e^+ e^-$ , $\gamma$ regions

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.5 90 ALEXANDER 99B CLE2  $e^+ e^- \approx \gamma(4S)$

### $\Xi_c(2815)^0$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.54±0.18±0.17</b>	1258		YELTON 16	BELL	$e^+ e^-$ , $\gamma$ regions

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.5 90 ALEXANDER 99B CLE2  $e^+ e^- \approx \gamma(4S)$

## $\Xi_c(2815)$ DECAY MODES

The  $\Xi_c \pi\pi$  modes are consistent with being entirely via  $\Xi_c(2645)\pi$ .

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \Xi_c \pi$	seen
$\Gamma_2 \quad \Xi_c^+ \pi^+ \pi^-$	seen
$\Gamma_3 \quad \Xi_c^0 \pi^+ \pi^-$	seen

## $\Xi_c(2815)$ REFERENCES

YELTON	16	PR D94 052011	J. Yelton <i>et al.</i>	(BELLE Collab.)
LESIAK	08	PL B665 9	T. Lesiak <i>et al.</i>	(BELLE Collab.)
ALEXANDER	99B	PRL 83 3390	J.P. Alexander <i>et al.</i>	(CLEO Collab.)