

$D^\pm$ 

$$I(J^P) = \frac{1}{2}(0^-)$$

## $D^\pm$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^{*}(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1869.65 ± 0.05 OUR FIT</b>				
<b>1869.5 ± 0.4 OUR AVERAGE</b>				
1869.53 ± 0.49 ± 0.20	110 ± 15	ANASHIN	10A	KEDR $e^+ e^-$ at $\psi(3770)$
1870.0 ± 0.5 ± 1.0	317	BARLAG	90C	ACCM $\pi^-$ Cu 230 GeV
1869.4 ± 0.6		<sup>1</sup> TRILLING	81	RVUE $e^+ e^-$ 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1875 ± 10	9	ADAMOVICH	87	EMUL Photoproduction
1860 ± 16	6	ADAMOVICH	84	EMUL Photoproduction
1863 ± 4		DERRICK	84	HRS $e^+ e^-$ 29 GeV
1868.4 ± 0.5		<sup>1</sup> SCHINDLER	81	MRK2 $e^+ e^-$ 3.77 GeV
1874 ± 5		GOLDHABER	77	MRK1 $D^0$ , $D^+$ recoil spectra
1868.3 ± 0.9		<sup>1</sup> PERUZZI	77	LGW $e^+ e^-$ 3.77 GeV
1874 ± 11		PICCOLO	77	MRK1 $e^+ e^-$ 4.03, 4.41 GeV
1876 ± 15	50	PERUZZI	76	MRK1 $K^\mp \pi^\pm \pi^\pm$

<sup>1</sup> PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision  $J/\psi(1S)$  and  $\psi(2S)$  measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted.

## $D^\pm$ MEAN LIFE

Measurements with an error  $> 100 \times 10^{-15}$  s have been omitted from the Listings.

VALUE ( $10^{-15}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1040 ± 7 OUR AVERAGE</b>				
1039.4 ± 4.3 ± 7.0	110k	LINK	02F	FOCS $\gamma$ nucleus, $\approx 180$ GeV
1033.6 ± 22.1 ± 9.9	3.7k	BONVICINI	99	CLEO $e^+ e^- \approx \gamma(4S)$
1048 ± 15 ± 11	9k	FRABETTI	94D	E687 $D^+ \rightarrow K^- \pi^+ \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1075 ± 40 ± 18	2.4k	FRABETTI	91	E687 $\gamma$ Be, $D^+ \rightarrow K^- \pi^+ \pi^+$
1030 ± 80 ± 60	200	ALVAREZ	90	NA14 $\gamma$ , $D^+ \rightarrow K^- \pi^+ \pi^+$
1050 ± 77 ± 72	317	<sup>1</sup> BARLAG	90C	ACCM $\pi^-$ Cu 230 GeV
1050 ± 80 ± 70	363	ALBRECHT	88I	ARG $e^+ e^-$ 10 GeV
1090 ± 30 ± 25	2.9k	RAAB	88	E691 Photoproduction

<sup>1</sup> BARLAG 90C estimates the systematic error to be negligible.

## **$D^+$ DECAY MODES**

Most decay modes (other than the semileptonic modes) that involve a neutral  $K$  meson are now given as  $K_S^0$  modes, not as  $\bar{K}^0$  modes. Nearly always it is a  $K_S^0$  that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that  $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$ .

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
<b>Inclusive modes</b>		
$\Gamma_1 e^+ \text{ semileptonic}$	$(16.07 \pm 0.30) \%$	
$\Gamma_2 \mu^+ \text{ anything}$	$(17.6 \pm 3.2) \%$	
$\Gamma_3 K^- \text{ anything}$	$(25.7 \pm 1.4) \%$	
$\Gamma_4 \bar{K}^0 \text{ anything} + K^0 \text{ anything}$	$(61 \pm 5) \%$	
$\Gamma_5 K^+ \text{ anything}$	$(5.9 \pm 0.8) \%$	
$\Gamma_6 K^*(892)^- \text{ anything}$	$(6 \pm 5) \%$	
$\Gamma_7 \bar{K}^*(892)^0 \text{ anything}$	$(23 \pm 5) \%$	
$\Gamma_8 K^*(892)^0 \text{ anything}$	$< 6.6 \%$	CL=90%
$\Gamma_9 \eta \text{ anything}$	$(6.3 \pm 0.7) \%$	
$\Gamma_{10} \eta' \text{ anything}$	$(1.04 \pm 0.18) \%$	
$\Gamma_{11} \phi \text{ anything}$	$(1.03 \pm 0.12) \%$	
<b>Leptonic and semileptonic modes</b>		
$\Gamma_{12} e^+ \nu_e$	$< 8.8 \times 10^{-6}$	CL=90%
$\Gamma_{13} \gamma e^+ \nu_e$	$< 3.0 \times 10^{-5}$	CL=90%
$\Gamma_{14} \mu^+ \nu_\mu$	$(3.74 \pm 0.17) \times 10^{-4}$	
$\Gamma_{15} \tau^+ \nu_\tau$	$< 1.2 \times 10^{-3}$	CL=90%
$\Gamma_{16} \bar{K}^0 e^+ \nu_e$	$(8.73 \pm 0.10) \%$	
$\Gamma_{17} \bar{K}^0 \mu^+ \nu_\mu$	$(8.74 \pm 0.19) \%$	
$\Gamma_{18} K^- \pi^+ e^+ \nu_e$	$(3.89 \pm 0.13) \%$	S=2.1
$\Gamma_{19} \bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(3.66 \pm 0.12) \%$	
$\Gamma_{20} (K^- \pi^+) [0.8-1.0] \text{GeV} e^+ \nu_e$	$(3.39 \pm 0.09) \%$	
$\Gamma_{21} (K^- \pi^+)_{S-wave} e^+ \nu_e$	$(2.28 \pm 0.11) \times 10^{-3}$	
$\Gamma_{22} \bar{K}^*(1410)^0 e^+ \nu_e, \bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	$< 6 \times 10^{-3}$	CL=90%
$\Gamma_{23} \bar{K}_2^*(1430)^0 e^+ \nu_e, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{24} K^- \pi^+ e^+ \nu_e \text{ nonresonant}$	$< 7 \times 10^{-3}$	CL=90%
$\Gamma_{25} K^- \pi^+ \mu^+ \nu_\mu$	$(3.65 \pm 0.34) \%$	
$\Gamma_{26} \bar{K}^*(892)^0 \mu^+ \nu_\mu, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(3.52 \pm 0.10) \%$	
$\Gamma_{27} K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant}$	$(1.9 \pm 0.5) \times 10^{-3}$	

$\Gamma_{28}$	$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	$< 1.5 \times 10^{-3}$	CL=90%
$\Gamma_{29}$	$\pi^0 e^+ \nu_e$	$(3.72 \pm 0.17) \times 10^{-3}$	S=2.0
$\Gamma_{30}$	$\eta e^+ \nu_e$	$(1.14 \pm 0.10) \times 10^{-3}$	
$\Gamma_{31}$	$\rho^0 e^+ \nu_e$	$(2.18^{+0.17}_{-0.25}) \times 10^{-3}$	
$\Gamma_{32}$	$\rho^0 \mu^+ \nu_\mu$	$(2.4 \pm 0.4) \times 10^{-3}$	
$\Gamma_{33}$	$\omega e^+ \nu_e$	$(1.69 \pm 0.11) \times 10^{-3}$	
$\Gamma_{34}$	$\eta'(958) e^+ \nu_e$	$(2.2 \pm 0.5) \times 10^{-4}$	
$\Gamma_{35}$	$\phi e^+ \nu_e$	$< 1.3 \times 10^{-5}$	CL=90%
$\Gamma_{36}$	$D^0 e^+ \nu_e$	$< 1.0 \times 10^{-4}$	CL=90%

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\Gamma_{37}$	$\bar{K}^*(892)^0 e^+ \nu_e$	$(5.40 \pm 0.10) \%$	S=1.1
$\Gamma_{38}$	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	$(5.25 \pm 0.15) \%$	
$\Gamma_{39}$	$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	$< 2.3 \times 10^{-4}$	CL=90%
$\Gamma_{40}$	$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	$< 1.5 \times 10^{-3}$	CL=90%

### Hadronic modes with a $\bar{K}$ or $\bar{K}\bar{K}\bar{K}$

$\Gamma_{41}$	$K_S^0 \pi^+$	$(1.47 \pm 0.08) \%$	S=3.0
$\Gamma_{42}$	$K_L^0 \pi^+$	$(1.46 \pm 0.05) \%$	
$\Gamma_{43}$	$K^- 2\pi^+$	[a] $(8.98 \pm 0.28) \%$	S=2.2
$\Gamma_{44}$	$(K^- \pi^+)_{S\text{-wave}} \pi^+$	$(7.20 \pm 0.25) \%$	
$\Gamma_{45}$	$\bar{K}_0^*(700)^0 \pi^+, \bar{K}_0^*(700) \rightarrow K^- \pi^+$		
$\Gamma_{46}$	$\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[b] $(1.19 \pm 0.07) \%$	
$\Gamma_{47}$	$\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(10.0 \pm 1.1) \times 10^{-3}$	
$\Gamma_{48}$	$\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+$	not seen	
$\Gamma_{49}$	$\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[b] $(2.2 \pm 0.7) \times 10^{-4}$	
$\Gamma_{50}$	$\bar{K}^*(1680)^0 \pi^+, \bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[b] $(2.1 \pm 1.0) \times 10^{-4}$	
$\Gamma_{51}$	$K^- (2\pi^+)_{I=2}$	$(1.39 \pm 0.26) \%$	
$\Gamma_{52}$	$K^- 2\pi^+ \text{ nonresonant}$		
$\Gamma_{53}$	$K_S^0 \pi^+ \pi^0$	[a] $(7.05 \pm 0.27) \%$	
$\Gamma_{54}$	$K_S^0 \rho^+$	$(5.9^{+0.6}_{-0.4}) \%$	
$\Gamma_{55}$	$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.5^{+1.1}_{-1.4}) \times 10^{-3}$	
$\Gamma_{56}$	$\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	$(2.52 \pm 0.31) \times 10^{-3}$	

$\Gamma_{57}$	$\bar{K}_0^*(1430)^0\pi^+, \bar{K}_0^{*0} \rightarrow K_S^0\pi^0$	$(2.6 \pm 0.9) \times 10^{-3}$
$\Gamma_{58}$	$\bar{K}_0^*(1680)^0\pi^+, \bar{K}_0^{*0} \rightarrow K_S^0\pi^0$	$(9 \pm 7) \times 10^{-4}$
$\Gamma_{59}$	$\bar{\kappa}^0\pi^+, \bar{\kappa}^0 \rightarrow K_S^0\pi^0$	$(5.4 \pm 5.0) \times 10^{-3}$
$\Gamma_{60}$	$K_S^0\pi^+\pi^0$ nonresonant	$(3 \pm 4) \times 10^{-3}$
$\Gamma_{61}$	$K_S^0\pi^+\pi^0$ nonresonant and $\bar{\kappa}^0\pi^+$	$(1.31 \pm 0.21) \%$
$\Gamma_{62}$	$(K_S^0\pi^0)_{S\text{-wave}}\pi^+$	$(1.22 \pm 0.26) \%$
$\Gamma_{63}$	$K^-2\pi^+\pi^0$	[c] $(5.98 \pm 0.23) \%$
$\Gamma_{64}$	$K_S^02\pi^+\pi^-$	[c] $(2.97 \pm 0.11) \%$
$\Gamma_{65}$	$K^-3\pi^+\pi^-$	[a] $(5.5 \pm 0.5) \times 10^{-3}$
$\Gamma_{66}$	$\bar{K}^*(892)^02\pi^+\pi^-, \bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(1.2 \pm 0.4) \times 10^{-3}$
$\Gamma_{67}$	$\bar{K}^*(892)^0\rho^0\pi^+, \bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(2.2 \pm 0.4) \times 10^{-3}$
$\Gamma_{68}$	$\bar{K}^*(892)^0a_1(1260)^+$	[d] $(8.9 \pm 1.8) \times 10^{-3}$
$\Gamma_{69}$	$\bar{K}^*(892)^02\pi^+\pi^-$ no- $\rho$ , $\bar{K}^*(892)^0 \rightarrow K^-\pi^+$	
$\Gamma_{70}$	$K^-\rho^02\pi^+$	$(1.65 \pm 0.27) \times 10^{-3}$
$\Gamma_{71}$	$K^-3\pi^+\pi^-$ nonresonant	$(3.9 \pm 2.8) \times 10^{-4}$
$\Gamma_{72}$	$K^+2K_S^0$	$(2.54 \pm 0.13) \times 10^{-3}$
$\Gamma_{73}$	$K^+K^-K_S^0\pi^+$	$(2.3 \pm 0.5) \times 10^{-4}$

**Pionic modes**

$\Gamma_{74}$	$\pi^+\pi^0$	$(1.17 \pm 0.06) \times 10^{-3}$
$\Gamma_{75}$	$2\pi^+\pi^-$	$(3.13 \pm 0.19) \times 10^{-3}$
$\Gamma_{76}$	$\rho^0\pi^+$	$(8.0 \pm 1.4) \times 10^{-4}$
$\Gamma_{77}$	$\pi^+(\pi^+\pi^-)_{S\text{-wave}}$	$(1.75 \pm 0.16) \times 10^{-3}$
$\Gamma_{78}$	$\sigma\pi^+, \sigma \rightarrow \pi^+\pi^-$	$(1.32 \pm 0.12) \times 10^{-3}$
$\Gamma_{79}$	$f_0(980)\pi^+, f_0(980) \rightarrow \pi^+\pi^-$	$(1.50 \pm 0.32) \times 10^{-4}$
$\Gamma_{80}$	$f_0(1370)\pi^+, f_0(1370) \rightarrow \pi^+\pi^-$	$(8 \pm 4) \times 10^{-5}$
$\Gamma_{81}$	$f_2(1270)\pi^+, f_2(1270) \rightarrow \pi^+\pi^-$	$(4.8 \pm 0.8) \times 10^{-4}$
$\Gamma_{82}$	$\rho(1450)^0\pi^+, \rho(1450)^0 \rightarrow \pi^+\pi^-$	$< 8 \times 10^{-5}$
$\Gamma_{83}$	$f_0(1500)\pi^+, f_0(1500) \rightarrow \pi^+\pi^-$	$(1.1 \pm 0.4) \times 10^{-4}$
$\Gamma_{84}$	$f_0(1710)\pi^+, f_0(1710) \rightarrow \pi^+\pi^-$	$< 5 \times 10^{-5}$

CL=95%

$\Gamma_{85}$	$f_0(1790)\pi^+, f_0(1790) \rightarrow \pi^+\pi^-$	< 6	$\times 10^{-5}$	CL=95%
$\Gamma_{86}$	$(\pi^+\pi^+)_{S\text{-wave}}\pi^-$	< 1.2	$\times 10^{-4}$	CL=95%
$\Gamma_{87}$	$2\pi^+\pi^-$ nonresonant	< 1.1	$\times 10^{-4}$	CL=95%
$\Gamma_{88}$	$\pi^+2\pi^0$	$(4.5 \pm 0.4) \times 10^{-3}$		
$\Gamma_{89}$	$2\pi^+\pi^-\pi^0$	$(1.11 \pm 0.08)\%$		
$\Gamma_{90}$	$3\pi^+2\pi^-$	$(1.59 \pm 0.16) \times 10^{-3}$		S=1.1
$\Gamma_{91}$	$\eta\pi^+$	$(3.33 \pm 0.21) \times 10^{-3}$		S=1.4
$\Gamma_{92}$	$\eta\pi^+\pi^0$	$(1.38 \pm 0.35) \times 10^{-3}$		
$\Gamma_{93}$	$\omega\pi^+$	$(2.8 \pm 0.6) \times 10^{-4}$		
$\Gamma_{94}$	$\eta'(958)\pi^+$	$(4.60 \pm 0.31) \times 10^{-3}$		
$\Gamma_{95}$	$\eta'(958)\pi^+\pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$		

### Hadronic modes with a $K\bar{K}$ pair

$\Gamma_{96}$	$K^+K_S^0$	$(2.83 \pm 0.16) \times 10^{-3}$		S=2.8
$\Gamma_{97}$	$K^+K^-\pi^+$	[a]	$(9.51 \pm 0.34) \times 10^{-3}$	S=1.6
$\Gamma_{98}$	$\phi\pi^+, \phi \rightarrow K^+K^-$	$(2.64 \pm 0.11) \times 10^{-3}$		
$\Gamma_{99}$	$K^+\bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(2.44^{+0.11}_{-0.15}) \times 10^{-3}$		
$\Gamma_{100}$	$K^+\bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow$ $K^-\pi^+$	$(1.79 \pm 0.34) \times 10^{-3}$		
$\Gamma_{101}$	$K^+\bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow$ $K^-\pi^+$	$(1.6^{+1.2}_{-0.8}) \times 10^{-4}$		
$\Gamma_{102}$	$K^+\bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^-\pi^+$	$(6.7^{+3.4}_{-2.1}) \times 10^{-4}$		
$\Gamma_{103}$	$a_0(1450)^0\pi^+, a_0^0 \rightarrow K^+K^-$	$(4.4^{+7.0}_{-1.8}) \times 10^{-4}$		
$\Gamma_{104}$	$\phi(1680)\pi^+, \phi \rightarrow K^+K^-$	$(4.9^{+4.0}_{-1.9}) \times 10^{-5}$		
$\Gamma_{105}$	$K_S^0K_S^0\pi^+$	$(2.70 \pm 0.13) \times 10^{-3}$		
$\Gamma_{106}$	$K^+K_S^0\pi^+\pi^-$	$(1.67 \pm 0.18) \times 10^{-3}$		
$\Gamma_{107}$	$K_S^0K^-2\pi^+$	$(2.28 \pm 0.18) \times 10^{-3}$		
$\Gamma_{108}$	$K^+K^-2\pi^+\pi^-$	$(2.2 \pm 1.2) \times 10^{-4}$		

A few poorly measured branching fractions:

$\Gamma_{109}$	$\phi\pi^+\pi^0$	$(2.3 \pm 1.0)\%$		
$\Gamma_{110}$	$\phi\rho^+$	< 1.4	%	CL=90%
$\Gamma_{111}$	$K^+K^-\pi^+\pi^0$ non- $\phi$	$(1.5^{+0.7}_{-0.6})\%$		
$\Gamma_{112}$	$K^*(892)^+K_S^0$	$(1.6 \pm 0.7)\%$		

**Doubly Cabibbo-suppressed modes**

$\Gamma_{113}$	$K^+ \pi^0$	$(1.81 \pm 0.27) \times 10^{-4}$	S=1.4
$\Gamma_{114}$	$K^+ \eta$	$(1.02 \pm 0.16) \times 10^{-4}$	
$\Gamma_{115}$	$K^+ \eta'(958)$	$(1.73 \pm 0.22) \times 10^{-4}$	
$\Gamma_{116}$	$K^+ \pi^+ \pi^-$	$(5.19 \pm 0.26) \times 10^{-4}$	
$\Gamma_{117}$	$K^+ \rho^0$	$(2.0 \pm 0.5) \times 10^{-4}$	
$\Gamma_{118}$	$K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow K^+ \pi^-$	$(2.4 \pm 0.4) \times 10^{-4}$	
$\Gamma_{119}$	$K^+ f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$	$(4.6 \pm 2.8) \times 10^{-5}$	
$\Gamma_{120}$	$K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow K^+ \pi^-$	$(4.2 \pm 2.8) \times 10^{-5}$	
$\Gamma_{121}$	$K^+ \pi^+ \pi^-$ nonresonant	not seen	
$\Gamma_{122}$	$2K^+ K^-$	$(8.5 \pm 2.0) \times 10^{-5}$	

 **$\Delta C = 1$  weak neutral current ( $C1$ ) modes, or****Lepton Family number ( $LF$ ) or Lepton number ( $L$ ) violating modes**

$\Gamma_{123}$	$\pi^+ e^+ e^-$	$C1$	$< 1.1 \times 10^{-6}$	CL=90%
$\Gamma_{124}$	$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[e]	$(1.7 \pm 1.4) \times 10^{-6}$	
$\Gamma_{125}$	$\pi^+ \mu^+ \mu^-$	$C1$	$< 7.3 \times 10^{-8}$	CL=90%
$\Gamma_{126}$	$\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$	[e]	$(1.8 \pm 0.8) \times 10^{-6}$	
$\Gamma_{127}$	$\rho^+ \mu^+ \mu^-$	$C1$	$< 5.6 \times 10^{-4}$	CL=90%
$\Gamma_{128}$	$K^+ e^+ e^-$	[f]	$< 1.0 \times 10^{-6}$	CL=90%
$\Gamma_{129}$	$K^+ \mu^+ \mu^-$	[f]	$< 4.3 \times 10^{-6}$	CL=90%
$\Gamma_{130}$	$\pi^+ e^+ \mu^-$	$LF$	$< 2.9 \times 10^{-6}$	CL=90%
$\Gamma_{131}$	$\pi^+ e^- \mu^+$	$LF$	$< 3.6 \times 10^{-6}$	CL=90%
$\Gamma_{132}$	$K^+ e^+ \mu^-$	$LF$	$< 1.2 \times 10^{-6}$	CL=90%
$\Gamma_{133}$	$K^+ e^- \mu^+$	$LF$	$< 2.8 \times 10^{-6}$	CL=90%
$\Gamma_{134}$	$\pi^- 2e^+$	$L$	$< 1.1 \times 10^{-6}$	CL=90%
$\Gamma_{135}$	$\pi^- 2\mu^+$	$L$	$< 2.2 \times 10^{-8}$	CL=90%
$\Gamma_{136}$	$\pi^- e^+ \mu^+$	$L$	$< 2.0 \times 10^{-6}$	CL=90%
$\Gamma_{137}$	$\rho^- 2\mu^+$	$L$	$< 5.6 \times 10^{-4}$	CL=90%
$\Gamma_{138}$	$K^- 2e^+$	$L$	$< 9 \times 10^{-7}$	CL=90%
$\Gamma_{139}$	$K^- 2\mu^+$	$L$	$< 1.0 \times 10^{-5}$	CL=90%
$\Gamma_{140}$	$K^- e^+ \mu^+$	$L$	$< 1.9 \times 10^{-6}$	CL=90%
$\Gamma_{141}$	$K^*(892)^- 2\mu^+$	$L$	$< 8.5 \times 10^{-4}$	CL=90%
$\Gamma_{142}$	Unaccounted decay modes		$(63.7 \pm 0.6) \%$	S=1.6

[a] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.

[b] These subfractions of the  $K^- 2\pi^+$  mode are uncertain: see the Particle Listings.

[c] Submodes of the  $D^+ \rightarrow K^- 2\pi^+ \pi^0$  and  $K_S^0 2\pi^+ \pi^-$  modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.

[d] The unseen decay modes of the resonances are included.

[e] This is *not* a test for the  $\Delta C=1$  weak neutral current, but leads to the  $\pi^+ \ell^+ \ell^-$  final state.

[f] This mode is not a useful test for a  $\Delta C=1$  weak neutral current because both quarks must change flavor in this decay.

## CONSTRAINED FIT INFORMATION

An overall fit to 22 branching ratios uses 33 measurements and one constraint to determine 14 parameters. The overall fit has a  $\chi^2 = 45.6$  for 20 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_{18}$	0												
$x_{31}$	0	0											
$x_{37}$	0	0	0										
$x_{38}$	8	0	0	0									
$x_{41}$	0	42	0	0	0								
$x_{43}$	0	72	0	0	0	59							
$x_{65}$	0	25	0	0	0	20	34						
$x_{90}$	0	23	0	0	0	19	32	77					
$x_{91}$	0	24	0	0	0	19	33	11	10				
$x_{96}$	0	40	0	0	0	85	56	19	18	18			
$x_{97}$	0	63	0	0	0	52	88	30	28	29			
$x_{113}$	0	13	0	0	0	11	19	6	6	6			
$x_{142}$	-34	-72	-3	-18	-28	-61	-85	-39	-35	-31			
	$x_{17}$	$x_{18}$	$x_{31}$	$x_{37}$	$x_{38}$	$x_{41}$	$x_{43}$	$x_{65}$	$x_{90}$	$x_{91}$			
$x_{97}$	49												
$x_{113}$	10	16											
$x_{142}$	-57	-76	-16										
	$x_{96}$	$x_{97}$	$x_{113}$										

## **$D^+$ BRANCHING RATIOS**

Some now-obsolete measurements have been omitted from these Listings.

### — c-quark decays —

#### **$\Gamma(c \rightarrow e^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$**

For the Summary Table, we only use the average of  $e^+$  and  $\mu^+$  measurements from  $Z^0 \rightarrow c\bar{c}$  decays; see the second data block below.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.103 \pm 0.009^{+0.009}_{-0.008}</math></b>	378	<sup>1</sup> ABBIENDI	99K OPAL	$Z^0 \rightarrow c\bar{c}$

<sup>1</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0 \pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

#### **$\Gamma(c \rightarrow \mu^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$**

For the Summary Table, we only use the average of  $e^+$  and  $\mu^+$  measurements from  $Z^0 \rightarrow c\bar{c}$  decays; see the next data block.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.082 \pm 0.005</math> OUR AVERAGE</b>				
$0.073 \pm 0.008 \pm 0.002$	73	KAYIS-TOPAK.05	CHRS	$\nu_\mu$ emulsion
$0.095 \pm 0.007^{+0.014}_{-0.013}$	2829	ASTIER	00D NOMD	$\nu_\mu$ Fe $\rightarrow \mu^- \mu^+ X$
$0.090 \pm 0.007^{+0.007}_{-0.006}$	476	<sup>1</sup> ABBIENDI	99K OPAL	$Z^0 \rightarrow c\bar{c}$
$0.086 \pm 0.017^{+0.008}_{-0.007}$	69	<sup>2</sup> ALBRECHT	92F ARG	$e^+ e^- \approx 10$ GeV
$0.078 \pm 0.009 \pm 0.012$		ONG	88 MRK2	$e^+ e^-$ 29 GeV
$0.078 \pm 0.015 \pm 0.02$		BARTEL	87 JADE	$e^+ e^-$ 34.6 GeV
$0.082 \pm 0.012^{+0.02}_{-0.01}$		ALTHOFF	84G TASS	$e^+ e^-$ 34.5 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.093 \pm 0.009 \pm 0.009$	88	KAYIS-TOPAK.02	CHRS	See KAYIS-TOPAKSU 05
$0.089 \pm 0.018 \pm 0.025$		BARTEL	85J JADE	See BARTEL 87

<sup>1</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0 \pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

<sup>2</sup> ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed  $D^*(2010)^+ \rightarrow D^0 \pi^+$  decays.

#### **$\Gamma(c \rightarrow \ell^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$**

This is an average (not a sum) of  $e^+$  and  $\mu^+$  measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.096 \pm 0.004</math> OUR AVERAGE</b>				
$0.0958 \pm 0.0042 \pm 0.0028$	1828	<sup>1</sup> ABREU	00O DLPH	$Z^0 \rightarrow c\bar{c}$
$0.095 \pm 0.006^{+0.007}_{-0.006}$	854	<sup>2</sup> ABBIENDI	99K OPAL	$Z^0 \rightarrow c\bar{c}$

<sup>1</sup> ABREU 00O uses leptons opposite fully reconstructed  $D^*(2010)^+$ ,  $D^+$ , or  $D^0$  mesons.

<sup>2</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0 \pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

$\Gamma(c \rightarrow D^*(2010)^+ \text{anything})/\Gamma(c \rightarrow \text{anything})$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.255 <math>\pm</math> 0.015 <math>\pm</math> 0.008</b>	2371	1 ABREU	000 DLPH	$Z^0 \rightarrow c\bar{c}$

<sup>1</sup> ABREU 000 uses slow pions opposite fully reconstructed  $D^*(2010)^+$ ,  $D^+$ , or  $D^0$  mesons as a signal of  $D^*(2010)^-$  production.

**Inclusive modes**

 $\Gamma(e^+ \text{semileptonic})/\Gamma_{\text{total}}$ 

The sum of our  $\overline{K}^0 e^+ \nu_e$ ,  $\overline{K}^*(892) e^+ \nu_e$ ,  $\pi^0 e^+ \nu_e$ ,  $\eta e^+ \nu_e$ ,  $\rho^0 e^+ \nu_e$ , and  $\omega e^+ \nu_e$  branching fractions is  $15.3 \pm 0.4\%$ .

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.07 <math>\pm</math> 0.30 OUR AVERAGE</b>				

$16.13 \pm 0.10 \pm 0.29$	$26.2 \pm 0.2k$	<sup>1</sup> ASNER	10 CLEO	$e^+ e^-$ at 3774 MeV
$15.2 \pm 0.9 \pm 0.8$	$521 \pm 32$	ABLIKIM	07G BES2	$e^+ e^- \approx \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$16.13 \pm 0.20 \pm 0.33$	$8798 \pm 105$	<sup>2</sup> ADAM	06A CLEO	See ASNER 10
$17.0 \pm 1.9 \pm 0.7$	158	BALTRUSAIT..85B MRK3	$e^+ e^-$ 3.77 GeV	

<sup>1</sup> Using the  $D^+$  and  $D^0$  lifetimes, ASNER 10 finds that the ratio of the  $D^+$  and  $D^0$  semileptonic widths is  $0.985 \pm 0.015 \pm 0.024$ .

<sup>2</sup> Using the  $D^+$  and  $D^0$  lifetimes, ADAM 06A finds that the ratio of the  $D^+$  and  $D^0$  inclusive  $e^+$  widths is  $0.985 \pm 0.028 \pm 0.015$ , consistent with the isospin-invariance prediction of 1.

 $\Gamma(\mu^+ \text{anything})/\Gamma_{\text{total}}$ 
 $\Gamma_2/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17.6 <math>\pm</math> 2.7 <math>\pm</math> 1.8</b>	$100 \pm 12$	<sup>1</sup> ABLIKIM	08L BES2	$e^+ e^- \approx \psi(3772)$

<sup>1</sup> ABLIKIM 08L finds the ratio of  $D^+ \rightarrow \mu^+ X$  and  $D^0 \rightarrow \mu^+ X$  branching fractions to be  $2.59 \pm 0.70 \pm 0.25$ , in accord with the ratio of  $D^+$  and  $D^0$  lifetimes,  $2.54 \pm 0.02$ .

 $\Gamma(K^- \text{anything})/\Gamma_{\text{total}}$ 
 $\Gamma_3/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.7 <math>\pm</math> 1.4 OUR AVERAGE</b>				
$24.7 \pm 1.3 \pm 1.2$	$631 \pm 33$	ABLIKIM	07G BES2	$e^+ e^- \approx \psi(3770)$
$27.8^{+3.6}_{-3.1}$		BARLAG	92C ACCM	$\pi^-$ Cu 230 GeV
$27.1 \pm 2.3 \pm 2.4$		COFFMAN	91 MRK3	$e^+ e^-$ 3.77 GeV

 $[\Gamma(\overline{K}^0 \text{anything}) + \Gamma(K^0 \text{anything})]/\Gamma_{\text{total}}$ 
 $\Gamma_4/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>61 <math>\pm</math> 5 OUR AVERAGE</b>				
$60.5 \pm 5.5 \pm 3.3$	$244 \pm 22$	ABLIKIM	06U BES2	$e^+ e^-$ at 3773 MeV
$61.2 \pm 6.5 \pm 4.3$		COFFMAN	91 MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(K^+ \text{anything})/\Gamma_{\text{total}}$ 
 $\Gamma_5/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.9 <math>\pm</math> 0.8 OUR AVERAGE</b>				
$6.1 \pm 0.9 \pm 0.4$	$189 \pm 27$	ABLIKIM	07G BES2	$e^+ e^- \approx \psi(3770)$
$5.5 \pm 1.3 \pm 0.9$		COFFMAN	91 MRK3	$e^+ e^-$ 3.77 GeV



• • • We do not use the following data for averages, fits, limits, etc. • • •

12.2	$\frac{+11.1}{5.3}$	$\pm 1.0$	3	<sup>3</sup> ABLIKIM	05D BES	$e^+ e^- \approx 3.773 \text{ GeV}$
4.40	$\pm 0.66$	$\frac{+0.09}{-0.12}$	47	$\pm 7$	<sup>4</sup> ARTUSO	05A CLEO See EISENSTEIN 08
3.5	$\pm 1.4$	$\pm 0.6$	7	<sup>5</sup> BONVICINI	04A CLEO	Incl. in ARTUSO 05A
8	$\frac{+16}{-5}$	$\frac{+5}{-2}$	1	<sup>6</sup> BAI	98B BES	$e^+ e^- \rightarrow D^* D^-$

<sup>1</sup> ABLIKIM 14F obtain  $|V_{cd}| \cdot f_{D^+} = (45.75 \pm 1.20 \pm 0.39) \text{ MeV}$ , and using  $|V_{cd}| = 0.22520 \pm 0.00065$  gets  $f_{D^+} = (203.2 \pm 5.3 \pm 1.8) \text{ MeV}$ .

<sup>2</sup> EISENSTEIN 08, using the  $D^+$  lifetime and assuming  $|V_{cd}| = |V_{us}|$ , gets  $f_{D^+} = (205.8 \pm 8.5 \pm 2.5) \text{ MeV}$  from this measurement.

<sup>3</sup> ABLIKIM 05D finds a background-subtracted  $2.67 \pm 1.74 D^+ \rightarrow \mu^+ \nu_\mu$  events, and from this obtains  $f_{D^+} = 371 \frac{+129}{-119} \pm 25 \text{ MeV}$ .

<sup>4</sup> ARTUSO 05A obtains  $f_{D^+} = 222.6 \pm 16.7 \frac{+2.8}{-3.4} \text{ MeV}$  from this measurement.

<sup>5</sup> BONVICINI 04A finds eight events with an estimated background of one, and from the branching fraction obtains  $f_{D^+} = 202 \pm 41 \pm 17 \text{ MeV}$ .

<sup>6</sup> BAI 98B obtains  $f_{D^+} = (300 \frac{+180}{-150} \frac{+80}{-40}) \text{ MeV}$  from this measurement.

### $\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.2 \times 10^{-3}</math></b>	90	EISENSTEIN 08	CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<2.1 \times 10^{-3}$	90	RUBIN	06A CLEO	See EISENSTEIN 08

### $\Gamma_{15}/\Gamma$

### $\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.73 <math>\pm 0.10</math> OUR AVERAGE</b>				
8.60 $\pm 0.06$	$\pm 0.15$	26k	ABLIKIM	17S BES3 Using $\bar{K}^0 \rightarrow \pi^+ \pi^-$
8.59 $\pm 0.14$	$\pm 0.21$	5013	ABLIKIM	16V BES3 Using $\bar{K}^0 \rightarrow 2\pi^0$
$8.962 \pm 0.054 \pm 0.206$	40k	<sup>1</sup> ABLIKIM	15AF BES3	from $D^+ \rightarrow K_L e^+ \nu_e$
8.83 $\pm 0.10$	$\pm 0.20$	8.5k	<sup>2</sup> BESSON	09 CLEO from $D^+ \rightarrow K_S e^+ \nu_e$
8.95 $\pm 1.59$	$\pm 0.67$	34	<sup>3</sup> ABLIKIM	05A BES from $D^+ \rightarrow K_S e^+ \nu_e$

### $\Gamma_{16}/\Gamma$

$8.53 \pm 0.13 \pm 0.23$	<sup>4</sup> DOBBS 08	CLEO	See BESSON 09
8.71 $\pm 0.38 \pm 0.37$	545	HUANG	05B CLEO See DOBBS 08

<sup>1</sup> ABLIKIM 15AF report  $\Gamma(D^+ \rightarrow K_L e^+ \nu_e)/\Gamma_{\text{total}} = (4.481 \pm 0.027 \pm 0.103)\%$ . See also the form-factor parameters near the end of this  $D^+$  Listing.

<sup>2</sup> See the form-factor parameters near the end of this  $D^+$  Listing.

<sup>3</sup> The ABLIKIM 05A result together with the  $D^0 \rightarrow K^- e^+ \nu_e$  branching fraction of ABLIKIM 04C and Particle Data Group lifetimes gives  $\Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.08 \pm 0.22 \pm 0.07$ ; isospin invariance predicts the ratio is 1.0.

<sup>4</sup> DOBBS 08 establishes  $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_\pi^\pi(0)}{f_K^K(0)}| = 0.188 \pm 0.008 \pm 0.002$  from the  $D^+$  and  $D^0$  decays to  $\bar{K} e^+ \nu_e$  and  $\pi e^+ \nu_e$ . It also finds  $\Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.06 \pm 0.02 \pm 0.03$ ; isospin invariance predicts the ratio is 1.0.

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8.74 \pm 0.19</math> OUR FIT</b>				

**$8.72 \pm 0.07 \pm 0.18$**       21k      ABLIKIM      16G BES3       $e^+ e^-$  at 3773 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.3 \pm 2.3 \pm 0.8$       29  $\pm$  6      ABLIKIM      07 BES2       $e^+ e^-$  at 3773 MeV

 $\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma(K^- 2\pi^+)$   $\Gamma_{17}/\Gamma_{43}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.97 \pm 0.04</math> OUR FIT</b>				Error includes scale factor of 1.5.
<b><math>1.019 \pm 0.076 \pm 0.065</math></b>	$555 \pm 39$	LINK	04E FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(K^- \pi^+ e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.89 \pm 0.13</math> OUR FIT</b>				Error includes scale factor of 2.1.

**$3.77 \pm 0.03 \pm 0.08$**       18.3k      ABLIKIM      16F BES3       $e^+ e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.50 \pm 0.75 \pm 0.27$       29      ABLIKIM      060 BES2       $e^+ e^-$  at 3773 MeV

$3.5 \begin{matrix} +1.2 \\ -0.7 \end{matrix} \pm 0.4$       14      BAI      91 MRK3       $e^+ e^- \approx 3.77$  GeV

 $\Gamma(K^- \pi^+ e^+ \nu_e)/\Gamma(K^- 2\pi^+)$   $\Gamma_{18}/\Gamma_{43}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.433 \pm 0.011</math> OUR FIT</b>				Error includes scale factor of 2.1.
<b><math>0.4380 \pm 0.0036 \pm 0.0042</math></b>	$70k \pm 363$	DEL-AMO-SA..11l	BABR	$e^+ e^- \approx 10.6$ GeV

 $\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

Unseen decay modes of  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

**$5.40 \pm 0.10$  OUR FIT** Error includes scale factor of 1.1.

**$5.40 \pm 0.10$  OUR AVERAGE** Error includes scale factor of 1.1.

$5.31 \pm 0.05 \pm 0.12$       16.2k      ABLIKIM      16F BES3       $e^+ e^-$  at  $\psi(3770)$

$5.52 \pm 0.07 \pm 0.13$        $\approx 5k$       BRIERE      10 CLEO       $e^+ e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.06 \pm 1.21 \pm 0.40$       28  $\pm$  7      ABLIKIM      060 BES2       $e^+ e^-$  at 3773 MeV

$5.56 \pm 0.27 \pm 0.23$        $422 \pm 21$       <sup>1</sup>HUANG      05B CLEO       $e^+ e^-$  at  $\psi(3770)$

<sup>1</sup>HUANG 05B finds  $\Gamma(D^0 \rightarrow K^{*-} e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e) = 0.98 \pm 0.08 \pm 0.04$ ; isospin invariance predicts the ratio is 1.0.

 $\Gamma((K^- \pi^+)_{[0.8-1.0]\text{GeV}} e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.39 \pm 0.03 \pm 0.08</math></b>	16.2k	ABLIKIM	16F BES3	$e^+ e^-$ at $\psi(3770)$

$$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma(K^- 2\pi^+) \quad \Gamma_{37}/\Gamma_{43}$$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$0.74 \pm 0.04 \pm 0.05$		BRANDENB... 02	CLEO	$e^+ e^- \approx \gamma(4S)$
$0.62 \pm 0.15 \pm 0.09$	35	ADAMOVICH 91	OMEG	$\pi^-$ 340 GeV
$0.55 \pm 0.08 \pm 0.10$	880	ALBRECHT 91	ARG	$e^+ e^- \approx 10.4$ GeV
$0.49 \pm 0.04 \pm 0.05$		ANJOS 89B	E691	Photoproduction

$$\Gamma((K^- \pi^+) S\text{-wave} e^+ \nu_e) / \Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.28 \pm 0.08 \pm 0.08</math></b>	ABLIKIM	16F	$e^+ e^-$ at $\psi(3770)$

$$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- \pi^+ e^+ \nu_e) \quad \Gamma_{19}/\Gamma_{18}$$

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>93.94 \pm 0.27</math> OUR AVERAGE</b>			
$93.93 \pm 0.22 \pm 0.18$	ABLIKIM	16F	$e^+ e^-$ at $\psi(3770)$
$94.11 \pm 0.74 \pm 0.75$	DEL-AMO-SA..11I	BABR	$e^+ e^- \approx 10.6$ GeV

$$\Gamma((K^- \pi^+) S\text{-wave} e^+ \nu_e) / \Gamma(K^- \pi^+ e^+ \nu_e) \quad \Gamma_{21}/\Gamma_{18}$$

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.89 \pm 0.17</math> OUR AVERAGE</b>			
$6.05 \pm 0.22 \pm 0.18$	ABLIKIM	16F	$e^+ e^-$ at $\psi(3770)$
$5.79 \pm 0.16 \pm 0.15$	DEL-AMO-SA..11I	BABR	$e^+ e^- \approx 10.6$ GeV

$$\Gamma(\bar{K}^*(1410)^0 e^+ \nu_e, \bar{K}^*(1410)^0 \rightarrow K^- \pi^+) / \Gamma_{\text{total}} \quad \Gamma_{22}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;6 \times 10^{-3}</math></b>	90	DEL-AMO-SA..11I	BABR	$e^+ e^- \approx 10.6$ GeV

$$\Gamma(\bar{K}_2^*(1430)^0 e^+ \nu_e, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+) / \Gamma_{\text{total}} \quad \Gamma_{23}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;5 \times 10^{-4}</math></b>	90	DEL-AMO-SA..11I	BABR	$e^+ e^- \approx 10.6$ GeV

$$\Gamma(K^- \pi^+ e^+ \nu_e \text{ nonresonant}) / \Gamma_{\text{total}} \quad \Gamma_{24}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;0.007</math></b>	90	ANJOS 89B	E691	Photoproduction

$$\Gamma(K^- \pi^+ \mu^+ \nu_\mu) / \Gamma(\bar{K}^0 \mu^+ \nu_\mu) \quad \Gamma_{25}/\Gamma_{17}$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.417 \pm 0.030 \pm 0.023</math></b>	$555 \pm 39$	LINK	04E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma_{\text{total}} \quad \Gamma_{38}/\Gamma$$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.25 \pm 0.15</math> OUR FIT</b>				
<b><math>5.27 \pm 0.07 \pm 0.14</math></b>	$\approx 5k$	BRIERE	10	CLEO $e^+ e^-$ at $\psi(3770)$

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^0 \mu^+ \nu_\mu)$  $\Gamma_{38}/\Gamma_{17}$ 

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.600 \pm 0.021</math> OUR FIT</b>				
<b><math>0.594 \pm 0.043 \pm 0.033</math></b>	$555 \pm 39$	LINK	04E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(K^- 2\pi^+)$  $\Gamma_{38}/\Gamma_{43}$ 

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.584 \pm 0.025</math> OUR FIT</b>				Error includes scale factor of 1.4.
<b><math>0.57 \pm 0.06</math> OUR AVERAGE</b>				Error includes scale factor of 1.2.
0.72 $\pm 0.10$ $\pm 0.05$		BRANDENB... 02	CLEO	$e^+ e^- \approx \gamma(4S)$
0.56 $\pm 0.04$ $\pm 0.06$	875	FRABETTI 93E	E687	$\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV
0.46 $\pm 0.07$ $\pm 0.08$	224	KODAMA 92C	E653	$\pi^-$ emulsion 600 GeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.602 $\pm 0.010 \pm 0.021$	12k	<sup>1</sup> LINK	02J	FOCS $\gamma$ nucleus, $\approx 180$ GeV

<sup>1</sup>This LINK 02J result includes the effects of an interference of a small  $S$ -wave  $K^- \pi^+$  amplitude with the dominant  $\bar{K}^{*0}$  amplitude. (The interference effect is reported in LINK 02E.) This result is redundant with results of LINK 04E elsewhere in these Listings.

 $\Gamma(K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant}) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$  $\Gamma_{27}/\Gamma_{25}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.0530 \pm 0.0074 \begin{array}{l} +0.0099 \\ -0.0096 \end{array}</math></b>	14k	LINK	05I	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(K^- \pi^+ \pi^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$  $\Gamma_{28}/\Gamma_{25}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.042</b>	90	FRABETTI 93E	E687	$\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV

 $\Gamma(\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$  $\Gamma_{39}/\Gamma_{25}$ 

Unseen decay modes of the  $\bar{K}_0^*(1430)^0$  are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.0064</b>	90	LINK	05I	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\bar{K}^*(1680)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$  $\Gamma_{40}/\Gamma_{25}$ 

Unseen decay modes of the  $\bar{K}^*(1680)^0$  are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.04</b>	90	LINK	05I	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

### $\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$

$\Gamma_{29}/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.372 <math>\pm 0.017</math> OUR AVERAGE</b>	Error includes scale factor of 2.0.			
0.363 $\pm 0.008 \pm 0.005$	3.4k	ABLIKIM 17S	BES3	Using $\pi^0 \rightarrow 2\gamma$
0.405 $\pm 0.016 \pm 0.009$	838	<sup>1</sup> BESSON 09	CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.373 $\pm 0.022 \pm 0.013$		<sup>2</sup> DOBBS 08	CLEO	See BESSON 09
0.44 $\pm 0.06 \pm 0.03$	63 $\pm$ 9	HUANG 05B	CLEO	See DOBBS 08

<sup>1</sup> See the form-factor parameters near the end of this  $D^+$  Listing.

<sup>2</sup> DOBBS 08 establishes  $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}| = 0.188 \pm 0.008 \pm 0.002$  from the  $D^+$  and  $D^0$  decays to  $\bar{K}e^+ \nu_e$  and  $\pi e^+ \nu_e$ . It finds  $\Gamma(D^0 \rightarrow \pi^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \pi^0 e^+ \nu_e) = 2.03 \pm 0.14 \pm 0.08$ ; isospin invariance predicts the ratio is 2.0.

### $\Gamma(\eta e^+ \nu_e)/\Gamma_{\text{total}}$

$\Gamma_{30}/\Gamma$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.4 <math>\pm 0.9 \pm 0.4</math></b>		YELTON 11	CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

13.3  $\pm 2.0 \pm 0.6$       46  $\pm$  8      MITCHELL 09B      CLEO      See YELTON 11

### $\Gamma(\rho^0 e^+ \nu_e)/\Gamma_{\text{total}}$

$\Gamma_{31}/\Gamma$

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.18 <math>\pm 0.17 \pm 0.25</math> OUR FIT</b>				

**2.17  $\pm 0.12 \pm 0.22$**       447  $\pm$  25      <sup>1</sup> DOBBS      13      CLEO       $e^+ e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1  $\pm 0.4 \pm 0.1$       27  $\pm$  6      <sup>2</sup> HUANG      05B      CLEO      See DOBBS 13

<sup>1</sup> DOBBS 13 finds  $\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e) / 2 \Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e) = 1.03 \pm 0.09^{+0.08}_{-0.02}$ ; isospin invariance predicts the ratio is 1.0.

<sup>2</sup> HUANG 05B finds  $\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e) / 2 \Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e) = 1.2^{+0.4}_{-0.3} \pm 0.1$ ; isospin invariance predicts the ratio is 1.0.

### $\Gamma(\rho^0 e^+ \nu_e)/\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)$

$\Gamma_{31}/\Gamma_{37}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0404 <math>\pm 0.0033 \pm 0.0050</math> OUR FIT</b>				

**0.045  $\pm 0.014 \pm 0.009$**       49      <sup>1</sup> AITALA      97      E791       $\pi^-$  nucleus, 500 GeV

<sup>1</sup> AITALA 97 explicitly subtracts  $D^+ \rightarrow \eta' e^+ \nu_e$  and other backgrounds to get this result.

### $\Gamma(\rho^0 \mu^+ \nu_\mu)/\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$

$\Gamma_{32}/\Gamma_{38}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.045 <math>\pm 0.007</math> OUR AVERAGE</b>	Error includes scale factor of 1.1.			

0.041  $\pm 0.006 \pm 0.004$       320  $\pm$  44      LINK      06B      FOCS       $\gamma$  A,  $\bar{E}_\gamma \approx 180$  GeV

0.051  $\pm 0.015 \pm 0.009$       54      <sup>1</sup> AITALA      97      E791       $\pi^-$  nucleus, 500 GeV

0.079  $\pm 0.019 \pm 0.013$       39      <sup>2</sup> FRABETTI      97      E687       $\gamma$  Be,  $\bar{E}_\gamma \approx 220$  GeV

<sup>1</sup> AITALA 97 explicitly subtracts  $D^+ \rightarrow \eta' \mu^+ \nu_\mu$  and other backgrounds to get this result.

<sup>2</sup> Because the reconstruction efficiency for photons is low, this FRABETTI 97 result also includes any  $D^+ \rightarrow \eta' \mu^+ \nu_\mu \rightarrow \gamma \rho^0 \mu^+ \nu_\mu$  events in the numerator.

### $\Gamma(\omega e^+ \nu_e)/\Gamma_{\text{total}}$

$\Gamma_{33}/\Gamma$

<u>VALUE</u> (units $10^{-3}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.69 <math>\pm</math> 0.11 OUR AVERAGE</b>				
1.63 $\pm$ 0.11 $\pm$ 0.08	491 $\pm$ 32	ABLIKIM	15W BES3	292 $\text{fb}^{-1}$ , 3773 MeV
1.82 $\pm$ 0.18 $\pm$ 0.07	129 $\pm$ 13	DOBBS	13 CLEO	$e^+ e^-$ at $\psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.6 $^{+0.7}_{-0.6}$ $\pm$ 0.1	7.6 $^{+3.3}_{-2.7}$	HUANG	05B CLEO	See DOBBS 13

### $\Gamma(\eta'(958)e^+\nu_e)/\Gamma_{\text{total}}$

$\Gamma_{34}/\Gamma$

<u>VALUE</u> (units $10^{-4}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.16 <math>\pm</math> 0.53 <math>\pm</math> 0.07</b>				
YELTON 11 CLEO $e^+ e^-$ at $\psi(3770)$				

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.5

90

MITCHELL

09B

CLEO

See YELTON 11

### $\Gamma(\phi e^+\nu_e)/\Gamma_{\text{total}}$

$\Gamma_{35}/\Gamma$

Unseen decay modes of the  $\phi$  are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;1.3 \times 10^{-5}</math></b>	90	ABLIKIM	15W BES3	292 $\text{fb}^{-1}$ , 3773 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.9 $\times 10^{-4}$ 90 YELTON 11 CLEO $e^+ e^-$ at $\psi(3770)$				
<1.6 $\times 10^{-4}$	90	MITCHELL	09B CLEO	See YELTON 11
<0.0201	90	ABLIKIM	06P BES2	$e^+ e^-$ at 3773 MeV
<0.0209	90	BAI	91 MRK3	$e^+ e^- \approx 3.77 \text{ GeV}$

### $\Gamma(D^0 e^+\nu_e)/\Gamma_{\text{total}}$

$\Gamma_{36}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;1.0 \times 10^{-4}</math></b>	90	ABLIKIM	17AD BES3	$e^+ e^-$ at 3.773 GeV

—— Hadronic modes with a  $\bar{K}$  or  $\bar{K}K\bar{K}$  ——

### $\Gamma(K_S^0\pi^+)/\Gamma_{\text{total}}$

$\Gamma_{41}/\Gamma$

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.526 $\pm$ 0.022 $\pm$ 0.038		<sup>1</sup> DOBBS 07	CLEO	See MENDEZ 10
1.55 $\pm$ 0.05 $\pm$ 0.06	2.2k	<sup>1</sup> HE 05	CLEO	See DOBBS 07
1.6 $\pm$ 0.3 $\pm$ 0.1	161	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

### $\Gamma(K_S^0\pi^+)/\Gamma(K^-2\pi^+)$

$\Gamma_{41}/\Gamma_{43}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.164 <math>\pm</math> 0.007 OUR FIT</b>		Error includes scale factor of 3.9.		
<b>0.162 <math>\pm</math> 0.009 OUR AVERAGE</b>		Error includes scale factor of 4.5.		
0.171 $\pm$ 0.002 $\pm$ 0.002		BONVICINI 14	CLEO	All CLEO-c runs
0.1530 $\pm$ 0.0023 $\pm$ 0.0016	10.6k	LINK 02B	FOCS	$\gamma$ nucleus, $\overline{E}_\gamma \approx 180 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.1682 \pm 0.0012 \pm 0.0037$	30k	MENDEZ	10	CLEO	See BONVICINI 14
$0.174 \pm 0.012 \pm 0.011$	473	<sup>1</sup> BISHAI	97	CLEO	$e^+ e^- \approx \Upsilon(4S)$
$0.137 \pm 0.015 \pm 0.016$	264	ANJOS	90C	E691	Photoproduction

<sup>1</sup> See BISHAI 97 for an isospin analysis of  $D^+ \rightarrow \bar{K}\pi$  amplitudes.

### $\Gamma(K_L^0 \pi^+)/\Gamma_{\text{total}}$

### $\Gamma_{42}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.460 \pm 0.040 \pm 0.035</math></b>	$2023 \pm 54$	<sup>1</sup> HE	08	CLEO $e^+ e^-$ at $\psi(3770)$

<sup>1</sup> The difference of CLEO  $D^+ \rightarrow K_S^0 \pi^+$  and  $K_L^0 \pi^+$  branching fractions over the sum (DOBBS 07 and HE 08) is  $+0.022 \pm 0.016 \pm 0.018$ .

### $\Gamma(K^- 2\pi^+)/\Gamma_{\text{total}}$

### $\Gamma_{43}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>8.98 \pm 0.28</math> OUR FIT</b>		Error includes scale factor of 2.2.		

**$9.224 \pm 0.059 \pm 0.157$**  BONVICINI 14 CLEO All CLEO-c runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.14 \pm 0.10 \pm 0.17$		<sup>1</sup> DOBBS	07	CLEO See BONVICINI 14
$9.5 \pm 0.2 \pm 0.3$	15.1k	<sup>1</sup> HE	05	CLEO See DOBBS 07
$9.3 \pm 0.6 \pm 0.8$	1502	<sup>2</sup> BALEST	94	CLEO $e^+ e^- \approx \Upsilon(4S)$
$6.4 \begin{matrix} +1.5 \\ -1.4 \end{matrix}$		<sup>3</sup> BARLAG	92C	ACCM $\pi^-$ Cu 230 GeV
$9.1 \pm 1.3 \pm 0.4$	1164	ADLER	88C	MRK3 $e^+ e^-$ 3.77 GeV
$9.1 \pm 1.9$	239	<sup>4</sup> SCHINDLER	81	MRK2 $e^+ e^-$ 3.771 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

<sup>2</sup> BALEST 94 measures the ratio of  $D^+ \rightarrow K^- \pi^+ \pi^+$  and  $D^0 \rightarrow K^- \pi^+$  branching fractions to be  $2.35 \pm 0.16 \pm 0.16$  and uses their absolute measurement of the  $D^0 \rightarrow K^- \pi^+$  fraction (AKERIB 93).

<sup>3</sup> BARLAG 92C computes the branching fraction by topological normalization.

<sup>4</sup> SCHINDLER 81 (MARK-2) measures  $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$  branching fraction to be  $0.38 \pm 0.05$  nb. We use the MARK-3 (ADLER 88C) value of  $\sigma = 4.2 \pm 0.6 \pm 0.3$  nb.

## See the related review(s):

### Review of Multibody Charm Analyses

### $\Gamma((K^- \pi^+)_{S-\text{wave}} \pi^+)/\Gamma(K^- 2\pi^+)$

### $\Gamma_{44}/\Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis. The  $K^- \pi^+$  S-wave includes a broad scalar  $\kappa$  ( $\bar{K}_0^*(700)$ ), the  $\bar{K}_0^*(1430)^0$ , and non-resonant background.

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.801 \pm 0.012</math> OUR AVERAGE</b>			
$0.8024 \pm 0.0138 \pm 0.0043$	<sup>1</sup> LINK	09	FOCS MIPWA fit, 53k evts
$0.838 \pm 0.038$	<sup>2</sup> BONVICINI	08A	CLEO QMIPWA fit, 141k evts
$0.786 \pm 0.014 \pm 0.018$	AITALA	06	E791 Dalitz fit, 15.1k events
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.8323 \pm 0.0150 \pm 0.0008$	<sup>3</sup> LINK	07B	FOCS See LINK 09

- <sup>1</sup>This LINK 09 model-independent partial-wave analysis of the  $K^- \pi^+$  S-wave slices the  $K^- \pi^+$  mass range into 39 bins.  
<sup>2</sup>The BONVICINI 08A QMIPWA (quasi-model-independent partial-wave analysis) of the  $K^- \pi^+$  S-wave amplitude slices the  $K^- \pi^+$  mass range into 26 bins but keeps the Breit-Wigner  $\bar{K}_0^*(1430)^0$ .  
<sup>3</sup>This LINK 07B fit uses a K matrix. The  $K^- \pi^+$  S-wave fit fraction given above breaks down into  $(207.3 \pm 25.5 \pm 12.4)\%$  isospin-1/2 and  $(40.5 \pm 9.6 \pm 3.2)\%$  isospin-3/2 — with large interference between the two. The isospin-1/2 component includes the  $\kappa$  (or  $\bar{K}_0^*(700)^0$ ) and  $\bar{K}_0^*(1430)^0$ .

### $\Gamma(\bar{K}_0^*(700)^0 \pi^+, \bar{K}_0^*(700) \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$ $\Gamma_{45}/\Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
$0.478 \pm 0.121 \pm 0.053$	AITALA	02 E791	See AITALA 06

### $\Gamma(\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$ $\Gamma_{47}/\Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.111 ± 0.012 OUR AVERAGE</b> Error includes scale factor of 3.7.			
$0.1236 \pm 0.0034 \pm 0.0034$	LINK	09 FOCS	MIPWA fit, 53k evts
$0.0988 \pm 0.0046$	BONVICINI	08A CLEO	QMIPWA fit, 141k evts
$0.119 \pm 0.002 \pm 0.020$	AITALA	06 E791	Dalitz fit, 15.1k events
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
$0.1361 \pm 0.0041 \pm 0.0030$	<sup>1</sup> LINK	07B FOCS	See LINK 09
$0.123 \pm 0.010 \pm 0.009$	AITALA	02 E791	See AITALA 06
$0.137 \pm 0.006 \pm 0.009$	FRABETTI	94G E687	Dalitz fit, 8800 evts
$0.170 \pm 0.009 \pm 0.034$	ANJOS	93 E691	$\gamma$ Be 90–260 GeV
$0.14 \pm 0.04 \pm 0.04$	ALVAREZ	91B NA14	Photoproduction
$0.13 \pm 0.01 \pm 0.07$	ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV

<sup>1</sup>The statistical error on this LINK 07B value is corrected in LINK 09.

### $\Gamma(\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$ $\Gamma_{48}/\Gamma_{43}$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>			
<b>not seen</b>			
$4.8 \pm 2.1 \pm 1.7$	LINK	07B FOCS	See LINK 09

### $\Gamma(\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$ $\Gamma_{46}/\Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.1330 ± 0.0062</b>			
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
$0.125 \pm 0.014 \pm 0.005$	AITALA	02 E791	See AITALA 06
$0.284 \pm 0.022 \pm 0.059$	FRABETTI	94G E687	Dalitz fit, 8800 evts
$0.248 \pm 0.019 \pm 0.017$	ANJOS	93 E691	$\gamma$ Be 90–260 GeV


$\Gamma(\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$

$\Gamma_{49}/\Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.24 ± 0.08 OUR AVERAGE</b>			Error includes scale factor of 2.2. See the ideogram below.
0.58 ± 0.10 ± 0.06	LINK 09	FOCS	MIPWA fit, 53k evts
0.204 ± 0.040	BONVICINI 08A	CLEO	QMIPWA fit, 141k evts
0.2 ± 0.1 ± 0.1	AITALA 06	E791	Dalitz fit, 15.1k events
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.39 ± 0.09 ± 0.05	LINK 07B	FOCS	See LINK 09
0.5 ± 0.1 ± 0.2	AITALA 02	E791	See AITALA 06

WEIGHTED AVERAGE  
0.24±0.08 (Error scaled by 2.2)



$$\Gamma(\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+) \quad \Gamma_{49}/\Gamma_{43}$$

(units  $10^{-2}$ )

$\Gamma(\bar{K}^*(1680)^0 \pi^+, \bar{K}^*(1680)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$

$\Gamma_{50}/\Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.23 ± 0.12 OUR AVERAGE</b>			
1.75 ± 0.62 ± 0.54	LINK 09	FOCS	MIPWA fit, 53k evts
0.196 ± 0.118	BONVICINI 08A	CLEO	QMIPWA fit, 141k evts
1.2 ± 0.6 ± 1.2	AITALA 06	E791	Dalitz fit, 15.1k events
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.90 ± 0.63 ± 0.43	LINK 07B	FOCS	See LINK 09
2.5 ± 0.7 ± 0.3	AITALA 02	E791	See AITALA 06
4.7 ± 0.6 ± 0.7	FRAZETTI 94G	E687	Dalitz fit, 8800 evts
3.0 ± 0.4 ± 1.3	ANJOS 93	E691	$\gamma$ Be 90–260 GeV

$\Gamma(K^-(2\pi^+)_{I=2})/\Gamma(K^-2\pi^+)$  $\Gamma_{51}/\Gamma_{43}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.155±0.028</b>	BONVICINI	08A	CLEO QMIPWA fit, 141k evts

 $\Gamma(K^-2\pi^+ \text{ nonresonant})/\Gamma(K^-2\pi^+)$  $\Gamma_{52}/\Gamma_{43}$ 

This is the “fit fraction” from the Dalitz-plot analysis. Later analyses find little need for this decay mode.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.130±0.058±0.044	AITALA	02	E791 See AITALA 06
0.998±0.037±0.072	FRABETTI	94G	E687 Dalitz fit, 8800 evts
0.838±0.088±0.275	ANJOS	93	E691 $\gamma$ Be 90–260 GeV
0.79 ±0.07 ±0.15	ADLER	87	MRK3 $e^+ e^-$ 3.77 GeV

 $\Gamma(K_S^0\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.99±0.09±0.25		1 DOBBS	07	CLEO See BONVICINI 14
7.2 ±0.2 ±0.4	5.1k	1 HE	05	CLEO See DOBBS 07
5.1 ±1.3 ±0.8	159	ADLER	88C	MRK3 $e^+ e^-$ 3.77 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

 $\Gamma(K_S^0\pi^+\pi^0)/\Gamma(K^-2\pi^+)$  $\Gamma_{53}/\Gamma_{43}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.785±0.007±0.016</b>	BONVICINI	14	CLEO All CLEO-c runs

 $\Gamma(K_S^0\rho^+)/\Gamma(K_S^0\pi^+\pi^0)$  $\Gamma_{54}/\Gamma_{53}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>83.4±2.2<sup>+7.1</sup><sub>-3.6</sub></b>	1 ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

68 ±8 ±12 ADLER 87 MRK3  $e^+ e^-$  3.77 GeV

<sup>1</sup> Fit fraction from Dalitz plot analysis of 142k  $D^+ \rightarrow K_S^0\pi^+\pi^0$  events.

 $\Gamma(K_S^0\rho(1450)^+, \rho^+ \rightarrow \pi^+\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$  $\Gamma_{55}/\Gamma_{53}$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>2.1±0.3<sup>+1.6</sup><sub>-1.9</sub></b>	ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

 $\Gamma(\bar{K}^*(892)^0\pi^+, \bar{K}^*(892)^0 \rightarrow K_S^0\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$  $\Gamma_{56}/\Gamma_{53}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.58±0.17<sup>+0.39</sup><sub>-0.38</sub></b>	1 ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

19 ±6 ±6 ADLER 87 MRK3  $e^+ e^-$  3.77 GeV

<sup>1</sup> Fit fraction from Dalitz plot analysis of 142k  $D^+ \rightarrow K_S^0\pi^+\pi^0$  events.

$\Gamma(\bar{K}_0^*(1430)^0\pi^+, \bar{K}_0^{*0} \rightarrow K_S^0\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{57}/\Gamma_{53}$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$3.7 \pm 0.6 \pm 1.1$	ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

 $\Gamma(\bar{K}_0^*(1680)^0\pi^+, \bar{K}_0^{*0} \rightarrow K_S^0\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{58}/\Gamma_{53}$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$1.3 \pm 0.2^{+0.9}_{-1.3}$	ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

 $\Gamma(\bar{\kappa}^0\pi^+, \bar{\kappa}^0 \rightarrow K_S^0\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{59}/\Gamma_{53}$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$7.7 \pm 1.2^{+6.5}_{-4.8}$	ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

 $\Gamma(K_S^0\pi^+\pi^0 \text{ nonresonant})/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{60}/\Gamma_{53}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 0.7^{+5.4}_{-5.1}$	<sup>1</sup> ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13  $\pm 7$   $\pm 8$  ADLER 87 MRK3  $e^+ e^-$  3.77 GeV

<sup>1</sup> Fit fraction from Dalitz plot analysis of 142k  $D^+ \rightarrow K_S^0\pi^+\pi^0$  events.

 $\Gamma(K_S^0\pi^+\pi^0 \text{ nonresonant and } \bar{\kappa}^0\pi^+)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{61}/\Gamma_{53}$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$18.6 \pm 1.7^{+2.3}_{-4.6}$	ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

 $\Gamma((K_S^0\pi^0)_{S-\text{wave}}\pi^+)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{62}/\Gamma_{53}$ 

The numerator here is the coherent sum of the  $\bar{K}_0^*(1430)^0\pi^+$ ,  $\bar{\kappa}^0\pi^+$ , and nonresonant contributions.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$17.3 \pm 1.4^{+3.4}_{-4.3}$	ABLIKIM	14E	BES3 $e^+ e^-$ at $\psi(3770)$

 $\Gamma(K^-2\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

See our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode. There is nothing new since 1992, and the two papers, ANJOS 92C, with  $91 \pm 12$  events above background, and COFFMAN 92B, with  $142 \pm 20$  such events, could not determine submode fractions with much accuracy.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.98 \pm 0.08 \pm 0.16$		<sup>1</sup> DOBBS 07 CLEO		See BONVICINI 14
$6.0 \pm 0.2 \pm 0.2$	4.8k	<sup>1</sup> HE 05 CLEO		See DOBBS 07
$5.8 \pm 1.2 \pm 1.2$	142	COFFMAN 92B MRK3	$e^+ e^-$ 3.77 GeV	
$6.3^{+1.4}_{-1.3} \pm 1.2$	175	BALTRUSAIT..86E MRK3		See COFFMAN 92B

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$\Gamma(K^- 2\pi^+ \pi^0)/\Gamma(K^- 2\pi^+)$ 
 $\Gamma_{63}/\Gamma_{43}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.666±0.006±0.014</b>	BONVICINI 14	CLEO	All CLEO-c runs

 $\Gamma(K_S^0 2\pi^+ \pi^-)/\Gamma_{\text{total}}$ 
 $\Gamma_{64}/\Gamma$ 

See our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode. There is nothing new since 1992, and the two papers, ANJOS 92C, with  $229 \pm 17$  events above background, and COFFMAN 92B, with  $209 \pm 20$  such events, could not determine submode fractions with much accuracy.

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

$3.122 \pm 0.046 \pm 0.096$	${}^1 \text{DOBBS}$	07	CLEO	See BONVICINI 14
$3.2 \pm 0.1 \pm 0.2$	3.2k	${}^1 \text{HE}$	05	CLEO See DOBBS 07
$2.1^{+1.0}_{-0.9}$		${}^2 \text{BARLAG}$	92C	ACCM $\pi^-$ Cu 230 GeV
$3.3 \pm 0.8 \pm 0.2$	168	ADLER	88C	MRK3 $e^+ e^-$ 3.77 GeV

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

<sup>2</sup> BARLAG 92C computes the branching fraction by topological normalization.

 $\Gamma(K_S^0 2\pi^+ \pi^-)/\Gamma(K^- 2\pi^+)$ 
 $\Gamma_{64}/\Gamma_{43}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.331±0.004±0.006</b>	BONVICINI 14	CLEO	All CLEO-c runs

 $\Gamma(K^- 3\pi^+ \pi^-)/\Gamma(K^- 2\pi^+)$ 
 $\Gamma_{65}/\Gamma_{43}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.061±0.005 OUR FIT</b>				Error includes scale factor of 1.1.
<b>0.062±0.008 OUR AVERAGE</b>				Error includes scale factor of 1.3.
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.058 \pm 0.002 \pm 0.006$	2923	LINK	03D FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
$0.077 \pm 0.008 \pm 0.010$	239	FRABETTI	97C E687	$\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV
$0.09 \pm 0.01 \pm 0.01$	113	ANJOS	90D E691	Photoproduction

 $\Gamma(\bar{K}^*(892)^0 2\pi^+ \pi^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 3\pi^+ \pi^-)$ 
 $\Gamma_{66}/\Gamma_{65}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.21±0.04±0.06</b>	LINK	03D FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 3\pi^+ \pi^-)$ 
 $\Gamma_{67}/\Gamma_{65}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.40±0.03±0.06</b>	LINK	03D FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$ 
 $\Gamma_{67}/\Gamma_{43}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.016 \pm 0.007 \pm 0.004$	FRABETTI	97C E687	$\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(\bar{K}^*(892)^0 2\pi^+ \pi^- \text{ no-}\rho, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+) \quad \Gamma_{69}/\Gamma_{43}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
0.032 ± 0.010 ± 0.008	FRABETTI 97C E687	γ Be, $\bar{E}_\gamma$ ≈ 200 GeV	

$$\Gamma(K^- \rho^0 2\pi^+)/\Gamma(K^- 3\pi^+ \pi^-) \quad \Gamma_{70}/\Gamma_{65}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.30 ± 0.04 ± 0.01</b>	LINK 03D FOCS	γ A, $\bar{E}_\gamma$ ≈ 180 GeV	

$$\Gamma(K^- \rho^0 2\pi^+)/\Gamma(K^- 2\pi^+) \quad \Gamma_{70}/\Gamma_{43}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
0.034 ± 0.009 ± 0.005	FRABETTI 97C E687	γ Be, $\bar{E}_\gamma$ ≈ 200 GeV	

$$\Gamma(\bar{K}^*(892)^0 a_1(1260)^+)/\Gamma(K^- 2\pi^+) \quad \Gamma_{68}/\Gamma_{43}$$

Unseen decay modes of the  $\bar{K}^*(892)^0$  and  $a_1(1260)^+$  are included.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.099 ± 0.008 ± 0.018</b>	LINK 03D FOCS	γ A, $\bar{E}_\gamma$ ≈ 180 GeV	

$$\Gamma(K^- 3\pi^+ \pi^- \text{ nonresonant})/\Gamma(K^- 3\pi^+ \pi^-) \quad \Gamma_{71}/\Gamma_{65}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.07 ± 0.05 ± 0.01</b>		LINK 03D FOCS	γ A, $\bar{E}_\gamma$ ≈ 180 GeV	
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
<0.026	90	FRABETTI 97C E687	γ Be, $\bar{E}_\gamma$ ≈ 200 GeV	

$$\Gamma(K^+ 2K_S^0)/\Gamma_{\text{total}} \quad \Gamma_{72}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.4 ± 0.5 ± 1.2</b>	3551	ABLIKIM 17A BES3	e <sup>+</sup> e <sup>-</sup> → $\psi(3770)$	

$$\Gamma(K^+ 2K_S^0)/\Gamma(K^- 2\pi^+) \quad \Gamma_{72}/\Gamma_{43}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
0.035 ± 0.010 ± 0.005	39 ± 9	ALBRECHT 94I ARG	e <sup>+</sup> e <sup>-</sup> ≈ 10 GeV	
0.085 ± 0.018	70 ± 12	AMMAR 91 CLEO	e <sup>+</sup> e <sup>-</sup> ≈ 10.5 GeV	

$$\Gamma(K^+ K^- K_S^0 \pi^+)/\Gamma(K_S^0 2\pi^+ \pi^-) \quad \Gamma_{73}/\Gamma_{64}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.7 ± 1.5 ± 0.9</b>	35 ± 7	LINK 01C FOCS	γ nucleus, $\bar{E}_\gamma$ ≈ 180 GeV	

### Pionic modes


$$\Gamma(\pi^+ \pi^0)/\Gamma(K^- 2\pi^+) \quad \Gamma_{74}/\Gamma_{43}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.31 ± 0.06 OUR AVERAGE</b>				
1.29 ± 0.04 ± 0.05	2649 ± 76	MENDEZ 10 CLEO	e <sup>+</sup> e <sup>-</sup> at 3774 MeV	
1.33 ± 0.11 ± 0.09	1229 ± 99	AUBERT,B 06F BABR	e <sup>+</sup> e <sup>-</sup> ≈ $\Upsilon(4S)$	
1.44 ± 0.19 ± 0.10	171 ± 22	ARMS 04 CLEO	e <sup>+</sup> e <sup>-</sup> ≈ 10 GeV	
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
1.33 ± 0.07 ± 0.06	914 ± 46	RUBIN 06 CLEO	See MENDEZ 10	

$\Gamma(2\pi^+\pi^-)/\Gamma(K^-\bar{2}\pi^+)$  $\Gamma_{75}/\Gamma_{43}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.48±0.19 OUR AVERAGE</b>	Error includes scale factor of 1.4. See the ideogram below.			
3.52±0.11±0.12	3303 ± 95	RUBIN	06	CLEO $e^+e^-$ at $\psi(3770)$
4.1 ± 1.1 ± 0.3	85 ± 22	ABLIKIM	05F	BES $e^+e^- \approx \psi(3770)$
3.11±0.18 <sup>+0.16</sup> <sub>-0.26</sub>	1172	AITALA	01B E791	$\pi^-$ nucleus, 500 GeV
4.3 ± 0.3 ± 0.3	236	FRAZETTI	97D E687	$\gamma$ Be ≈ 200 GeV
3.5 ± 0.7 ± 0.3	83	ANJOS	89 E691	Photoproduction

WEIGHTED AVERAGE  
 $3.48\pm0.19$  (Error scaled by 1.4)




$$\Gamma(2\pi^+\pi^-)/\Gamma(K^-\bar{2}\pi^+) \text{ (units } 10^{-2})$$

 $\Gamma(\rho^0\pi^+)/\Gamma(2\pi^+\pi^-)$  $\Gamma_{76}/\Gamma_{75}$ 

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.25 ±0.04 OUR AVERAGE</b>	Error includes scale factor of 2.4. See the ideogram below.		
0.200 ± 0.023 ± 0.009	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts
0.3082±0.0314±0.0230	LINK	04	FOCS Dalitz fit, $1527 \pm 51$ evts
0.336 ± 0.032 ± 0.022	AITALA	01B E791	Dalitz fit, 1172 evts



### $\Gamma(\pi^+(\pi^+\pi^-)S\text{-wave})/\Gamma(2\pi^+\pi^-)$

### $\Gamma_{77}/\Gamma_{75}$

This is the “fit fraction” from the Dalitz-plot analysis. See also the next three data blocks.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.5600±0.0324±0.0214</b>	1 LINK	04	FOCS Dalitz fit, 1527 ± 51 evts

<sup>1</sup>LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full  $\pi\pi$  S-wave isoscalar scattering amplitude to describe the  $\pi^+\pi^-$  S-wave component of the  $\pi^+\pi^+\pi^-$  state. The fit fraction given above is a sum over five  $f_0$  mesons, the  $f_0(980)$ ,  $f_0(1300)$ ,  $f_0(1200\text{--}1600)$ ,  $f_0(1500)$ , and  $f_0(1750)$ . See LINK 04 for details and discussion.

### $\Gamma(\sigma\pi^+, \sigma \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$

### $\Gamma_{78}/\Gamma_{75}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.422±0.027 OUR AVERAGE</b>			
0.418±0.014±0.025	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts
0.463±0.090±0.021	ITALA	01B	E791 Dalitz fit, 1172 evts

### $\Gamma(f_0(980)\pi^+, f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$

### $\Gamma_{79}/\Gamma_{75}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.048±0.010 OUR AVERAGE</b>			Error includes scale factor of 1.3.
0.041±0.009±0.003	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts
0.062±0.013±0.004	ITALA	01B	E791 Dalitz fit, 1172 evts

$$\Gamma(f_0(1370)\pi^+, f_0(1370) \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ \pi^-) \quad \Gamma_{80}/\Gamma_{75}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.024±0.013 OUR AVERAGE</b>			
0.026±0.018±0.006	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts
0.023±0.015±0.008	AITALA	01B E791	Dalitz fit, 1172 evts


$$\Gamma(f_2(1270)\pi^+, f_2(1270) \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ \pi^-) \quad \Gamma_{81}/\Gamma_{75}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.154 ± 0.025 OUR AVERAGE</b>			Error includes scale factor of 1.9. See the ideogram below.
0.182 ± 0.026 ± 0.007	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts
0.1174±0.0190±0.0029	LINK	04	FOCS Dalitz fit, 1527 ± 51 evts
0.194 ± 0.025 ± 0.004	AITALA	01B E791	Dalitz fit, 1172 evts

WEIGHTED AVERAGE

0.154±0.025 (Error scaled by 1.9)



$$\Gamma(f_2(1270)\pi^+, f_2(1270) \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ \pi^-)$$

$$\Gamma(\rho(1450)^0\pi^+, \rho(1450)^0 \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ \pi^-) \quad \Gamma_{82}/\Gamma_{75}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.024	95	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.007±0.007±0.003		AITALA	01B E791	Dalitz fit, 1172 evts

$$\Gamma(f_0(1500)\pi^+, f_0(1500) \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ \pi^-) \quad \Gamma_{83}/\Gamma_{75}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.034±0.010±0.008</b>	BONVICINI	07	CLEO Dalitz fit, ≈ 2240 evts

$\Gamma(f_0(1710)\pi^+, f_0(1710) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{84}/\Gamma_{75}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.016	95	BONVICINI	07	CLEO Dalitz fit, $\approx 2240$ evts

 $\Gamma(f_0(1790)\pi^+, f_0(1790) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{85}/\Gamma_{75}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	95	BONVICINI	07	CLEO Dalitz fit, $\approx 2240$ evts

 $\Gamma((\pi^+\pi^+)_{S-\text{wave}}\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{86}/\Gamma_{75}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.037	95	BONVICINI	07	CLEO Dalitz fit, $\approx 2240$ evts

 $\Gamma(2\pi^+\pi^- \text{ nonresonant})/\Gamma(2\pi^+\pi^-)$   $\Gamma_{87}/\Gamma_{75}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.035	95	BONVICINI	07	CLEO Dalitz fit, $\approx 2240$ evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.078  $\pm$  0.060  $\pm$  0.027 AITALA 01B E791 Dalitz fit, 1172 evts $\Gamma(\pi^+ 2\pi^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{88}/\Gamma_{43}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.0 <math>\pm</math> 0.3 <math>\pm</math> 0.3</b>	1535 $\pm$ 89	RUBIN	06	CLEO $e^+ e^-$ at $\psi(3770)$

 $\Gamma(2\pi^+\pi^-\pi^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{89}/\Gamma_{43}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.4 <math>\pm</math> 0.5 <math>\pm</math> 0.6</b>	5701 $\pm$ 205	RUBIN	06	CLEO $e^+ e^-$ at $\psi(3770)$

 $\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$ Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>33.3 <math>\pm</math> 2.1 OUR FIT</b>				Error includes scale factor of 1.4.

**30.7  $\pm$  2.2  $\pm$  1.3** 258 ABLIKIM 16D BES3  $e^+ e^-$  at 3773 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

34.3  $\pm$  1.4  $\pm$  1.7 1033  $\pm$  42 ARTUSO 08 CLEO See MENDEZ 10 $\Gamma(\eta\pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{91}/\Gamma_{43}$ Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.71 <math>\pm</math> 0.23 OUR FIT</b>				Error includes scale factor of 1.3.

**3.87  $\pm$  0.09  $\pm$  0.19** 2940  $\pm$  68 MENDEZ 10 CLEO  $e^+ e^-$  at 3774 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.81  $\pm$  0.26  $\pm$  0.21 377  $\pm$  26 RUBIN 06 CLEO See ARTUSO 08



$\Gamma(K^+ K_S^0)/\Gamma(K_S^0 \pi^+)$   $\Gamma_{96}/\Gamma_{41}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.192 ± 0.006 OUR FIT</b>	Error includes scale factor of 2.6.			
<b>0.1901 ± 0.0024 OUR AVERAGE</b>				
0.1899 ± 0.0011 ± 0.0022	101k ± 561	WON	09	BELL $e^+ e^-$ at $\Upsilon(4S)$
0.1892 ± 0.0155 ± 0.0073	278 ± 21	ARMS	04	CLEO $e^+ e^- \approx 10$ GeV
0.1996 ± 0.0119 ± 0.0096	949	LINK	02B	FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
0.222 ± 0.037 ± 0.013	63 ± 10	ABLIKIM	05F	BES $e^+ e^- \approx \psi(3770)$
0.222 ± 0.041 ± 0.019	70	BISHAI	97	CLEO See ARMS 04
0.25 ± 0.04 ± 0.02	129	FRABETTI	95	E687 $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV
0.271 ± 0.065 ± 0.039	69	ANJOS	90C	E691 $\gamma$ Be
0.317 ± 0.086 ± 0.048	31	BALTRUSAIT..85E	MRK3	$e^+ e^-$ 3.77 GeV
0.25 ± 0.15	6	SCHINDLER	81	MRK2 $e^+ e^-$ 3.771 GeV

 $\Gamma(K^+ K_S^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{96}/\Gamma_{43}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.15 ± 0.15 OUR FIT</b>	Error includes scale factor of 3.2.			
<b>3.35 ± 0.06 ± 0.07</b>				
5161 ± 86	MENDEZ	10	CLEO	$e^+ e^-$ at 3774 MeV
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
3.02 ± 0.18 ± 0.15	949	<sup>1</sup> LINK	02B	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

<sup>1</sup> This LINK 02B result is redundant with a result in the previous datablock.

 $\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
0.935 ± 0.017 ± 0.024		<sup>1</sup> DOBBS	07	CLEO See BONVICINI 14
0.97 ± 0.04 ± 0.04	1250 ± 40	<sup>1</sup> HE	05	CLEO See DOBBS 07

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

 $\Gamma(K^+ K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{97}/\Gamma_{43}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.1059 ± 0.0018 OUR FIT</b>				
<b>0.1059 ± 0.0018 OUR AVERAGE</b>				
0.106 ± 0.002 ± 0.003		BONVICINI	14	CLEO All CLEO-c runs
0.117 ± 0.013 ± 0.007	181 ± 20	ABLIKIM	05F	BES $e^+ e^- \approx \psi(3770)$
0.107 ± 0.001 ± 0.002	43k	AUBERT	05S	BABR $e^+ e^- \approx \Upsilon(4S)$
0.093 ± 0.010 <sup>+0.008</sup> <sub>-0.006</sub>		JUN	00	SELX $\Sigma^-$ nucleus, 600 GeV
0.0976 ± 0.0042 ± 0.0046		FRABETTI	95B	E687 $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV

 $\Gamma(\phi\pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+)$   $\Gamma_{98}/\Gamma_{97}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>27.8 ± 0.4 <sup>+0.2</sup> <sub>-0.5</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458 ± 163 evts
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
29.2 ± 3.1 ± 3.0	FRABETTI	95B	E687 Dalitz fit, 915 evts

$$\Gamma(K^+ \bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{99}/\Gamma_{97}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>25.7±0.5<sup>+0.4</sup><sub>-1.2</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458±163 evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
30.1±2.0±2.5	FRABETTI	95B E687	Dalitz fit, 915 evts

$$\Gamma(K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{100}/\Gamma_{97}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>18.8±1.2<sup>+3.3</sup><sub>-3.4</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458±163 evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
37.0±3.5±1.8	FRABETTI	95B E687	Dalitz fit, 915 evts

$$\Gamma(K^+ \bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{101}/\Gamma_{97}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>1.7±0.4<sup>+1.2</sup><sub>-0.7</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458±163 evts

$$\Gamma(K^+ \bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{102}/\Gamma_{97}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>7.0±0.8<sup>+3.5</sup><sub>-2.0</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458±163 evts

$$\Gamma(a_0(1450)^0 \pi^+, a_0^0 \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{103}/\Gamma_{97}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>4.6±0.6<sup>+7.2</sup><sub>-1.8</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458±163 evts

$$\Gamma(\phi(1680) \pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{104}/\Gamma_{97}$$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.51±0.11<sup>+0.37</sup><sub>-0.16</sub></b>	RUBIN	08	CLEO Dalitz fit, 19,458±163 evts

$$\Gamma(K^*(892)^+ K_S^0)/\Gamma(K_S^0 \pi^+) \quad \Gamma_{112}/\Gamma_{41}$$

Unseen decay modes of the  $K^*(892)^+$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.1±0.3±0.4</b>	67	FRABETTI	95	E687 $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(K_S^0 K_S^0 \pi^+)/\Gamma_{\text{total}} \quad \Gamma_{105}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>27.0±0.5±1.2</b>	4897	ABLIKIM	17A BES3	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\phi\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{109}/\Gamma$ Unseen decay modes of the  $\phi$  are included.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.023±0.010</b>	1 BARLAG	92C ACCM	$\pi^-$ Cu 230 GeV

1 BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(\phi\rho^+)/\Gamma(K^-2\pi^+)$  $\Gamma_{110}/\Gamma_{43}$ Unseen decay modes of the  $\phi$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.16</b>	90	DAOUDI	92 CLEO	$e^+e^- \approx 10.5$ GeV

 $\Gamma(K^+K^-\pi^+\pi^0\text{non-}\phi)/\Gamma_{\text{total}}$  $\Gamma_{111}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.015<sup>+0.007</sup><sub>-0.006</sub></b>	1 BARLAG	92C ACCM	$\pi^-$ Cu 230 GeV

1 BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(K^+K^-\pi^+\pi^0\text{non-}\phi)/\Gamma(K^-2\pi^+)$  $\Gamma_{111}/\Gamma_{43}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

 $\Gamma(K^+K_S^0\pi^+\pi^-)/\Gamma(K_S^02\pi^+\pi^-)$  $\Gamma_{106}/\Gamma_{64}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.62±0.39±0.40</b>	$469 \pm 32$	LINK	01C FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(K_S^0K^-2\pi^+)/\Gamma(K_S^02\pi^+\pi^-)$  $\Gamma_{107}/\Gamma_{64}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.68±0.41±0.32</b>	$670 \pm 35$	LINK	01C FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(K^+K^-2\pi^+\pi^-)/\Gamma(K^-3\pi^+\pi^-)$  $\Gamma_{108}/\Gamma_{65}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.040±0.009±0.019</b>	38	LINK	03D FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV

---

 Doubly Cabibbo-suppressed modes 

---

 $\Gamma(K^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{113}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.81±0.27 OUR FIT</b>	Error includes scale factor of 1.4.			
<b>2.52±0.47±0.26</b>	$189 \pm 37$	AUBERT,B	06F BABR	$e^+e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

2.28±0.36±0.17       $148 \pm 23$       DYTMAN      06      CLEO      See MENDEZ 10 $\Gamma(K^+\pi^0)/\Gamma(K^-2\pi^+)$  $\Gamma_{113}/\Gamma_{43}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.01±0.30 OUR FIT</b>	Error includes scale factor of 1.4.			
<b>1.9 ± 0.2 ± 0.1</b>	$343 \pm 37$	MENDEZ	10	CLEO $e^+e^-$ at 3774 MeV

$\Gamma(K^+\eta)/\Gamma(\eta\pi^+)$  $\Gamma_{114}/\Gamma_{91}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.06±0.43±0.14</b>	166 ± 23	WON	11	BELL $e^+e^- \approx \gamma(4S)$

 $\Gamma(K^+\eta)/\Gamma(K^-2\pi^+)$  $\Gamma_{114}/\Gamma_{43}$ Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.15	90	MENDEZ	10	CLEO $e^+e^-$ at 3774 MeV

 $\Gamma(K^+\eta'(958))/\Gamma(\eta'(958)\pi^+)$  $\Gamma_{115}/\Gamma_{94}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.77±0.39±0.10</b>	180 ± 19	WON	11	BELL $e^+e^- \approx \gamma(4S)$

 $\Gamma(K^+\eta'(958))/\Gamma(K^-2\pi^+)$  $\Gamma_{115}/\Gamma_{43}$ Unseen decay modes of the  $\eta'(958)$  are included.

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.20	90	MENDEZ	10	CLEO $e^+e^-$ at 3774 MeV

 $\Gamma(K^+\pi^+\pi^-)/\Gamma(K^-2\pi^+)$  $\Gamma_{116}/\Gamma_{43}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.77±0.22 OUR AVERAGE</b>				
5.69±0.18±0.14	2638 ± 84	KO	09	BELL $e^+e^-$ at $\gamma(4S)$
6.5 ± 0.8 ± 0.4	189 ± 24	LINK	04F	FOCS $\gamma A$ , $\overline{E}_\gamma \approx 180$ GeV
7.7 ± 1.7 ± 0.8	59 ± 13	AITALA	97C	E791 $\pi^- A$ , 500 GeV
7.2 ± 2.3 ± 1.7	21	FRABETTI	95E	E687 $\gamma Be$ , $\overline{E}_\gamma = 220$ GeV

 $\Gamma(K^+\rho^0)/\Gamma(K^+\pi^+\pi^-)$  $\Gamma_{117}/\Gamma_{116}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.39 ±0.09 OUR AVERAGE</b>			
0.3943±0.0787±0.0815	LINK	04F	FOCS Dalitz fit, 189 evts
0.37 ± 0.14 ± 0.07	AITALA	97C	E791 Dalitz fit, 59 evts

 $\Gamma(K^+f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(K^+\pi^+\pi^-)$  $\Gamma_{119}/\Gamma_{116}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0892±0.0333±0.0412</b>	LINK	04F	FOCS Dalitz fit, 189 evts

 $\Gamma(K^*(892)^0\pi^+, K^*(892)^0 \rightarrow K^+\pi^-)/\Gamma(K^+\pi^+\pi^-)$  $\Gamma_{118}/\Gamma_{116}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.47 ±0.08 OUR AVERAGE</b>			
0.5220±0.0684±0.0638	LINK	04F	FOCS Dalitz fit, 189 evts
0.35 ± 0.14 ± 0.01	AITALA	97C	E791 Dalitz fit, 59 evts

$\Gamma(K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow K^+ \pi^-)/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{120}/\Gamma_{116}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.0803 \pm 0.0372 \pm 0.0391$	LINK	04F	FOCS Dalitz fit, 189 evts

 $\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{121}/\Gamma_{116}$ 

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.36 \pm 0.14 \pm 0.07$	<sup>1</sup> AITALA	97C E791	Dalitz fit, 59 evts
<sup>1</sup> LINK 04F, with three times as many events, finds no need for a nonresonant amplitude.			

 $\Gamma(2K^+ K^-)/\Gamma(K^- 2\pi^+)$   $\Gamma_{122}/\Gamma_{43}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$9.49 \pm 2.17 \pm 0.22$	65	<sup>1</sup> LINK	02I	FOCS $\gamma$ nucleus, $\approx 180$ GeV

<sup>1</sup> LINK 02I finds little evidence for  $\phi K^+$  or  $f_0(980) K^+$  submodes.

---

Rare or forbidden modes

---

 $\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{123}/\Gamma$ A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<5.9 \times 10^{-6}$	90	<sup>1</sup> RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$
$<7.4 \times 10^{-6}$	90	HE	05A CLEO	See RUBIN 10
$<5.2 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<1.1 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<6.6 \times 10^{-5}$	90	AITALA	96 E791	$\pi^- N$ 500 GeV
$<2.5 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
$<2.6 \times 10^{-3}$	90	HAAS	88 CLEO	$e^+ e^-$ 10 GeV

<sup>1</sup> This RUBIN 10 limit is for the  $e^+ e^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block. $\Gamma(\pi^+ \phi, \phi \rightarrow e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{124}/\Gamma$ This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ e^+ e^-$  final state.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$(1.7^{+1.4}_{-0.9} \pm 0.1) \times 10^{-6}$	4	<sup>1</sup> RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(2.7^{+3.6}_{-1.8} \pm 0.2) \times 10^{-6}$	2	HE	05A CLEO	See RUBIN 10
--	---	----	----------	--------------

<sup>1</sup> This RUBIN 10 result is consistent with the known  $D^+ \rightarrow \phi \pi^+$  and  $\phi \rightarrow e^+ e^-$  fractions.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_{125}/\Gamma$ 

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-8}$	90	AAIJ	13AF LHCb	$p\bar{p}$ at 7 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<6.5 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$
$<3.9 \times 10^{-6}$	90	<sup>1</sup> ABAZOV	08D D0	$p\bar{p}, E_{\text{cm}} = 1.96 \text{ TeV}$
$<8.8 \times 10^{-6}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_{\gamma} \approx 180 \text{ GeV}$
$<1.5 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<8.9 \times 10^{-5}$	90	FRAEBETTI	97B E687	$\gamma Be, \bar{E}_{\gamma} \approx 220 \text{ GeV}$
$<1.8 \times 10^{-5}$	90	AITALA	96 E791	$\pi^- N$ 500 GeV
$<2.2 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV
$<5.9 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
$<2.9 \times 10^{-3}$	90	HAAS	88 CLEO	$e^+ e^-$ 10 GeV

<sup>1</sup> This ABAZOV 08D limit is for the  $\mu^+ \mu^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block.

 $\Gamma(\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_{126}/\Gamma$ 

This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ \mu^+ \mu^-$  final state.

VALUE	DOCUMENT ID	TECN	COMMENT
$(1.8 \pm 0.5 \pm 0.6) \times 10^{-6}$	<sup>1</sup> ABAZOV	08D D0	$p\bar{p}, E_{\text{cm}} = 1.96 \text{ TeV}$

<sup>1</sup> This ABAZOV 08D value is consistent with the known  $D^+ \rightarrow \phi \pi^+$  and  $\phi \rightarrow \mu^+ \mu^-$  fractions.

 $\Gamma(\rho^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_{127}/\Gamma$ 

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV

 $\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{128}/\Gamma$ 

Both quarks would have to change flavor for this decay to occur.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<3.0 \times 10^{-6}$	90	RUBIN	10 CLEO	$e^+ e^-$ at $\psi(3770)$
$<6.2 \times 10^{-6}$	90	HE	05A CLEO	See RUBIN 10
$<2.0 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<2.0 \times 10^{-4}$	90	FRAEBETTI	97B E687	$\gamma Be, \bar{E}_{\gamma} \approx 220 \text{ GeV}$
$<4.8 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

 $\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_{129}/\Gamma$ 

Both quarks would have to change flavor for this decay to occur.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.2 \times 10^{-6}$	90	LINK	03F	FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV
$<4.4 \times 10^{-5}$	90	AITALA	99G	E791	$\pi^- N$ 500 GeV
$<9.7 \times 10^{-5}$	90	FRABETTI	97B	E687	$\gamma Be, \bar{E}_\gamma \approx 220$ GeV
$<3.2 \times 10^{-4}$	90	KODAMA	95	E653	$\pi^-$ emulsion 600 GeV
$<9.2 \times 10^{-3}$	90	WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

### $\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$

$\Gamma_{130}/\Gamma$

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	LEES	11G	BABR $e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-4}$	90	FRABETTI	97B	E687 $\gamma Be, \bar{E}_\gamma \approx 220$ GeV
$<3.3 \times 10^{-3}$	90	WEIR	90B	MRK2 $e^+ e^-$ 29 GeV

### $\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$

$\Gamma_{131}/\Gamma$

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6 \times 10^{-6}$	90	LEES	11G	BABR $e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-4}$	90	FRABETTI	97B	E687 $\gamma Be, \bar{E}_\gamma \approx 220$ GeV
$<3.3 \times 10^{-3}$	90	WEIR	90B	MRK2 $e^+ e^-$ 29 GeV

### $\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$

$\Gamma_{132}/\Gamma$

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-6}$	90	LEES	11G	BABR $e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-4}$	90	FRABETTI	97B	E687 $\gamma Be, \bar{E}_\gamma \approx 220$ GeV
$<3.4 \times 10^{-3}$	90	WEIR	90B	MRK2 $e^+ e^-$ 29 GeV

### $\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$

$\Gamma_{133}/\Gamma$

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-6}$	90	LEES	11G	BABR $e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-4}$	90	FRABETTI	97B	E687 $\gamma Be, \bar{E}_\gamma \approx 220$ GeV
$<3.4 \times 10^{-3}$	90	WEIR	90B	MRK2 $e^+ e^-$ 29 GeV

### $\Gamma(\pi^- 2e^+)/\Gamma_{\text{total}}$

$\Gamma_{134}/\Gamma$

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-6}$	90	RUBIN	10	CLEO $e^+ e^-$ at $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.9 \times 10^{-6}$	90	LEES	11G	BABR $e^+ e^- \approx \gamma(4S)$
$<3.6 \times 10^{-6}$	90	HE	05A	CLEO See RUBIN 10
$<9.6 \times 10^{-5}$	90	AITALA	99G	E791 $\pi^- N$ 500 GeV
$<1.1 \times 10^{-4}$	90	FRABETTI	97B	E687 $\gamma Be, \bar{E}_\gamma \approx 220$ GeV
$<4.8 \times 10^{-3}$	90	WEIR	90B	MRK2 $e^+ e^-$ 29 GeV



$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{140}/\Gamma$ 

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.9 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.3 \times 10^{-4}$	90	FRABETTI	97B E687	$\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV
$<4.0 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV

 $\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$  $\Gamma_{141}/\Gamma$ 

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-4}$	90	KODAMA	95	E653 $\pi^-$ emulsion 600 GeV

 $D^\pm CP$ -VIOLATING DECAY-RATE ASYMMETRIES

This is the difference between  $D^+$  and  $D^-$  partial widths for the decay to state  $f$ , divided by the sum of the widths:

$$A_{CP}(f) = [\Gamma(D^+ \rightarrow f) - \Gamma(D^- \rightarrow \bar{f})]/[\Gamma(D^+ \rightarrow f) + \Gamma(D^- \rightarrow \bar{f})].$$

 $A_{CP}(\mu^\pm \nu)$  in  $D^+ \rightarrow \mu^+ \nu_\mu, D^- \rightarrow \mu^- \bar{\nu}_\mu$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$+8 \pm 8$	EISENSTEIN	08	CLEO $e^+ e^-$ at $\psi(3770)$

 $A_{CP}(K_L^0 e^\pm \nu)$  in  $D^+ \rightarrow K_L^0 e^+ \nu_e, D^- \rightarrow K_L^0 e^- \bar{\nu}_e$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-0.59 \pm 0.60 \pm 1.48$	ABLIKIM	15AF BES3	$e^+ e^-$ 3773 MeV

 $A_{CP}(K_S^0 \pi^\pm)$  in  $D^\pm \rightarrow K_S^0 \pi^\pm$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.41 \pm 0.09</math> OUR AVERAGE</b>				
-1.1	$\pm 0.6$	$\pm 0.2$	BONVICINI	14 CLEO All CLEO-c runs
$-0.363 \pm 0.094 \pm 0.067$	1738k	<sup>1</sup> KO	12A BELL	$e^+ e^- \approx \gamma(nS)$
$-0.44 \pm 0.13 \pm 0.10$	807k	DEL-AMO-SA..11H	BABR	$e^+ e^- \approx \gamma(4S)$
-1.6	$\pm 1.5$	$\pm 0.9$	LINK	02B FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
-0.71	$\pm 0.19$	$\pm 0.20$	KO	10 BELL See KO 12A
-1.3	$\pm 0.7$	$\pm 0.3$	MENDEZ	10 CLEO See BONVICINI 14
-0.6	$\pm 1.0$	$\pm 0.3$	DOBBS	07 CLEO See MENDEZ 10

<sup>1</sup> KO 12A finds that after subtracting the contribution due to  $K^0 - \bar{K}^0$  mixing, the  $CP$  asymmetry due to the change of charm is  $(-0.024 \pm 0.094 \pm 0.067)\%$ , consistent with zero.

<sup>2</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

$A_{CP}(K^\mp 2\pi^\pm)$  in  $D^+ \rightarrow K^- 2\pi^+$ ,  $D^- \rightarrow K^+ 2\pi^-$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.18±0.16 OUR AVERAGE</b>				
-0.16±0.15±0.09	2.3M	ABAZOV	14L D0	$p\bar{p}$ , $\sqrt{s} = 1.96$ TeV
-0.3 ± 0.2 ± 0.4		BONVICINI	14 CLEO	All CLEO-c runs
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.1 ± 0.4 ± 0.9	231k	MENDEZ	10 CLEO	See BONVICINI 14
-0.5 ± 0.4 ± 0.9		DOBBS	07 CLEO	See MENDEZ 10

 $A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0)$  in  $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$ ,  $D^- \rightarrow K^+ \pi^- \pi^- \pi^0$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-0.3±0.6±0.4</b>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.0±0.9±0.9	DOBBS	07 CLEO	See BONVICINI 14

 $A_{CP}(K_S^0 \pi^\pm \pi^0)$  in  $D^+ \rightarrow K_S^0 \pi^+ \pi^0$ ,  $D^- \rightarrow K_S^0 \pi^- \pi^0$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-0.1±0.7±0.2</b>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.3±0.9±0.3	DOBBS	07 CLEO	See BONVICINI 14

 $A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-)$  in  $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$ ,  $D^- \rightarrow K_S^0 \pi^- \pi^- \pi^+$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.0±1.2±0.3</b>			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.1±1.1±0.6	DOBBS	07 CLEO	See BONVICINI 14

 $A_{CP}(\pi^\pm \pi^0)$  in  $D^\pm \rightarrow \pi^\pm \pi^0$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4 ±1.2 OUR AVERAGE</b>				
2.31±1.24±0.23	108k	BABU	18 BELL	At/near $\gamma(4S)$ , $\gamma(5S)$
2.9 ± 2.9 ± 0.3	2.6k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

 $A_{CP}(\pi^\pm \eta)$  in  $D^\pm \rightarrow \pi^\pm \eta$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.0 ±1.5 OUR AVERAGE</b>				
Error includes scale factor of 1.4.				
+1.74±1.13±0.19		WON	11 BELL	$e^+ e^- \approx \gamma(4S)$
-2.0 ± 2.3 ± 0.3	2.9k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

 $A_{CP}(\pi^\pm \eta'(958))$  in  $D^\pm \rightarrow \pi^\pm \eta'(958)$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.6 ±0.7 OUR AVERAGE</b>				
-0.61±0.72±0.54	63k	AAIJ	17AF LHCb	$p\bar{p}$ at 7, 8 TeV
-0.12±1.12±0.17		WON	11 BELL	$e^+ e^- \approx \gamma(4S)$
-4.0 ± 3.4 ± 0.3	1.0k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV

**$A_{CP}(\bar{K}^0/K^0 K^\pm)$  in  $D^+ \rightarrow \bar{K}^0 K^+$ ,  $D^- \rightarrow K^0 K^-$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.11 \pm 0.17</math> OUR AVERAGE</b>				
0.03 $\pm 0.17 \pm 0.14$	1.0M	<sup>1</sup> AAIJ	14BD LHCb	$p\ p$ at 7, 8 TeV
0.08 $\pm 0.28 \pm 0.14$	277k	KO	13 BELL	$e^+ e^-$ at $\Upsilon(4S)$
0.46 $\pm 0.36 \pm 0.25$	159k	LEES	13E BABR	$e^+ e^-$ at $\Upsilon(4S)$

<sup>1</sup> AAIJ 14BD reports its result as  $A_{CP}(D^\pm \rightarrow K_S^0 \pi^\pm)$  with  $CP$ -violation effects in the  $K^0 - \bar{K}^0$  system subtracted. It also measures  $A_{CP}(D^\pm \rightarrow \bar{K}^0/K^0 K^\pm) + A_{CP}(D_s^\pm \rightarrow \bar{K}^0/K^0 \pi^\pm) = (0.41 \pm 0.49 \pm 0.26)\%$ .

 **$A_{CP}(K_S^0 K^\pm)$  in  $D^\pm \rightarrow K_S^0 K^\pm$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.11 \pm 0.25</math> OUR AVERAGE</b>				
-0.25 $\pm 0.28 \pm 0.14$	277k	KO	13 BELL	$e^+ e^-$ at $\Upsilon(nS)$
0.13 $\pm 0.36 \pm 0.25$	159k	LEES	13E BABR	$e^+ e^-$ at $\Upsilon(4S)$
-0.2 $\pm 1.5 \pm 0.9$	5.2k	MENDEZ	10 CLEO	$e^+ e^-$ at 3774 MeV
7.1 $\pm 6.1 \pm 1.2$	949	<sup>1</sup> LINK	02B FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.16 $\pm 0.58 \pm 0.25$	KO	10 BELL	$e^+ e^- \approx \Upsilon(4S)$
6.9 $\pm 6.0 \pm 1.5$	949	<sup>2</sup> LINK	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

<sup>1</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 K^+)/N(D^+ \rightarrow K_S^0 \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

<sup>2</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 K^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

 **$A_{CP}(K^+ K^- \pi^\pm)$  in  $D^\pm \rightarrow K^+ K^- \pi^\pm$** 

See also AAIJ 11G for a search for  $CP$  asymmetry in the  $D^\pm \rightarrow K^+ K^- \pi^\pm$  Dalitz plots using 370k decays and four different binning schemes. No evidence for  $CP$  asymmetry was found.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.37 \pm 0.29</math> OUR AVERAGE</b>				
0.37 $\pm 0.30 \pm 0.15$	224k	<sup>1</sup> LEES	13F BABR	$e^+ e^-$ at $\Upsilon(4S)$
-0.03 $\pm 0.84 \pm 0.29$		RUBIN	08 CLEO	$e^+ e^-$ at 3774 MeV
1.4 $\pm 1.0 \pm 0.8$	43k	<sup>2</sup> AUBERT	05S BABR	$e^+ e^-$ at $\Upsilon(4S)$
0.6 $\pm 1.1 \pm 0.5$	14k	<sup>3</sup> LINK	00B FOCS	
-1.4 $\pm 2.9$		<sup>3</sup> AITALA	97B E791	$-0.062 < A_{CP} < +0.034$ (90% CL)
-3.1 $\pm 6.8$		<sup>3</sup> FRABETTI	94I E687	$-0.14 < A_{CP} < +0.081$ (90% CL)

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.1 $\pm 0.9 \pm 0.4$	<sup>4</sup> BONVICINI	14 CLEO	See RUBIN 08
-0.1 $\pm 1.5 \pm 0.8$	DOBBS	07 CLEO	See BONVICINI 14 and RUBIN 08

<sup>1</sup> This is the integrated  $CP$  asymmetry. LEES 13F also searches for  $CP$  asymmetries in four regions of the Dalitz plots (two of which are listed below); in comparisons of binned  $D^+$  and  $D^-$  Dalitz plots; in parametrized fits to those plots, including 2-body submodes; and in comparisons of Legendre-polynomial distributions for the  $K^+ K^-$  and  $K^- \pi^+$  systems.

- <sup>2</sup>AUBERT 05S measures  $N(D^+ \rightarrow K^+ K^- \pi^+)/N(D_s^+ \rightarrow K^+ K^- \pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .
- <sup>3</sup>FRABETTI 94I, AITALA 98C, and LINK 00B measure  $N(D^+ \rightarrow K^- K^+ \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .
- <sup>4</sup>RUBIN 08 performs a dedicated analysis of this decay mode on the same dataset, with slightly better precision. We therefore take it that BONVICINI 14 does not supersede RUBIN 08's  $A_{CP}$  result.

### $A_{CP}(K^\pm K^{*0})$ in $D^+ \rightarrow K^+ \bar{K}^{*0}$ , $D^- \rightarrow K^- K^{*0}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.3 ± 0.4 OUR AVERAGE</b>				
- 0.3 ± 0.4 ± 0.2	73k	<sup>1</sup> LEES	13F BABR	$e^+ e^-$ at $\Upsilon(4S)$
- 0.4 ± 2.0 ± 0.6		RUBIN	08 CLEO	Fit-fraction asymmetry
+ 0.9 ± 1.7 ± 0.7	11k	<sup>2</sup> AUBERT	05S BABR	$e^+ e^-$ at $\Upsilon(4S)$
- 1.0 ± 5.0		<sup>3</sup> AITALA	97B E791	$-0.092 < A_{CP} < +0.072$ (90% CL)
-12 ± 13		<sup>3</sup> FRABETTI	94I E687	$-0.33 < A_{CP} < +0.094$ (90% CL)

<sup>1</sup>This LEES 13F result is for the  $K^\mp \pi^\pm$  mass-squared between 0.4 and 1.0 GeV<sup>2</sup>, and does not actually separate out the  $K^*$ .

<sup>2</sup>AUBERT 05S measures  $N(D^+ \rightarrow K^+ \bar{K}^{*0})/N(D_s^+ \rightarrow K^+ K^- \pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup>FRABETTI 94I and AITALA 97B measure  $N(D^+ \rightarrow K^+ \bar{K}^*(892)^0)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(\phi \pi^\pm)$ in $D^\pm \rightarrow \phi \pi^\pm$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.09 ± 0.19 OUR AVERAGE</b>				
		Error includes scale factor of 1.2.		
-0.04 ± 0.14 ± 0.14	1.58M	AAIJ	13W LHCb	$p\ p$ at 7 TeV
-0.3 ± 0.3 ± 0.5	97k	<sup>1</sup> LEES	13F BABR	$e^+ e^-$ at $\Upsilon(4S)$
+0.51 ± 0.28 ± 0.05	237k	STARIC	12 BELL	Mainly at $\Upsilon(4S)$
-1.8 ± 1.6 ± 0.2		RUBIN	08 CLEO	Fit-fraction asymmetry
-0.4				
+0.2 ± 1.5 ± 0.6	10k	<sup>2</sup> AUBERT	05S BABR	$e^+ e^-$ at $\Upsilon(4S)$
-2.8 ± 3.6		<sup>3</sup> AITALA	97B E791	$-0.087 < A_{CP} < +0.031$ (90% CL)
+6.6 ± 8.6		<sup>3</sup> FRABETTI	94I E687	$-0.075 < A_{CP} < +0.21$ (90% CL)

<sup>1</sup>This LEES 13F result is for the  $K^+ K^-$  mass-squared less than 1.3 GeV<sup>2</sup> and the  $K^\mp \pi^\pm$  mass-squared above 1.0 GeV<sup>2</sup>, and does not actually separate out the  $\phi$ .

<sup>2</sup>AUBERT 05S measures  $N(D^+ \rightarrow \phi \pi^+)/N(D_s^+ \rightarrow K^+ K^- \pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup>FRABETTI 94I and AITALA 97B measure  $N(D^+ \rightarrow \phi \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(K^\pm K_0^*(1430)^0)$ in $D^+ \rightarrow K^+ \bar{K}_0^*(1430)^0$ , $D^- \rightarrow K^- K_0^*(1430)^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+8 ± 6 ± 4	RUBIN	08 CLEO	Fit-fraction asymmetry

**$A_{CP}(K^\pm K_2^*(1430)^0)$  in  $D^+ \rightarrow K^+ \bar{K}_2^*(1430)^0$ ,  $D^- \rightarrow K^- K_2^*(1430)^0$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$+43 \pm 19^{+5}_{-18}$	RUBIN	08	CLEO Fit-fraction asymmetry

 **$A_{CP}(K^\pm K_0^*(700))$  in  $D^+ \rightarrow K^+ \bar{K}_0^*(700)$ ,  $D^- \rightarrow K^- K_0^*(700)$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-12 \pm 11^{+14}_{-6}$	RUBIN	08	CLEO Fit-fraction asymmetry

 **$A_{CP}(a_0(1450)^0 \pi^\pm)$  in  $D^\pm \rightarrow a_0(1450)^0 \pi^\pm$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-19 \pm 12^{+8}_{-11}$	RUBIN	08	CLEO Fit-fraction asymmetry

 **$A_{CP}(\phi(1680) \pi^\pm)$  in  $D^\pm \rightarrow \phi(1680) \pi^\pm$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-9 \pm 22 \pm 14$	RUBIN	08	CLEO Fit-fraction asymmetry

 **$A_{CP}(\pi^+ \pi^- \pi^\pm)$  in  $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$** 

See also AAIJ 14C for a search for  $CP$  violation in  $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$  Dalitz plots using model-independent binned and unbinned methods. No evidence was found.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-1.7 \pm 4.2$	<sup>1</sup> AITALA	97B E791	$-0.086 < A_{CP} < +0.052$ (90% CL)

<sup>1</sup> AITALA 97B measure  $N(D^+ \rightarrow \pi^+ \pi^- \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

 **$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-)$  in  $D^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$-4.2 \pm 6.4 \pm 2.2$	$523 \pm 32$	LINK	05E FOCS	$\gamma A$ , $\bar{E}_\gamma \approx 180$ GeV

 **$A_{CP}(K^\pm \pi^0)$  in  $D^\pm \rightarrow K^\pm \pi^0$** 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$-3.5 \pm 10.7 \pm 0.9$	$343 \pm 37$	MENDEZ	10	CLEO $e^+ e^-$ at 3774 MeV

 **$D^\pm \chi^2$  TESTS OF CP-VIOLATION (CPV)**

We list model-independent searches for local  $CP$  violation in phase-space distributions of multi-body decays.

Most of these searches divide phase space (Dalitz plot for 3-body decays, five-dimensional equivalent for 4-body decays) into bins, and perform a  $\chi^2$  test comparing normalised yields  $N_i$ ,  $\bar{N}_i$  in  $CP$ -conjugate bin pairs  $i$ :  $\chi^2 = \sum_i (N_i - \alpha \bar{N}_i)/\sigma(N_i - \alpha \bar{N}_i)$ . The factor  $\alpha = (\sum_i N_i)/(\sum_i \bar{N}_i)$  removes the dependence on phase-space-integrated rate asymmetries. The result is used to obtain the probability (p-value) to obtain the measured  $\chi^2$  or larger under the assumption of  $CP$  conservation [AUBERT 08A0, BEDIAGA 09]. Alternative methods obtain p-values from other test variables based on unbinned analyses [WILLIAMS 11, AAIJ 14C]. Results can be combined using Fisher's method [MOSTELLER 48].

**Local CPV in  $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$** 

<i>p</i> -value (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>78.1</b>	3.1M	<sup>1</sup> AAIJ	14C LHCb	$\chi^2$

<sup>1</sup> AAIJ 14C uses binned and unbinned methods, and finds slightly better sensitivity with the former. We took the first value in the table of results for the binned method.

**Local CPV in  $D^\pm \rightarrow K^+ K^- \pi^\pm$** 

<i>p</i> -value (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>31 OUR EVALUATION</b>				
72	224k	LEES	13F BABR	$\chi^2$

<i>p</i> -value (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.7	370k	<sup>1</sup> AAIJ	11G LHCb	$\chi^2$

<sup>1</sup> AAIJ 11G publishes results for several binning schemes. We picked the first value in their table of results.

**CP VIOLATING ASYMMETRIES OF *P*-ODD (*T*-ODD) MOMENTS** **$A_{T\text{Viol}}(K_S^0 K^\pm \pi^+ \pi^-)$  in  $D^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$** 

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$  is a parity-odd correlation of the  $K^+$ ,  $\pi^+$ , and  $\pi^-$  momenta for the  $D^+$ .  $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$  is the corresponding quantity for the  $D^-$ . Then

$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$ , and

$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]$ , and

$A_{T\text{Viol}} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ .  $C_T$  and  $\bar{C}_T$  are commonly referred to as *T*-odd moments, because they are odd under *T* reversal. However, the *T*-conjugate process  $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D^\pm$  is not accessible, while the *P*-conjugate process is.


VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-12.0 \pm 10.0 \pm 4.6</math></b>	$21.2 \pm 0.4$ k	LEES	11E BABR	$e^+ e^- \approx \gamma(4S)$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
23 $\pm 62$ $\pm 22$	$523 \pm 32$	LINK	05E FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

 **$D^+ \rightarrow (\bar{K}^0/\pi^0/\eta/\omega/\rho^0/\bar{K}^{*0})\ell^+\nu_\ell$  FORM FACTORS** **$f_+(0)|V_{cs}|$  in  $D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.719 <math>\pm 0.011</math> OUR AVERAGE</b>			Error includes scale factor of 1.6. See the ideogram below.
0.7053 $\pm 0.0040 \pm 0.0112$	ABLIKIM	17S BES3	$K_S^0 e^+ \nu_e$ 2-parameter fit
0.737 $\pm 0.006 \pm 0.009$	<sup>1</sup> ABLIKIM	15AF BES3	$K_L e^+ \nu_e$ 3-parameter fit
0.707 $\pm 0.010 \pm 0.009$	<sup>2</sup> BESSON	09 CLEO	$K_S^0 e^+ \nu_e$ 3-parameter fit

<sup>1</sup> ABLIKIM 15AF finds  $0.728 \pm 0.006 \pm 0.011$  for a 2-parameter fit.

<sup>2</sup> BESSON 09 finds  $0.716 \pm 0.007 \pm 0.009$  for a 2-parameter fit.



$$f_+(0)|V_{cs}| \text{ in } D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$$

$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-2.13<math>\pm</math>0.14 OUR AVERAGE</b>				
-2.18 $\pm$ 0.14 $\pm$ 0.05		ABLIKIM	17S BES3	$K_S^0 e^+ \nu_e$ 2-parameter fit
-2.23 $\pm$ 0.42 $\pm$ 0.53	40k	<sup>1</sup> ABLIKIM	15AF BES3	$K_L e^+ \nu_e$ 3-parameter fit
-1.66 $\pm$ 0.44 $\pm$ 0.10		<sup>2</sup> BESSON	09 CLEO	$K_S^0 e^+ \nu_e$ 3-parameter fit

<sup>1</sup> ABLIKIM 15AF finds  $r_1 = -1.91 \pm 0.33 \pm 0.28$  for a 2-parameter fit.

<sup>2</sup> BESSON 09 finds  $r_1 = -2.10 \pm 0.25 \pm 0.08$  for 2-parameter fit.

$r_2 \equiv a_2/a_0$  in  $D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-3<math>\pm</math>12 OUR AVERAGE</b> Error includes scale factor of 1.5.				
+11 $\pm$ 9 $\pm$ 9	40k	ABLIKIM	15AF BES3	$K_L e^+ \nu_e$ 3-parameter fit
-14 $\pm$ 11 $\pm$ 1		BESSON	09 CLEO	$K_S^0 e^+ \nu_e$ 3-parameter fit

$f_+(0)|V_{cd}|$  in  $D^+ \rightarrow \pi^0 \ell^+ \nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.1407<math>\pm</math>0.0025 OUR AVERAGE</b>			
0.1400 $\pm$ 0.0026 $\pm$ 0.0007	ABLIKIM	17S BES3	$\pi^0 e^+ \nu_e$ 2-parameter fit
0.146 $\pm$ 0.007 $\pm$ 0.002	BESSON	09 CLEO	$\pi^0 e^+ \nu_e$ 3-parameter fit

$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \pi^0 \ell^+ \nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-2.00<math>\pm</math>0.13 OUR AVERAGE</b>			
-2.01 $\pm$ 0.13 $\pm$ 0.02	ABLIKIM	17S BES3	$\pi^0 e^+ \nu_e$ 2-parameter fit
-1.37 $\pm$ 0.88 $\pm$ 0.24	BESSON	09 CLEO	$\pi^0 e^+ \nu_e$ 3-parameter fit

$r_2 \equiv a_2/a_0$  in  $D^+ \rightarrow \pi^0 \ell^+ \nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-4±5±1</b>	BESSON 09	CLEO	$\pi^0 e^+ \nu_e$ 3-parameter fit

$f_+(0)|V_{cd}|$  in  $D^+ \rightarrow \eta e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.086±0.006±0.001</b>	YELTON 11	CLEO	$z$ expansion

$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \eta e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-1.83±2.23±0.28</b>	YELTON 11	CLEO	$z$ expansion

$r_v \equiv V(0)/A_1(0)$  in  $D^+ \rightarrow \omega e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.24±0.09±0.06</b>	ABLIKIM 15W	BES3	$292 \text{ fb}^{-1}$ , 3773 MeV

$r_2 \equiv A_2(0)/A_1(0)$  in  $D^+ \rightarrow \omega e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.06±0.15±0.05</b>	ABLIKIM 15W	BES3	$292 \text{ fb}^{-1}$ , 3773 MeV

$r_v \equiv V(0)/A_1(0)$  in  $D^+, D^0 \rightarrow \rho e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.48±0.15±0.05</b>	1 DOBBS 13	CLEO	$e^+ e^-$ at $\psi(3770)$

<sup>1</sup> Uses both  $D^+$  and  $D^0$  events. Using PDG 10 values of  $V_{cd}$  and lifetimes, DOBBS 13 gets  $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$ ,  $A_2(0) = 0.47 \pm 0.06 \pm 0.04$ , and  $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$ .

$r_2 \equiv A_2(0)/A_1(0)$  in  $D^+, D^0 \rightarrow \rho e^+ \nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.83±0.11±0.04</b>	1 DOBBS 13	CLEO	$e^+ e^-$ at $\psi(3770)$

<sup>1</sup> Uses both  $D^+$  and  $D^0$  events. Using PDG 10 values of  $V_{cd}$  and lifetimes, DOBBS 13 gets  $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$ ,  $A_2(0) = 0.47 \pm 0.06 \pm 0.04$ , and  $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$ .

$r_v \equiv V(0)/A_1(0)$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$


See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.49 ± 0.05 OUR AVERAGE</b>		Error includes scale factor of 2.1.		See the ideogram below.
1.411±0.058±0.007	16.2k	ABLIKIM 16F	BES3	$\bar{K}^*(892)^0 e^+ \nu_e$
1.463±0.017±0.031		1 DEL-AMO-SA..11I	BABR	
1.504±0.057±0.039	15k	2 LINK 02L	FOCS	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.45 ± 0.23 ± 0.07	763	ADAMOVICH 99	BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.90 ± 0.11 ± 0.09	3000	3AITALA 98B	E791	$\bar{K}^*(892)^0 e^+ \nu_e$
1.84 ± 0.11 ± 0.09	3034	AITALA 98F	E791	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.74 ± 0.27 ± 0.28	874	FRABETTI 93E	E687	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
2.00 $^{+0.34}_{-0.32}$ ± 0.16	305	KODAMA 92	E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
2.0 ± 0.6 ± 0.3	183	ANJOS 90E	E691	$\bar{K}^*(892)^0 e^+ \nu_e$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.6 ± 0.3 183 ANJOS 90E E691  $\bar{K}^*(892)^0 e^+ \nu_e$

- <sup>1</sup> DEL-AMO-SANCHEZ 11I finds the pole mass  $m_A = (2.63 \pm 0.10 \pm 0.13)$  GeV ( $m_V$  is fixed at 2 GeV).  
<sup>2</sup> LINK 02L includes the effects of interference with an *S*-wave background. This much improves the goodness of fit, but does not much shift the values of the form factors.  
<sup>3</sup> This is slightly different from the AITALA 98B value: see ref. [5] in AITALA 98F.



### $r_2 \equiv A_2(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.802±0.021 OUR AVERAGE</b>				
0.788±0.042±0.008	16.2k	ABLIKIM	16F	$\bar{K}^*(892)^0 e^+ \nu_e$
0.801±0.020±0.020		<sup>1</sup> DEL-AMO-SA... 11I	BABR	
0.875±0.049±0.064	15k	<sup>2</sup> LINK	02L	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.00 ± 0.15 ± 0.03	763	ADAMOVICH	99	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.71 ± 0.08 ± 0.09	3000	AITALA	98B	$\bar{K}^*(892)^0 e^+ \nu_e$
0.75 ± 0.08 ± 0.09	3034	AITALA	98F	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.78 ± 0.18 ± 0.10	874	FRABETTI	93E	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.82 ± 0.22 ± 0.11	305	KODAMA	92	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0 ± 0.5 ± 0.2	183	ANJOS	90E	$\bar{K}^*(892)^0 e^+ \nu_e$
-----------------	-----	-------	-----	------------------------------

<sup>1</sup> DEL-AMO-SANCHEZ 11I finds the pole mass  $m_A = (2.63 \pm 0.10 \pm 0.13)$  GeV ( $m_V$  is fixed at 2 GeV).

<sup>2</sup> LINK 02L includes the effects of interference with an *S*-wave background. This much improves the goodness of fit, but does not much shift the values of the form factors.

**$r_3 \equiv A_3(0)/A_1(0)$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$** See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.04±0.33±0.29</b>	3034	AITALA	98F E791	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$

 **$\Gamma_L/\Gamma_T$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$** See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.13±0.08 OUR AVERAGE</b>				
1.09±0.10±0.02	763	ADAMOVICH 99	BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.20±0.13±0.13	874	FRABETTI 93E	E687	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.18±0.18±0.08	305	KODAMA 92	E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.8 $^{+0.6}_{-0.4}$ ±0.3	183	ANJOS 90E	E691	$\bar{K}^*(892)^0 e^+ \nu_e$

 **$\Gamma_+/\Gamma_-$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$** See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.22±0.06 OUR AVERAGE</b> Error includes scale factor of 1.6.				
0.28±0.05±0.02	763	ADAMOVICH 99	BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.16±0.05±0.02	305	KODAMA 92	E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.15 $^{+0.07}_{-0.05}$ ±0.03	183	ANJOS 90E	E691	$\bar{K}^*(892)^0 e^+ \nu_e$

 **$D^\pm$  REFERENCES**

BABU	18	PR D97 011101	V. Babu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AF	PL B771 21	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17A	PL B765 231	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AD	PR D96 092002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17M	PR D95 071102	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17S	PR D96 012002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16D	PRL 116 082001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16F	PR D94 032001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16G	EPJ C76 369	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16V	CP C40 113001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15AF	PR D92 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15W	PR D92 071101	M. Ablikim <i>et al.</i>	(BES III Collab.)
AAIJ	14BD	JHEP 1410 025	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14C	PL B728 585	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	14L	PR D90 111102	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	14E	PR D89 052001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14F	PR D89 051104	M. Ablikim <i>et al.</i>	(BES III Collab.)
BONVICINI	14	PR D89 072002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AAIJ	13AF	PL B724 203	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13W	JHEP 1306 112	R. Aaij <i>et al.</i>	(LHCb Collab.)
DOBBS	13	PRL 110 131802	S. Dobbs <i>et al.</i>	(CLEO Collab.)
KO	13	JHEP 1302 098	B.R. Ko <i>et al.</i>	(BELLE Collab.)
LEES	13E	PR D87 052012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
KO	12A	PRL 109 119903 (errat.)	B.R. Ko <i>et al.</i>	(BELLE Collab.)
Also		PRL 109 021601	B.R. Ko <i>et al.</i>	(BELLE Collab.)
STARIC	12	PRL 108 071801	M. Staric <i>et al.</i>	(BELLE Collab.)
AAIJ	11G	PR D84 112008	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA... 11H	PR D83 071103	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	
DEL-AMO-SA... 11I	PR D83 072001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	

LEES	11E	PR D84 031103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11G	PR D84 072006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WILLIAMS	11	PR D84 054015	M. Williams	(LOIC)
WON	11	PRL 107 221801	E. Won <i>et al.</i>	(BELLE Collab.)
YELTON	11	PR D84 032001	J. Yelton <i>et al.</i>	(CLEO Collab.)
ANASHIN	10A	PL B686 84	V.V. Anashin <i>et al.</i>	(VEPP-4M KEDR Collab.)
ASNER	10	PR D81 052007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	10	PR D81 112001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
KO	10	PRL 104 181602	B.R. Ko <i>et al.</i>	(BELLÉ Collab.)
MENDEZ	10	PR D81 052013	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)
RUBIN	10	PR D82 092007	P. Rubin <i>et al.</i>	(CLEO Collab.)
BEDIAGA	09	PR D80 096006	I. Bediaga <i>et al.</i>	(CBPF, NDAM)
BESSON	09	PR D80 032005	D. Besson <i>et al.</i>	(CLEO Collab.)
Also		PR D79 052010	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KO	09	PRL 102 221802	B.R. Ko <i>et al.</i>	(BELLÉ Collab.)
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
MITCHELL	09B	PRL 102 081801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
WON	09	PR D80 111101	E. Won <i>et al.</i>	(BELLÉ Collab.)
ABAZOV	08D	PRL 100 101801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	08L	PL B665 16	M. Ablikim <i>et al.</i>	(BES Collab.)
ARTUSO	08	PR D77 092003	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	08AO	PR D78 051102	B. Aubert <i>et al.</i>	(BABAR Collab.)
BONVICINI	08	PR D77 091106	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
DOBBS	08	PR D77 112005	S. Dobbs <i>et al.</i>	(CLEO Collab.)
Also		PRL 100 251802	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
EISENSTEIN	08	PR D78 052003	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)
HE	08	PRL 100 091801	Q. He <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
RUBIN	08	PR D78 072003	P. Rubin <i>et al.</i>	(CLEO Collab.)
ABLIKIM	07	PL B644 20	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07G	PL B658 1	M. Ablikim <i>et al.</i>	(BES Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	06O	EPJ C47 31	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06P	EPJ C47 39	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06U	PL B643 246	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06A	PRL 97 251801	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AUBERT,B	06F	PR D74 011107	B. Aubert <i>et al.</i>	(BABAR Collab.)
DYTMAN	06	PR D74 071102	S.A. Dytman <i>et al.</i>	(CLEO Collab.)
HUANG	06B	PR D74 112005	G.S. Huang <i>et al.</i>	(CLEO Collab.)
LINK	06B	PL B637 32	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
RUBIN	06	PRL 96 081802	P. Rubin <i>et al.</i>	(CLEO Collab.)
RUBIN	06A	PR D73 112005	P. Rubin <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05A	PL B608 24	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05D	PL B610 183	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05F	PL B622 6	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05P	PL B625 196	M. Ablikim <i>et al.</i>	(BES Collab.)
ARTUSO	05A	PRL 95 251801	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	05S	PR D71 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)
Also		PRL 96 199903 (errat.)	Q. He <i>et al.</i>	(CLEO Collab.)
HE	05A	PRL 95 221802	Q. He <i>et al.</i>	(CLEO Collab.)
HUANG	05B	PR L 95 181801	G.S. Huang <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAK..05		PL B626 24	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)
LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	04C	PL B597 39	M. Ablikim <i>et al.</i>	(BEPC BES Collab.)
ARMS	04	PR D69 071102	K. Arms <i>et al.</i>	(CLEO Collab.)
BONVICINI	04A	PR D70 112004	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04E	PL B598 33	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04F	PL B601 10	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03D	PL B561 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03F	PL B572 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)

BRANDENB...	02	PRL 89 222001	G. Brandenburg <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAK..	02	PL B549 48	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)
LINK	02B	PRL 88 041602	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
Also		PRL 88 159903 (errat.)	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02E	PL B535 43	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02F	PL B537 192	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02I	PL B541 227	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02J	PL B541 243	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02L	PL B544 89	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
LINK	01C	PRL 87 162001	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABREU	000	EPJ C12 209	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ASTIER	00D	PL B486 35	P. Astier <i>et al.</i>	(CERN NOMAD Collab.)
JUN	00	PRL 84 1857	S.Y. Jun <i>et al.</i>	(FNAL SELEX Collab.)
LINK	00B	PL B491 232	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
Also		PL B495 443 (errat.)	J.M. Link <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99K	EPJ C8 573	G. Abbiendi <i>et al.</i>	(CERN BEATRICE Collab.)
ADAMOVICH	99	EPJ C6 35	M. Adamovich <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(CLEO Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98B	PRL 80 1393	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98C	PL B421 405	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	98F	PL B440 435	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BAI	98B	PL B429 188	J.Z. Bai <i>et al.</i>	(BEPC BES Collab.)
AITALA	97	PL B397 325	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	97B	PL B403 377	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	97C	PL B404 187	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BISHAI	97	PRL 78 3261	M. Bishai <i>et al.</i>	(CLEO Collab.)
FRAZETTI	97	PL B391 235	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	97B	PL B398 239	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	97C	PL B401 131	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AITALA	96	PRL 76 364	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
FRAZETTI	95	PL B346 199	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	95B	PL B351 591	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	95E	PL B359 403	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	94I	ZPHY C64 375	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	94	PRL 72 2328	R. Balest <i>et al.</i>	(CLEO Collab.)
FRAZETTI	94D	PL B323 459	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	94G	PL B331 217	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAZETTI	94I	PR D50 2953	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AKERIB	93	PRL 71 3070	D.S. Akerib <i>et al.</i>	(CLEO Collab.)
ANJOS	93	PR D48 56	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
FRAZETTI	93E	PL B307 262	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	92F	PL B278 202	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	92C	PR D46 1941	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
COFFMAN	92B	PR D45 2196	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
DAOUDI	92	PR D45 3965	M. Daoudi <i>et al.</i>	(CLEO Collab.)
KODAMA	92	PL B274 246	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
KODAMA	92C	PL B286 187	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ADAMOVICH	91	PL B268 142	M.I. Adamovich <i>et al.</i>	(WA82 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALVAREZ	91B	ZPHY C50 11	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
AMMAR	91	PR D44 3383	R. Ammar <i>et al.</i>	(CLEO Collab.)
BAI	91	PRL 66 1011	Z. Bai <i>et al.</i>	(Mark III Collab.)
COFFMAN	91	PL B263 135	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FRAZETTI	91	PL B263 584	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90C	PR D41 2705	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	90D	PR D42 2414	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	90E	PR L 65 2630	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
WEIR	90B	PR D41 1384	A.J. Weir <i>et al.</i>	(Mark II Collab.)
ANJOS	89	PRL 62 125	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	89B	PRL 62 722	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)
ALBRECHT	88I	PL B210 267	H. Albrecht <i>et al.</i>	(ARGUS Collab.)

HAAS	88	PRL 60 1614	P. Haas <i>et al.</i>	(CLEO Collab.)
ONG	88	PRL 60 2587	R.A. Ong <i>et al.</i>	(Mark II Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	M.I. Adamovich <i>et al.</i>	(Photon Emulsion Collab.)
ADLER	87	PL B196 107	J. Adler <i>et al.</i>	(Mark III Collab.)
BARTEL	87	ZPHY C33 339	W. Bartel <i>et al.</i>	(JADE Collab.)
BALTRUSAIT...	86E	PRL 56 2140	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	85B	PRL 54 1976	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	85E	PRL 55 150	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BARTEL	85J	PL 163B 277	W. Bartel <i>et al.</i>	(JADE Collab.)
ADAMOVICH	84	PL 140B 119	M.I. Adamovich <i>et al.</i>	(CERN WA58 Collab.)
ALTHOFF	84G	ZPHY C22 219	M. Althoff <i>et al.</i>	(TASSO Collab.)
DERRICK	84	PRL 53 1971	M. Derrick <i>et al.</i>	(HRS Collab.)
SCHINDLER	81	PR D24 78	R.H. Schindler <i>et al.</i>	(Mark II Collab.)
TRILLING	81	PRPL 75 57	G.H. Trilling	(LBL, UCB) J
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
PERUZZI	77	PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)
PICCOLO	77	PL 70B 260	M. Piccolo <i>et al.</i>	(Mark I Collab.)
PERUZZI	76	PRL 37 569	I. Peruzzi <i>et al.</i>	(Mark I Collab.)
MOSTELLER	48	Am.Stat. 3 No.5 30	R.A. Fisher, F. Mosteller	

## OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
ROSNER	95	CNPP 21 369	J. Rosner	(CHIC)