

**$K_2^*(1980)$** 

$I(J^P) = \frac{1}{2}(2^+)$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

 **$K_2^*(1980)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1974±26 OUR AVERAGE</b>					
2073±94 <sup>+245</sup> <sub>-240</sub>	4289	<sup>1</sup> AAIJ	17C	LHCb	$B^+ \rightarrow J/\psi \phi K^+$
1973± 8± 25		ASTON	87	LASS 0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2020±20		TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1978±40	241	BIRD	89	LASS -	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 $\sigma$ .					

 **$K_2^*(1980)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>376± 70 OUR AVERAGE</b>					
678±311 <sup>+1153</sup> <sub>-559</sub>	4289	<sup>2</sup> AAIJ	17C	LHCb	$B^+ \rightarrow J/\psi \phi K^+$
373± 33± 60		ASTON	87	LASS 0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
180± 70		TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
398± 47	241	BIRD	89	LASS -	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
<sup>2</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 $\sigma$ .					

 **$K_2^*(1980)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 K^*(892)\pi$	possibly seen
$\Gamma_2 K\rho$	possibly seen
$\Gamma_3 Kf_2(1270)$	possibly seen
$\Gamma_4 K\phi$	seen

 **$K_2^*(1980)$  BRANCHING RATIOS**

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	$\Gamma_1/\Gamma$
possibly seen	GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
<b><math>\Gamma(K\rho)/\Gamma_{\text{total}}</math></b>			
possibly seen	GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	$\Gamma_2/\Gamma_1$				
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
<b>1.49±0.24±0.09</b>	ASTON 87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
$\Gamma(K f_2(1270))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$				
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>possibly seen</b>	TIKHOMIROV 03	SPEC	40.0	$\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
$\Gamma(K\phi)/\Gamma_{\text{total}}$	$\Gamma_4/\Gamma$				
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>seen</b>	4289	<sup>3</sup> AAIJ	17C	LHCb $B^+ \rightarrow J/\psi \phi K^+$	
<sup>3</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 $\sigma$ .					

## **$K_2^*(1980)$ REFERENCES**

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)