

$a_2(1700)$ $I^G(J^{PC}) = 1^-(2^{++})$

OMITTED FROM SUMMARY TABLE

 $a_2(1700)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1732 ± 9 OUR AVERAGE		Error includes scale factor of 1.2.			
1726 $\pm 12 \pm 25$		¹ ABLIKIM	17K	BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$
1737 $\pm 5 \pm 7$		ABE	04	BELL	$10.6 e^+e^- \rightarrow e^+e^- K^+K^-$
1698 ± 44		² AMSLER	02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta$
1660 ± 40		ABELE	99B	CBAR	$1.94 \bar{p}p \rightarrow \pi^0\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1675 ± 25		ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$
1722 $\pm 9 \pm 15$	18k	³ SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
1702 ± 7	80k	⁴ UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
1721 $\pm 13 \pm 44$	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega\pi^-\pi^0 p$
1767 ± 14	221	⁵ ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
~ 1775		⁶ GRYGOREV	99	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1752 $\pm 21 \pm 4$		ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

¹ From an amplitude analysis using an isobar model.² T-matrix pole.³ From analysis of L3 data at 183–209 GeV.⁴ Statistical error only.⁵ Spin 2 dominant, isospin not determined, could also be $J=1$.⁶ Possibly two $J^P = 2^+$ resonances with isospins 0 and 1. **$a_2(1700)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
193 ± 27 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
190 $\pm 18 \pm 30$		⁷ ABLIKIM	17K	BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$
151 $\pm 22 \pm 24$		ABE	04	BELL	$10.6 e^+e^- \rightarrow e^+e^- K^+K^-$
265 ± 55		⁸ AMSLER	02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta$
280 ± 70		ABELE	99B	CBAR	$1.94 \bar{p}p \rightarrow \pi^0\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
270^{+50}_{-20}		ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$
336 $\pm 20 \pm 20$	18k	⁹ SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
417 ± 19	80k	¹⁰ UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
279 $\pm 49 \pm 66$	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega\pi^-\pi^0 p$
187 ± 60	221	¹¹ ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
150 $\pm 110 \pm 34$		ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$


⁷ From an amplitude analysis using an isobar model.

⁸ T-matrix pole.

⁹ From analysis of L3 data at 183–209 GeV.

¹⁰ Statistical error only.

¹¹ Spin 2 dominant, isospin not determined, could also be $J=1$.



a₂(1700) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta\pi$	seen
$\Gamma_2 \gamma\gamma$	
$\Gamma_3 \rho\pi$	
$\Gamma_4 f_2(1270)\pi$	
$\Gamma_5 K\bar{K}$	seen
$\Gamma_6 \omega\pi^-\pi^0$	seen
$\Gamma_7 \omega\rho$	seen

a₂(1700) PARTIAL WIDTHS

$\Gamma(\eta\pi)$	Γ_1
VALUE (MeV)	EVTS
<hr/>	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
9.5±2.0	870 ¹² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(\gamma\gamma)$ Γ_2

<u>VALUE</u> (keV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.30 ± 0.05	870	12 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

 $\Gamma(K\bar{K})$ Γ_5

<u>VALUE</u> (MeV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
5.0 ± 3.0	870	12 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
12 From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.				

 $a_2(1700) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$(\Gamma_3 + \Gamma_4)\Gamma_2/\Gamma$			
<u>VALUE</u> (keV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.29 \pm 0.04 \pm 0.02$		ACCIARRI	97T	L3 $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.37^{+0.12}_{-0.08} \pm 0.10$	18k	13 SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_2/\Gamma$

<u>VALUE</u> (eV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
20.6 \pm 4.2 \pm 4.6	14 ABE	04	BELL	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$
49 \pm 11 \pm 13	15 ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{ee} = 91, 183\text{--}209 \text{ GeV}$

 13 From analysis of L3 data at 183–209 GeV. 14 Assuming spin 2. 15 Spin 2 dominant, isospin not determined, could also be $J=1$. $a_2(1700) \text{ BRANCHING RATIOS}$ $\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$ Γ_3/Γ_4

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$3.4 \pm 0.4 \pm 0.1$	18k	16 SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

 16 From analysis of L3 data at 183–209 GeV.

$a_2(1700)$ REFERENCES

ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>	
ACCIARRI	97T	Translated from YAF 62 513.	M. Acciarri <i>et al.</i>	(L3 Collab.)
PL B413 147				