


$\chi_{b1}(2P)$
 $I^G(J^{PC}) = 0^+(1^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b1}(2P)$ MASSVALUE (MeV)
 $10255.46 \pm 0.22 \pm 0.50$ OUR EVALUATION From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV
 $m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$ VALUE (MeV)
 $23.5 \pm 0.7 \pm 0.7$ DOCUMENT ID TECN COMMENT
¹ HEINTZ 92 CSB2 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$
¹ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.
 γ ENERGY IN $\Upsilon(3S)$ DECAYVALUE (MeV)

<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
99.26 ± 0.22 OUR EVALUATION	Treating systematic errors as correlated		
99.53 ± 0.23 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
99.15 $\pm 0.07 \pm 0.25$	ARTUSO 05	CLEO	$\Upsilon(3S) \rightarrow \gamma X$
99 ± 1	CRAWFORD 92B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
100.1 ± 0.4	² HEINTZ 92	CSB2	$e^+ e^- \rightarrow \gamma X$
100.2 ± 0.5	³ HEINTZ 92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
99.5 $\pm 0.1 \pm 0.5$	MORRISON 91	CLE2	$e^+ e^- \rightarrow \gamma X$

WEIGHTED AVERAGE
 99.53 ± 0.23 (Error scaled by 1.3)


²A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

³A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

$\chi_{b1}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \omega \gamma(1S)$	(1.63 $^{+0.40}_{-0.34}$) %
$\Gamma_2 \gamma \gamma(2S)$	(18.1 ± 1.9) %
$\Gamma_3 \gamma \gamma(1S)$	(9.9 ± 1.0) %
$\Gamma_4 \pi\pi\chi_{b1}(1P)$	(9.1 ± 1.3) $\times 10^{-3}$
$\Gamma_5 D^0 X$	(8.8 ± 1.7) %
$\Gamma_6 \pi^+\pi^- K^+ K^- \pi^0$	(3.1 ± 1.0) $\times 10^{-4}$
$\Gamma_7 2\pi^+\pi^- K^- K_S^0$	(1.1 ± 0.5) $\times 10^{-4}$
$\Gamma_8 2\pi^+\pi^- K^- K_S^0 2\pi^0$	(7.7 ± 3.2) $\times 10^{-4}$
$\Gamma_9 2\pi^+ 2\pi^- 2\pi^0$	(5.9 ± 2.0) $\times 10^{-4}$
$\Gamma_{10} 2\pi^+ 2\pi^- K^+ K^-$	(10 ± 4) $\times 10^{-5}$
$\Gamma_{11} 2\pi^+ 2\pi^- K^+ K^- \pi^0$	(5.5 ± 1.8) $\times 10^{-4}$
$\Gamma_{12} 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(10 ± 4) $\times 10^{-4}$
$\Gamma_{13} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	(6.7 ± 2.6) $\times 10^{-4}$
$\Gamma_{14} 3\pi^+ 3\pi^-$	(1.2 ± 0.4) $\times 10^{-4}$
$\Gamma_{15} 3\pi^+ 3\pi^- 2\pi^0$	(1.2 ± 0.4) $\times 10^{-3}$
$\Gamma_{16} 3\pi^+ 3\pi^- K^+ K^-$	(2.0 ± 0.8) $\times 10^{-4}$
$\Gamma_{17} 3\pi^+ 3\pi^- K^+ K^- \pi^0$	(6.1 ± 2.2) $\times 10^{-4}$
$\Gamma_{18} 4\pi^+ 4\pi^-$	(1.7 ± 0.6) $\times 10^{-4}$
$\Gamma_{19} 4\pi^+ 4\pi^- 2\pi^0$	(1.9 ± 0.7) $\times 10^{-3}$

$\chi_{b1}(2P)$ BRANCHING RATIOS

$\Gamma(\omega \gamma(1S))/\Gamma_{\text{total}}$	Γ_1/Γ
$1.63^{+0.35+0.16}_{-0.31-0.15}$	$32.6^{+6.9}_{-6.1}$ ⁴ CRONIN-HEN..04 CLE3 $\gamma(3S) \rightarrow \gamma\omega \gamma(1S)$

⁴ Using $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.3 \pm 0.6)\%$ and $B(\gamma(1S) \rightarrow \ell^+\ell^-) = 2 B(\gamma(1S) \rightarrow \mu^+\mu^-) = 2 (2.48 \pm 0.06)\%$.

$\Gamma(\gamma \gamma(2S))/\Gamma_{\text{total}}$	Γ_2/Γ
0.181 ± 0.019 OUR AVERAGE	
$0.211 \pm 0.017 \pm 0.019$	^{5,6,7} LEES 14M BABR $\gamma(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$0.190 \pm 0.018 \pm 0.017$	^{4.3k} ⁸ LEES 11J BABR $\gamma(3S) \rightarrow X\gamma$
$0.206 \pm 0.035 \pm 0.019$	^{5,9} CRAWFORD 92B CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
$0.132 \pm 0.018 \pm 0.012$	^{5,10} HEINTZ 92 CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

⁵ Assuming $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

⁶ LEES 14M quotes $\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$ $= (2.66 \pm 0.22)\%$ combining the results from $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ samples with and without photon conversions.

⁷ LEES 14M reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (2.66 \pm 0.22) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \ell^+ \ell^-) = (10.23 \pm 1.20 \pm 1.26) 10^{-4}$.

¹⁰ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.099±0.010 OUR AVERAGE					
0.107±0.006±0.010	11,12,13	LEES	14M	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
0.098±0.005±0.009	15k	14	BABR	$\Upsilon(3S) \rightarrow X \gamma$	
0.103±0.023±0.009		11,15	CRAWFORD	92B	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
0.075±0.010±0.007		11,16	HEINTZ	92	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

¹¹ Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

¹² LEES 14M quotes $\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$ $= (13.48 \pm 0.72) \times 10^{-3}$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and without converted photons.

¹³ LEES 14M reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (13.48 \pm 0.72) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁴ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁵ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (6.47 \pm 1.12 \pm 0.82) 10^{-4}$.

¹⁶ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91.

$\Gamma(\pi \pi \chi_{b1}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
9.1±1.3 OUR AVERAGE					
9.2±1.1±0.8	31k	17	LEES	11C	$BABR \quad e^+ e^- \rightarrow \pi^+ \pi^- X$
8.6±2.3±2.1		18	CAWLFIELD	06	$CLE3 \quad \Upsilon(3S) \rightarrow 2(\gamma \pi \ell)$

¹⁷ LEES 11C measures $B(\gamma(3S) \rightarrow \chi_{b1}(2P)X \times B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$. We derive the value assuming $B(\gamma(3S) \rightarrow \chi_{b1}(2P)X) = B(\gamma(3S) \rightarrow \chi_{b1}(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$.

¹⁸ CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming I-spin conservation, no D -wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$	Γ_5/Γ			
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.8±1.5±0.8	2243	19 BRIERE	08 CLEO	$\gamma(3S) \rightarrow \gamma D^0 X$

¹⁹ For $p_{D^0} > 2.5$ GeV/c.

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.1±1.0±0.3	30	20 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$

²⁰ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (39 \pm 8 \pm 9) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$	Γ_7/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.1±0.5±0.1	10	21 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

²¹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (14 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$	Γ_8/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.7±3.1±0.7	15	22 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

²² ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (97 \pm 30 \pm 26) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$	Γ_9/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.9±2.0±0.5	36	23 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

²³ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (74 \pm 16 \pm 19) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.4±0.1	12	24 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$
24 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.5±1.7±0.5	38	25 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
25 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (69 \pm 13 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.6±3.5±0.9	27	26 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
26 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (121 \pm 29 \pm 33) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±2.5±0.6	17	27 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
27 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (85 \pm 23 \pm 22) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.4±0.1	18	28 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
28 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (15 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	44	29 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
29 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (150 \pm 30 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.0±0.7±0.2	16	30 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
30 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (25 \pm 7 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±2.1±0.6	25	31 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
31 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (77 \pm 17 \pm 21) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7±0.6±0.2	16	32 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
32 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (22 \pm 6 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19±7±2	41	33 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
33 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))] = (241 \pm 47 \pm 72) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\chi_{b1}(2P)$ Cross-Particle Branching Ratios

$$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}} \\ \Gamma_3/\Gamma \times \Gamma_{21}^{(3S)}/\Gamma^{(3S)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±0.3±0.6	15k	LEES	11J BABR	$\Gamma(3S) \rightarrow X\gamma$


$$B(\chi_{b1}(2P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Gamma(1S) \rightarrow \ell^+ \ell^-)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.9. See the ideogram below.
$3.52^{+0.28+0.17}_{-0.27-0.18}$		34 LEES	14M BABR	$\Gamma(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$3.24^{+0.56+0.41}_{-0.28-0.15}$	58	35 CRAWFORD	92B CLE2	$\Gamma(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
$2.34^{+0.28+0.15}_{-0.27-0.18}$		36 HEINTZ	92 CSB2	$\Gamma(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

³⁴ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.

³⁵ CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.

³⁶ Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$.



$$B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$$

(units 10^{-4})

$$\frac{\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S))/\Gamma_{\text{total}}}{\Gamma_2/\Gamma} \times \frac{\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))/\Gamma_{\text{total}}}{\Gamma_{21}/\Gamma_{21}} = \frac{\Gamma_2/\Gamma}{\Gamma_{21}/\Gamma_{21}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 0.1 \pm 0.2$	4.3k	LEES	11J	$\Upsilon(3S) \rightarrow X\gamma$

$$B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+\ell^-)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.

$4.95^{+0.75+1.01}_{-0.70-0.24}$ 37 LEES 14M BABR $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$


$5.12 \pm 0.60 \pm 0.63$ 111 38 CRAWFORD 92B CLE2 $\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

$3.30 \pm 0.33 \pm 0.20$ 39 HEINTZ 92 CSB2 $\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

³⁷ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.

³⁸ CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.

³⁹ Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$.



$$B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$$

(units 10^{-4})

$$B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P) \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b1}(2P) X)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.16±0.07±0.12	31k	LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$

$$B(\chi_{b2}(2P) \rightarrow pX + \bar{p}X) / B(\chi_{b1}(2P) \rightarrow pX + \bar{p}X)$$

VALUE	DOCUMENT ID	TECN	COMMENT
1.109±0.007±0.040	BRIERE	07	CLEO $\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

$$B(\chi_{b0}(2P) \rightarrow pX + \bar{p}X) / B(\chi_{b1}(2P) \rightarrow pX + \bar{p}X)$$

VALUE	DOCUMENT ID	TECN	COMMENT
1.082±0.025±0.060	BRIERE	07	CLEO $\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

$\chi_{b1}(2P)$ REFERENCES

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawfield <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HENNESSY	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)