

$\omega(1420)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\omega(1420)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
(1400–1450) OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1418 ± 30 ± 10	824	¹ AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
1470 ± 50	13.1k	² AULCHENKO 15A	SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1382 ± 23 ± 70		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1350 ± 20 ± 20		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1400 ± 50 ± 130	1.2M	³ ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1450 ± 10		⁴ HENNER 02	RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1373 ± 70	177	⁵ AKHMETSHIN 00D	CMD2	1.2–1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
1370 ± 25	5095	ANISOVICH 00H	SPEC	0.0 $p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
1400 ⁺¹⁰⁰ ₋₂₀₀		⁶ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
~1400		⁷ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
~1460		⁸ ACHASOV 98H	RVUE	$e^+e^- \rightarrow K^+K^-$
1440 ± 70		⁹ CLEGG 94	RVUE	
1419 ± 31	315	¹⁰ ANTONELLI 92	DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi$

¹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

² From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁵ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

⁶ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

⁷ Using the data from ANTONELLI 92.

⁸ Using the data from IVANOV 81 and BISELLO 88B.

⁹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹⁰ From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed (+, -, +) phases.

 $\omega(1420)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
(180–250) OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
104 ± 35 ± 10	824	¹ AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
880 ± 170	13.1k	² AULCHENKO 15A	SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

$130 \pm 50 \pm 100$	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
$450 \pm 70 \pm 70$	AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
$870^{+500}_{-300} \pm 450$	1.2M	3 ACHASOV	$0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
199 ± 15		4 HENNER	$02 RVUE 1.2-2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
188 ± 45	177	5 AKHMETSHIN	$00D CMD2 1.2-1.38 e^+ e^- \rightarrow \omega \pi^+ \pi^-$
360^{+100}_{-60}	5095	ANISOVICH	$00H SPEC 0.0 p\bar{p} \rightarrow \omega \pi^0 \pi^0 \pi^0$
240 ± 70		6 CLEGG	$94 RVUE$
174 ± 59	315	7 ANTONELLI	$92 DM2 1.34-2.4 e^+ e^- \rightarrow \rho \pi$

¹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

² From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁵ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho \pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

⁶ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁷ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed $(+, -, +)$ phases.

$\omega(1420)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho \pi$	dominant
$\Gamma_2 \omega \pi \pi$	seen
$\Gamma_3 \omega \eta$	
$\Gamma_4 b_1(1235) \pi$	seen
$\Gamma_5 e^+ e^-$	seen
$\Gamma_6 \pi^0 \gamma$	

$$\omega(1420) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$$

$\Gamma(\rho \pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma \times \Gamma_5/\Gamma$			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.73 ± 0.08	13.1k	1 AULCHENKO	15A SND	$1.05-1.80 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.82 $\pm 0.05 \pm 0.06$		AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
0.65 $\pm 0.13 \pm 0.21$	1.2M	2,3 ACHASOV	03D RVUE	$0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.625 ± 0.160		4,5 CLEGG	94 RVUE	
0.466 ± 0.178		6,7 ANTONELLI	92 DM2	$1.34-2.4 e^+ e^- \rightarrow \rho \pi$

- ¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.
² Calculated by us from the cross section at the peak.
³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
⁵ From the partial and leptonic width given by the authors.
⁶ From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed $(+,-,+)$ phases.
⁷ From the product of the leptonic width and partial branching ratio given by the authors.

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
19.7 ± 5.7	AUBERT 07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$	
1.9 ± 1.9	¹ AKHMETSHIN 00D CMD2	$1.2-2.4 e^+e^- \rightarrow \omega\pi^+\pi^-$	

¹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$5.0 \pm 2.6 \pm 0.3$	824	¹ AKHMETSHIN 17A CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$	
$1.6^{+0.9}_{-0.7}$	898	² ACHASOV 16B SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$	

¹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1420) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1420) \rightarrow e^+e^-) = 5.3 \pm 1.6$ eV.

² From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass and the width of $\omega(1420)$ are fixed to the 2014 edition (PDG 14) of this review.

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$2.03^{+0.70}_{-0.75}$	¹ AKHMETSHIN 05 CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$	

¹ Using 1420 MeV and 220 MeV for the $\omega(1420)$ mass and width.

$\omega(1420)$ BRANCHING RATIOS

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.301 ± 0.029 possibly seen	¹ HENNER 02 RVUE AKHMETSHIN 00D CMD2	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$ $e^+e^- \rightarrow \omega\pi^+\pi^-$	

$\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$ Γ_2/Γ_4

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.60 ± 0.16	5095	ANISOVICH	00H SPEC	$0.0 p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.699 ± 0.029	1 HENNER	02 RVUE	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 6.6	1.2M	2,3 ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
23 ± 1	1 HENNER	02 RVUE	1.2–2.0 e^+e^-	$\rightarrow \rho\pi, \omega\pi\pi$

¹ Assuming that the $\omega(1420)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.² Calculated by us from the cross section at the peak.³ Assuming that the $\omega(1420)$ decays into $\rho\pi$ only. **$\omega(1420)$ REFERENCES**

AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
		Translated from ZETF 148 34.		
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)