

$\psi(4160)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\psi(4160)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4191 ± 5 OUR AVERAGE			
4191 + 9 - 8	AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 ± 6.5	¹ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4193 ± 7	² MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4151 ± 4	³ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4155 ± 5	⁴ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C DASP	$e^+ e^-$

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

 $\psi(4160)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70 ±10 OUR AVERAGE			
65 +22 -16	AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8 ± 12.3	¹ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
79 ± 14	² MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
107 ± 10	³ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
107 ± 16	⁴ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
78 ± 20	BRANDELIK	78C DASP	$e^+ e^-$

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4160)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$	
$\Gamma_2 \mu^+ \mu^-$	seen	
$\Gamma_3 D\bar{D}$	seen	
$\Gamma_4 D^0\bar{D}^0$	seen	
$\Gamma_5 D^+ D^-$	seen	
$\Gamma_6 D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_7 D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_8 D^*(2010)^+ D^- + \text{c.c.}$	seen	
$\Gamma_9 D^*\bar{D}^*$	seen	
$\Gamma_{10} D^*(2007)^0\bar{D}^*(2007)^0$	seen	
$\Gamma_{11} D^*(2010)^+ D^*(2010)^-$	seen	
$\Gamma_{12} D^0 D^- \pi^+ + \text{c.c. (excl.)}$ $D^*(2007)^0\bar{D}^0 + \text{c.c.},$ $D^*(2010)^+ D^- + \text{c.c.})$	not seen	
$\Gamma_{13} D\bar{D}^* \pi + \text{c.c. (excl. } D^*\bar{D}^*)$	seen	
$\Gamma_{14} D^0 D^{*-} \pi^+ + \text{c.c. (excl.)}$ $D^*(2010)^+ D^*(2010)^-$	not seen	
$\Gamma_{15} D_s^+ D_s^-$	not seen	
$\Gamma_{16} D_s^{*+} D_s^- + \text{c.c.}$	seen	
$\Gamma_{17} J/\psi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{18} J/\psi \pi^0 \pi^0$	$< 3 \times 10^{-3}$	90%
$\Gamma_{19} J/\psi K^+ K^-$	$< 2 \times 10^{-3}$	90%
$\Gamma_{20} J/\psi \eta$	$< 8 \times 10^{-3}$	90%
$\Gamma_{21} J/\psi \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{22} J/\psi \eta'$	$< 5 \times 10^{-3}$	90%
$\Gamma_{23} J/\psi \pi^+ \pi^- \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{24} \psi(2S) \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{25} \chi_{c1} \gamma$	$< 5 \times 10^{-3}$	90%
$\Gamma_{26} \chi_{c2} \gamma$	$< 1.3 \%$	90%
$\Gamma_{27} \chi_{c1} \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{28} \chi_{c2} \pi^+ \pi^- \pi^0$	$< 8 \times 10^{-3}$	90%
$\Gamma_{29} h_c(1P) \pi^+ \pi^-$	$< 5 \times 10^{-3}$	90%
$\Gamma_{30} h_c(1P) \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{31} h_c(1P) \eta$	$< 2 \times 10^{-3}$	90%
$\Gamma_{32} h_c(1P) \pi^0$	$< 4 \times 10^{-4}$	90%

Γ_{33}	$\phi\pi^+\pi^-$	< 2	$\times 10^{-3}$	90%
Γ_{34}	$\gamma\chi_{c1}(3872) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 6.8	$\times 10^{-5}$	90%
Γ_{35}	$\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 1.36	$\times 10^{-4}$	90%
Γ_{36}	$\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 1.18	$\times 10^{-4}$	90%
Γ_{37}	$\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 1.47	$\times 10^{-4}$	90%
Γ_{38}	$\gamma\chi_{c1}(3872) \rightarrow \gamma\gamma J/\psi$	< 1.05	$\times 10^{-4}$	90%
Γ_{39}	$\gamma X(3915) \rightarrow \gamma\gamma J/\psi$	< 1.26	$\times 10^{-4}$	90%
Γ_{40}	$\gamma X(3930) \rightarrow \gamma\gamma J/\psi$	< 8.8	$\times 10^{-5}$	90%
Γ_{41}	$\gamma X(3940) \rightarrow \gamma\gamma J/\psi$	< 1.79	$\times 10^{-4}$	90%
Γ_{42}	K^+K^-			

$\psi(4160)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	Γ_1			
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
0.48±0.22	¹ ABLIKIM	08D	BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.4 to 1.1	² MO	10	RVUE	$e^+e^- \rightarrow$ hadrons
0.83 ± 0.08	³ SETH	05A	RVUE	$e^+e^- \rightarrow$ hadrons
0.84 ± 0.13	⁴ SETH	05A	RVUE	$e^+e^- \rightarrow$ hadrons
0.77 ± 0.23	BRANDELIK	78C	DASP	e^+e^-

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{25}\Gamma_1/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	¹ HAN	15	BELL 10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

<6.1	90	¹ HAN	15	BELL 10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$
--------	----	------------------	----	-------------------------------------------------

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$
<u>VALUE</u> (units 10^{-8})	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$2.8 \pm 0.9 \pm 0.9$	¹ WANG 13B BELL $e^+ e^- \rightarrow J/\psi\eta\gamma$
$12.8 \pm 1.7 \pm 2.0$	² WANG 13B BELL $e^+ e^- \rightarrow J/\psi\eta\gamma$
¹ Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.	
² Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.	

$\psi(4160)$ BRANCHING RATIOS

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	
1 AAIJ	13BC LHCb $B^+ \rightarrow K^+ \mu^+ \mu^-$
1 AAIJ 13BC report $B(B^+ \rightarrow K^+ \psi(4160)) B(\psi(4160) \rightarrow \mu^+ \mu^-) = (3.5^{+0.9}_{-0.8}) \times 10^{-9}$.	

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$	Γ_3/Γ_9
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.02 \pm 0.03 \pm 0.02$	AUBERT 09M BABR $e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	CRONIN-HEN..09 CLEO $e^+ e^- \rightarrow D^0\bar{D}^0$
seen	PAKHLOVA 08 BELL $e^+ e^- \rightarrow D^0\bar{D}^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
not seen	AUBERT 09M BABR $e^+ e^- \rightarrow D^0\bar{D}^0\gamma$

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	Γ_5/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	CRONIN-HEN..09 CLEO $e^+ e^- \rightarrow D^+ D^-$
seen	PAKHLOVA 08 BELL $e^+ e^- \rightarrow D^+ D^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
not seen	AUBERT 09M BABR $e^+ e^- \rightarrow D^+ D^- \gamma$

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	AUBERT 09M BABR $e^+ e^- \rightarrow D^{*0}\bar{D}^0\gamma$
seen	CRONIN-HEN..09 CLEO $e^+ e^- \rightarrow D^{*0}\bar{D}^0$

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	1 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$	
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$	

¹ Supersedes PAKHLOVA 07. $\Gamma(D^* \bar{D} + \text{c.c.})/\Gamma(D^* \bar{D}^*)$ Γ_6/Γ_9

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.34 ± 0.14 ± 0.05	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$	

 $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$	

 $\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	1 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	

¹ Supersedes PAKHLOVA 07. $\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.}))/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$	

 $\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$	

 $\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$	

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$	

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$

 Γ_{16}/Γ $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{17}/Γ $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{18}/Γ $\Gamma(J/\psi K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{19}/Γ $\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{20}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	¹ ABLIKIM	15L	BES3	$e^+ e^- \rightarrow J/\psi \eta$
seen	WANG	13B	BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

¹ An enhancement around 4.2 GeV is observed.

 $\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{21}/Γ $\Gamma(J/\psi \eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{22}/Γ $\Gamma(J/\psi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{23}/Γ $\Gamma(\psi(2S) \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{24}/Γ $\Gamma(\chi_{c1} \gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 Γ_{25}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 COAN 06 CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$					Γ_{26}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<13	90	COAN	06	CLEO	$4.12-4.2 \text{ e}^+ \text{e}^- \rightarrow \text{hadrons}$

$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{27}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2	90	COAN	06	CLEO	$4.12-4.2 \text{ e}^+ \text{e}^- \rightarrow \text{hadrons}$

$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{28}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<8	90	COAN	06	CLEO	$4.12-4.2 \text{ e}^+ \text{e}^- \rightarrow \text{hadrons}$

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{29}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<5	90	¹ PEDLAR	11	CLEO	$e^+ e^- \rightarrow h_c(1P)\pi^+\pi^-$

¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+ e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_{30}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2	90	¹ PEDLAR	11	CLEO	$e^+ e^- \rightarrow h_c(1P)\pi^0\pi^0$

¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+ e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$					Γ_{31}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	¹ PEDLAR	11	CLEO	$e^+ e^- \rightarrow h_c(1P)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 41 ² ABLIKIM 17R BES3 $e^+ e^- \rightarrow h_c(1P)\eta$

¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+ e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

² An enhancement around 4.2 GeV is observed.

$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$					Γ_{32}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.4	90	¹ PEDLAR	11	CLEO	$e^+ e^- \rightarrow h_c(1P)\pi^0$

¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+ e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{33}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2	90	COAN	06	CLEO	$4.12-4.2 \text{ e}^+ \text{e}^- \rightarrow \text{hadrons}$

$\Gamma(\gamma\chi_{c1}(3872) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<0.68 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.36 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.18 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.47 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma\chi_{c1}(3872) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.05 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3915) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.26 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3930) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<0.88 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3940) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.79 \times 10^{-4}$	90	¹ XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$<2 \times 10^{-5}$ 90 ¹DRUZHININ 15 RVUE $e^+e^- \rightarrow \psi(3770)$

¹ DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+e^- \rightarrow K^+K^-$ and $e^+e^- \rightarrow K_S^0K_L^0$.

$\psi(4160)$ REFERENCES

ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
XIAO	13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
