

## 25. Neutrinos in Cosmology

Written August 2017 by J. Lesgourgues (RWTH Aachen U.) and L. Verde (U. of Barcelona & ICREA).

### 25.1. Standard neutrino cosmology

Neutrino properties leave detectable imprints on cosmological observations that can then be used to constrain neutrino properties. This is a great example of the remarkable interconnection and interplay between nuclear physics, particle physics, astrophysics and cosmology (for general reviews see *e.g.*, [1,2,3,4]) . Present cosmological data are already providing constraints on neutrino properties not only complementary but also competitive with terrestrial experiments; for instance, upper bounds on the total neutrino mass have shrunk by a factor of about 10 in the past 15 years. Forthcoming cosmological data may soon provide key information, not obtainable in other ways like *e.g.*, a measurement of the absolute neutrino mass scale. This new section is motivated by this exciting prospect.

A relic neutrino background pervading the Universe (the Cosmic Neutrino background,  $C\nu B$ ) is a generic prediction of the standard hot Big Bang model (see Big Bang Nucleosynthesis – Chap. 23 of this *Review*). While it has not yet been detected directly, it has been indirectly confirmed by the accurate agreement of predictions and observations of: *a*) the primordial abundance of light elements (see Big Bang Nucleosynthesis – Chap. 23) of this *Review*; *b*) the power spectrum of Cosmic Microwave Background (CMB) anisotropies (see Cosmic Microwave Background – Chap. 28 of this *Review*); and *c*) the large scale clustering of cosmological structures. Within the hot Big Bang model such good agreement would fail dramatically without a  $C\nu B$  with properties matching closely those predicted by the standard neutrino decoupling process (*i.e.*, involving only weak interactions).

We will illustrate below that cosmology is sensitive to the following neutrino properties: their density, related to the number of active (*i.e.*, left-handed, see Neutrino Mass, Mixing, and Oscillations - Chap. 14 of this *Review*) neutrino species, and their masses. At first order, cosmology is sensitive to the total neutrino mass, but is blind to the mixing angles and CP violation phase as discussed in Neutrino Mass, Mixing, and Oscillations (Chap. 14 of this *Review*). This makes cosmological constraints nicely complementary to measurements from terrestrial neutrino experiments.

The minimal cosmological model,  $\Lambda$ CDM, currently providing a good fit to most cosmological data sets (up to moderate tensions discussed in The Cosmological Parameters Chap. 24 of this *Review*), assumes that the only massless or light (sub-keV) relic particles since the Big Bang Nucleosynthesis (BBN) epoch are photons and active neutrinos. Extended models with light sterile neutrinos, light thermal axions or other light relics –sometimes referred to as “dark radiation”– would produce effects similar to, and potentially degenerate with, those of active neutrinos. Thus neutrino bounds are often discussed together with limits on such scenarios. In case of anomalies in cosmological data, it might not be obvious to discriminate between interpretation in terms of active neutrinos with non-standard decoupling, additional production mechanisms, non-standard interactions, etc., or in terms of some additional light particles. At the moment, such extensions are, however, not required by the cosmological data at any significant level.

## 2 25. Neutrinos in cosmology

Hence neutrino density and mass bounds can be derived under the assumption of no additional massless or light relic particles, and the neutrino density measured in that way provides a test of standard (*i.e.*, involving only weak interactions) neutrino decoupling.

In that model, the three active neutrino types thermalize in the early Universe, with a negligible leptonic asymmetry. Then they can be viewed as three propagating mass eigenstates sharing the same temperature and identical Fermi-Dirac distributions, thus with no visible effects of flavour oscillations. Neutrinos decouple gradually from the thermal plasma at temperatures  $T \sim 2 \text{ MeV}$ . In the instantaneous neutrino decoupling limit, *i.e.*, assuming that neutrinos were fully decoupled at the time when electron-positrons annihilate and release entropy in the thermal bath, the neutrino-to-photon density ratio between the time of electron-positron annihilation and the non-relativistic transition of neutrinos would be given by

$$\frac{\rho_\nu}{\rho_\gamma} = \frac{7}{8} N_{\text{eff}} \left( \frac{4}{11} \right)^{4/3}, \quad (25.1)$$

with  $N_{\text{eff}} = 3$ , and the last factor comes from the fourth power of the temperature ratio  $T_\nu/T_\gamma = (4/11)^{1/3}$  (see Big Bang Cosmology – Chap. 21 in this *Review*). In the above formula,  $N_{\text{eff}}$  is called the effective number of neutrino species because it can be viewed as a convenient parametrisation of the relativistic energy density of the Universe beyond that of photons, in units of one neutrino in the instantaneous decoupling limit. Precise simulations of neutrino decoupling and electron-positron annihilation, taking into account flavor oscillations, provide precise predictions for the actual phase-space distribution of relic neutrinos [5,6,7,8]. These distributions differ from the instantaneous decoupling approximation through a combination of a small shift in the photon temperature and small non-thermal distortions, all at the percent level. The final result for the density ratio  $\rho_\nu/\rho_\gamma$  in the relativistic regime can always be expressed as in Eq. (25.1), but with a different value of  $N_{\text{eff}}$ . The most recent analysis, that includes the effect of neutrino oscillations with the present values of the mixing parameters and an improved calculation of the collision terms, gives  $N_{\text{eff}} = 3.045$  [8]. The precise number density ratio  $n_\nu/n_\gamma$  can also be derived from such studies, and is important for computing the ratio  $\Omega_\nu h^2 / \sum_i m_i$  (ratio of the physical density of neutrinos in units of the critical density to the sum of neutrino masses) in the non-relativistic regime.

The neutrino temperature today,  $T_\nu^0 \simeq 1.7 \times 10^{-4} \text{ eV} \simeq 1.9 \text{ K}$ , is smaller than at least two of the neutrino masses, since the two squared-mass differences are  $|\Delta m_{31}^2|^{1/2} > |\Delta m_{21}^2|^{1/2} > T_\nu^0$  (see Neutrino mass, Mixing, and oscillations – Chap. 14 of this *Review*). Thus at least two neutrino mass eigenstates are non-relativistic today and behave as a small “hot” fraction of the total dark matter (they cannot be all the dark matter, as explained in Chap. 26 in this *Review*). This fraction of hot dark matter can be probed by cosmological experiments, for two related reasons, as we now describe.

First, neutrinos are the only known particles behaving as radiation at early times (during the CMB acoustic oscillations) and dark matter at late times (during structure formation), which has consequences on the background evolution. Neutrinos become non-relativistic when their mass is equal to their average momentum, given for any

Fermi-Dirac-distributed particle by  $\langle p \rangle = 3.15 T$ . Thus the redshift of the non-relativistic transition is given by  $z_i^{\text{nr}} = m_i/(3.15 T_\nu^0) - 1 = m_i/[0.53 \text{ meV}] - 1$  for each eigenstate of mass  $m_i$ , giving for instance  $z_i^{\text{nr}} = 110$  for  $m_i = 60 \text{ meV}$ , corresponding to a time deep inside the matter-dominated regime. Second, until the non-relativistic transition, neutrinos travel at the speed of light, and later on they move at a typical velocity  $\langle v_i/c \rangle = 3.15 T_\nu(z)/m_i = 0.53(1+z) \text{ meV}/m_i$ , which is several orders of magnitude larger than that of the dominant cold (or even of possibly warm) dark matter component(s). This brings their characteristic diffusion scale, called the “free-streaming length”, to cosmological relevant values, with consequences on gravitational clustering and the growth of structure.

Once neutrinos are non-relativistic, their energy density is given by  $\rho_\nu \simeq \sum m_i n_i$ . Since the number densities  $n_i$  are equal to each other (up to negligible corrections coming from flavour effects in the decoupling phase), the total mass ( $\sum m_\nu$ ) =  $m_1 + m_2 + m_3$  can be factorized out. It is possible that the lightest neutrino is still relativistic today, in which case this relation is slightly incorrect, but given that the total density is always strongly dominated by that of non-relativistic neutrinos, the error made is completely negligible. Using the expression for  $n_i/n_\gamma$  obtained from precise neutrino decoupling studies, and knowing  $n_\gamma$  from the measurement of the CMB temperature, one can compute  $\rho_\nu^0$ , the total neutrino density today, in units of the critical density  $\rho_{\text{crit}}^0$  [7]:


$$\Omega_\nu = \frac{\rho_\nu^0}{\rho_{\text{crit}}^0} = \frac{\sum m_\nu}{93.14 h^2 \text{ eV}}, \quad (25.2)$$

and the total neutrino average number density today:  $n_\nu^0 = 339.5 \text{ cm}^{-3}$ . Here  $h$  is the Hubble constant in units of  $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

## 25.2. Effects of neutrino properties on cosmological observables

As long as they are relativistic, *i.e.*, until some time deep inside the matter-dominated regime for neutrinos with a mass  $m_i \ll 3.15 T_\nu^{\text{eq}} \sim 1.5 \text{ eV}$  (see Big Bang Cosmology, Chap. 21 in this *Review*), neutrinos enhance the density of radiation: this effect is parameterised by  $N_{\text{eff}}$  and can be discussed separately from the effect of the mass that will be described later in this section. Increasing  $N_{\text{eff}}$  impacts the observable spectra of CMB anisotropies and matter fluctuations through background and perturbation effects.

**25.2.1. Effect of  $N_{\text{eff}}$  on the CMB :** The background effects depend on what is kept fixed when increasing  $N_{\text{eff}}$ . If the densities of other species are kept fixed, a higher  $N_{\text{eff}}$  implies a smaller redshift of radiation-to-matter equality, with very strong effects on the CMB spectrum: when the amount of expansion between radiation-to-matter equality and photon decoupling is larger, the CMB peaks are suppressed. This effect is not truly characteristic of the neutrino density, since it can be produced by varying several other parameters. Hence, to characterise the effect of  $N_{\text{eff}}$ , it is more useful and illuminating to enhance the density of total radiation, of total matter and of  $\Lambda$  by exactly the same amount, in order to keep the redshift of radiation-to-matter equality  $z_{\text{eq}}$  and matter-to- $\Lambda$  equality  $z_\Lambda$  fixed [9,10,4]. The primordial spectrum parameters, the baryon density




**Figure 25.1:** Ratio of the CMB  $C_\ell^{\text{TT}}$  (left, including lensing effects) and matter power spectrum  $P(k)$  (right, computed for each model in units of  $(h^{-1}\text{Mpc})^3$ ) for different values of  $\Delta N_{\text{eff}} \equiv N_{\text{eff}} - 3.045$  over those of a reference model with  $\Delta N_{\text{eff}} = 0$ . In order to minimize and better characterise the effect of  $N_{\text{eff}}$  on the CMB, the parameters that are kept fixed are  $\{z_{\text{eq}}, z_\Lambda, \omega_b, \tau\}$  and the primordial spectrum parameters. Fixing  $\{z_{\text{eq}}, z_\Lambda\}$  is equivalent to fixing the fractional density of total radiation, of total matter and of cosmological constant  $\{\Omega_r, \Omega_m, \Omega_\Lambda\}$  while increasing the Hubble parameter as a function of  $N_{\text{eff}}$ . The statistical errors on the  $C_\ell$  are  $\sim 1\%$  for a band power of  $\Delta\ell = 30$  at  $\ell \sim 1000$ . The error on  $P(k)$  is estimated to be of the order of  $5\%$ .

$\omega_b \equiv \Omega_b h^2$  and the optical depth to reionization  $\tau$  can be kept fixed at the same time, since we can simply vary  $N_{\text{eff}}$  together with the Hubble parameter  $h$  with fixed  $\{\omega_b, \Omega_c, \Omega_\Lambda\}$ . The impact of such a transformation is shown in Fig. 25.1 for the CMB temperature spectrum  $C_\ell^{TT}$  (defined in Chap. 28 in this *Review*) and for the matter power spectrum  $P(k)$  (defined in Chap. 21 in this *Review*) for several representative values of  $N_{\text{eff}}$ . These effects are within the reach of cosmological observations given current error bars, as discussed in Section 25.3 (for instance, with the *Planck* satellite data, the statistical error on the  $C_\ell$ 's is of the order of one per cent for a band power of  $\Delta\ell = 30$  at  $\ell \sim 1000$ ).

With this transformation, the main background effect of  $N_{\text{eff}}$  is an increase in the diffusion scale (or Silk damping scale, see Cosmic Microwave Background – Chap. 28 in this *Review*) at the time of decoupling, responsible for the decrease in  $C_\ell^{TT}$  at high  $\ell$ , plus smaller effects coming from a slight increase in the redshift of photon decoupling [4,9,10]. At the level of perturbations, a higher  $N_{\text{eff}}$  implies that photons feel gravitational forces from a denser neutrino component; this tends to decrease the acoustic peaks (because neutrinos are distributed in a smoother way than photons) and to shift them to larger scales / smaller multipoles (because photon perturbations traveling at the speed of sound in the photon-baryon fluid feel some dragging effect from neutrino perturbations travelling at the speed of light) [9,4,11]. The combination of these effects is truly characteristic of the radiation density parameter  $N_{\text{eff}}$  and cannot be mimicked by other parameters; thus  $N_{\text{eff}}$  can be accurately measured from the CMB alone. However, there are correlations between  $N_{\text{eff}}$  and other parameters. In particular, we have seen (Fig. 25.1) that in order to minimise the effect of  $N_{\text{eff}}$  on the CMB spectrum, one should vary  $h$  at the same time, hence there is a correlation between  $N_{\text{eff}}$  and  $h$ , which implies that independent measurements reducing the error bar on  $h$  also reduce that on  $N_{\text{eff}}$ . Note that this correlation is not equivalent to a perfect degeneracy, so both parameters can anyway be constrained with CMB data alone.

**25.2.2. Effect of  $N_{\text{eff}}$  on the matter spectrum :** We have discussed the effect of increasing  $N_{\text{eff}}$  while keeping  $z_{\text{eq}}$  and  $\omega_b$  fixed, because the latter two quantities are very accurately constrained by CMB data. This implies that  $\omega_c$  increases with  $N_{\text{eff}}$ , and that the ratio  $\omega_b/\omega_c = \Omega_b/\Omega_c$  decreases. However, the ratio of baryonic-to-dark matter has a strong impact on the shape of the matter power spectrum, because until the time of decoupling of the baryons from the photons, CDM experiences gravitational collapse, while baryons are kept smoothly distributed by photon pressure and affected by acoustic oscillations. The decrease of  $\Omega_b/\Omega_c$  following from the increase of  $N_{\text{eff}}$  gives more weight to the most clustered of the two components, namely the dark matter one, and produces an enhancement of the small-scale matter power spectrum and a damping of the amplitude of baryon acoustic oscillations (BAOs), clearly visible in Fig. 25.1 (right plot). The scale of BAOs is also slightly shifted.

The increase in the small-scale matter power spectrum is also responsible for a last effect on the CMB spectra : the CMB last scattering surface is slightly more affected by weak lensing from large-scale structures. This tends to smooth the maxima, the minima, and the damping scale of the CMB spectra [12].



**Figure 25.2:** Ratio of the CMB  $C_\ell^{TT}$  and matter power spectrum  $P(k)$  (computed for each model in units of  $(h^{-1}\text{Mpc})^3$ ) for different values of  $\sum m_\nu$  over those of a reference model with massless neutrinos. In order to minimize and better characterise the effect of  $\sum m_\nu$  on the CMB, the parameters that are kept fixed are  $\omega_b$ ,  $\omega_c$ ,  $\tau$ , the angular scale of the sound horizon  $\theta_s$  and the primordial spectrum parameters (solid lines). This implies that we are increasing the Hubble parameter  $h$  as a function of  $\sum m_\nu$ . For the matter power spectrum, in order to single out the effect of neutrino free-streaming on  $P(k)$ , the dashed lines show the spectrum ratio when  $\{\omega_m, \omega_b, \Omega_\Lambda\}$  are kept fixed. For comparison, the error on  $P(k)$  is of the order of 5% with current observations, and the fractional  $C_\ell$  errors are of the order of  $1/\sqrt{\ell}$  at low  $\ell$ .

**25.2.3. Effect of neutrino masses on the CMB :** Neutrino eigenstates with a mass  $m_i \ll 0.57$  eV become non-relativistic after photon decoupling. They contribute to the non-relativistic matter budget today, but not at the time of equality or recombination. If we increase the neutrino mass while keeping fixed the density of baryons and dark matter ( $\omega_b$  and  $\omega_c$ ), the early cosmological evolution remains fixed and independent of the neutrino mass, until the time of the non-relativistic transition. Thus one might expect that the CMB temperature and polarisation power spectra are left invariant. This is not true for four reasons.

First, the neutrino density enhances the total non-relativistic density at late times,  $\omega_m = \omega_b + \omega_c + \omega_\nu$ , where  $\omega_\nu \equiv \Omega_\nu h^2$  is given as a function of the total mass  $\sum m_\nu$  by Eq. (25.2). The late background evolution impacts the CMB spectrum through the relation between scales on the last scattering surface and angles on the sky, and through the late ISW effect (see Cosmic Microwave Background – Chap. 28 of this *Review*). These two effects depend respectively on the angular diameter distance to recombination,  $d_A(z_{\text{rec}})$ , and on the redshift of matter-to- $\Lambda$  equality. Increasing  $\sum m_\nu$  tends to modify these two quantities. By playing with  $h$  and  $\Omega_\Lambda$ , it is possible to keep one of them fixed, but not both at the same time. Since the CMB measures the angular scale of acoustic oscillations with exquisite precision, and is only loosely sensitive to the late ISW effect due to cosmic variance, we choose in Fig. 25.2 to play with the Hubble parameter in order to maintain a fixed scale  $d_A(z_{\text{rec}})$ . With such a choice, an increase in neutrino mass comes together with a decrease in the late ISW effect explaining the depletion of the CMB spectrum for  $l \leq 20$ . The fact that both  $\sum m_\nu$  and  $h$  enter the expression of  $d_A(z_{\text{rec}})$  implies that measurements of the neutrino mass from CMB data are strongly correlated with  $h$ . Second, the non-relativistic transition of neutrinos affects the total pressure-to-density ratio of the universe, and causes a small variation of the metric fluctuations. If this transition takes place not too long after photon decoupling, this variation is observable through the early ISW effect [4,13,14]. It is responsible for the dip seen in Fig. 25.2 for  $20 \leq \ell \leq 200$ . Third, when the neutrino mass is higher, the CMB spectrum is less affected by the weak lensing effect induced by the large-scale structure at small redshift. This is due to a decrease in the matter power spectrum described in the next paragraphs. This reduced lensing effect is responsible for most of the oscillatory patterns visible in Fig. 25.2 (left plot) for  $\ell \geq 200$ . Fourth, the neutrinos with the smallest momenta start to be non-relativistic earlier than the average ones. The photon perturbations feel this through their gravitational coupling with neutrinos. This leads to a small enhancement of  $C_l^{TT}$  for  $\ell \geq 500$ , hardly visible on Fig. 25.2 because it is balanced by the lensing effect.

**25.2.4. Effect of neutrino masses on the matter spectrum :** The physical effect of neutrinos on the matter power spectrum is related to their velocity dispersion. Neutrinos free-stream over large distances without falling into small potential wells. The free-streaming scale is roughly defined as the distance travel by neutrinos over a Hubble time scale  $t_H = (a/\dot{a})$ , and approximates the scale below which neutrinos remain very smooth. On larger scales, they cluster in the same way as cold dark matter. The power spectrum of total matter fluctuations, related to the squared fluctuation  $\delta_m^2$  with  $\delta_m \equiv \delta_b + \delta_c + \delta_\nu$ , gets a negligible contribution from the neutrino component on small

scales, and is reduced by a factor  $(1 - 2f_\nu)$ , where  $f_\nu = \omega_\nu/\omega_m$ . Additionally, on scales below the free-streaming scale, the growth of ordinary cold dark matter and baryon fluctuations is modified by the fact that neutrinos contribute to the background density, but not to the density fluctuations. This changes the balance between the gravitational forces responsible for clustering, and the Hubble friction term slowing it down. Thus the growth rate of CDM and baryon fluctuations is reduced [15]. This results today in an additional suppression of the small-scale linear matter power spectrum by approximately  $(1 - 6f_\nu)$ . These two effects sum up to a factor  $(1 - 8f_\nu)$  [16] (more precise approximations can be found in [2,4]). The non-linear spectrum is even more suppressed on mildly non-linear scales [17,18,19,20,21,3].

This effect is often illustrated by plots of the matter power spectrum ratio with fixed parameters  $\{\omega_m, \omega_b, \Omega_\Lambda\}$  and varying  $f_\nu$ , *i.e.*, with the CDM density adjusted to get a fixed total dark matter density [2,4,16] (see Fig. 25.2, right plot, dashed lines). This transformation does not leave the redshift of equality  $z_{\text{eq}}$  invariant, and has very large effects on the CMB spectra. If one follows the logic of minimizing CMB variations and fixing  $z_{\text{eq}}$  like in the previous paragraphs, the increase in  $\sum m_\nu$  must take place together with an increase of  $h$ , which tends to suppress the large-scale power spectrum, by approximately the same amount as the neutrino free-streaming effect [22]. In that case, the impact of neutrino masses on the matter power spectrum appears as an overall amplitude suppression, which can be seen in Fig. 25.2 (right plot, solid lines). The oscillations on intermediate wavenumbers come from a small shift in the BAO scale [22]. This global effect is not degenerate with a variation of the primordial spectrum amplitude  $A_s$ , because it only affects the matter power spectrum, and not the CMB spectra. However, the amplitude of the CMB temperature and polarization spectrum is given by the combination  $A_s e^{-2\tau}$ . Hence a measurement of  $\tau$  is necessary in order to fix  $A_s$  from CMB data, and avoid a parameter degeneracy between  $\sum m_\nu$  and  $A_s$  [22,23,24].

A few of the neutrino mass effects described above—free-streaming scale, early ISW—depend on individual masses  $m_i$ , but most of them depend only on the total mass through  $f_\nu$ —suppression of the matter power spectrum, CMB lensing, shift in angular diameter distance—. Because the latter effects are easier to measure, cosmology is primarily sensitive to the total mass  $\sum m_\nu$  [25,26]. The possibility that future data sets might be able to measure individual masses or the mass hierarchy, despite systematic errors and parameter degeneracies, has recently become a subject of investigation [27,28].

### 25.3. Cosmological Constraints on neutrino properties

In this review we focus on cosmological constraints on the abundance and mass of ordinary active neutrinos. Several stringent but model-dependent constraints on non-standard neutrinos (*e.g.*, sterile neutrinos, active neutrinos with interactions beyond the weak force, unstable neutrinos with invisible decay, etc.) can also be found in the literature.



### 25.3.1. Neutrino abundance :

Table 25.1 shows a list of constraints on  $N_{\text{eff}}$  obtained with several combination of data sets. ‘P15’ denotes the *Planck* 2015 data, composed of a high- $\ell$  temperature likelihood (TT), low- $\ell$  temperature+polarization likelihood (lowP) and CMB lensing spectrum likelihood (lensing) based on lensing extraction from quadratic estimators [29]. ‘BAO’ refers to measurements of the BAO scale (and hence of the angular diameter distance) from various recent data sets, described in detail in the references given in the table. ‘JLA’ refers to the supernovae luminosity distance measurements [30]. ‘HST’ refers to the direct measurement of the Hubble scale from cepheids and supernovae [31].

Within the framework of a 7-parameter cosmological model ( $\Lambda$ CDM+ $N_{\text{eff}}$ ), the most conservative constraint on  $N_{\text{eff}}$  comes from the *Planck* 2015 data release with robust temperature and large-angle polarization information:  $N_{\text{eff}} = 3.13 \pm 0.32$  (68%CL). This number is perfectly compatible with the prediction of the standard neutrino decoupling model,  $N_{\text{eff}} = 3.045$ , and can be viewed as a proof of self-consistency of the cosmological model.

The bounds can be tightened by adding more CMB polarization data, but the *Planck* Collaboration warns that there might be some unremoved systematics in the full TT,TE,EE likelihood; or, more conservatively, by adding information on the low-redshift background expansion from BAOs, supernovae or direct  $H_0$  measurements. Finally, one can also add information on large scale structure (LSS), *i.e.*, on the growth rate and clustering amplitude of matter as a function of scale. However, LSS data are not very constraining for the  $N_{\text{eff}}$  parameter, and the only LSS data included in Table 25.1 is the measurement of the CMB lensing spectrum.

Combinations of *Planck* 2015 data with BAO, supernovae or CMB lensing constraints, all return measurements consistent with the standard expectation.

The situation is different with the inclusion of the direct measurement of  $H_0$  heavily relying on the *Hubble Space Telescope* (HST) data, [31], known to be in tension with *Planck* in the  $\Lambda$ CDM framework. As explained in Section 25.2, the positive correlation between  $N_{\text{eff}}$  and  $h$  means that inclusion of the  $H_0$  measurement pushes  $N_{\text{eff}}$  to higher values,  $N_{\text{eff}} = 3.41 \pm 0.22$  (68%CL, P15[TT+lowP+lensing] + BAO + JLA + HST), but still compatible with the standard expectation at the  $1.7\sigma$  level. It remains to be seen whether the  $3.2\sigma$  tension between CMB data and direct measurements of  $H_0$  results from systematics, or from a departure from the  $\Lambda$ CDM model [33].

The error bars on  $N_{\text{eff}}$  degrade when the data are analysed in the context of more extended cosmological scenarios. Adding the neutrino mass as an 8th free parameter, the P15[TT+lowP+lensing]+BAO data set of Ref. [29] returns  $N_{\text{eff}} = 3.2 \pm 0.5$  instead of  $3.07 \pm 0.23$  (68%CL). The authors of Ref. [34] take an extreme point of view and fit a 12-parameter model to P15[TT,TE,EE+lowP+lensing] data; they obtain  $N_{\text{eff}} = 2.93^{+0.51}_{-0.48}$  (95%CL), showing that it is very difficult with current cosmological data to accommodate shifts of more than 0.5 from the standard  $N_{\text{eff}}$  value, and to obtain good fits with, for instance, a fourth (sterile) thermalized neutrino. This is interesting since the anomalies in some oscillation data could be interpreted as evidence for at least one sterile neutrino with a large mixing angle, which would need to be thermalised unless

non-standard interactions come into play [36]. In other words cosmology disfavours the explanation of the oscillations anomalies in terms of 1 or more extra neutrinos if they are thermalized.

**Table 25.1:** Summary of  $N_{\text{eff}}$  constraints.

	Model	68%CL	Ref.
<b>CMB alone</b>			
Pl15[TT+lowP]	$\Lambda\text{CDM}+N_{\text{eff}}$	$3.13 \pm 0.32$	[29]
Pl15[TT+lowP]	$\Lambda\text{CDM}+N_{\text{eff}}+\sum m_\nu$	$3.08 \pm 0.31$	[35]
<b>CMB + probes of background evolution</b>			
Pl15[TT+lowP] + BAO	$\Lambda\text{CDM}+N_{\text{eff}}$	$3.15 \pm 0.23$	[29]
Pl15[TT+lowP] + BAO	$\Lambda\text{CDM}+N_{\text{eff}}+\sum m_\nu$	$3.18^{+0.24}_{-0.27}$	[35]
<b>CMB + probes of background evolution + LSS</b>			
Pl15[TT+lowP+lensing] + BAO	$\Lambda\text{CDM}+N_{\text{eff}}$	$3.08^{+0.22}_{-0.24}$	[35]
” + BAO + JLA + HST	$\Lambda\text{CDM}+N_{\text{eff}}$	$3.41 \pm 0.22$	[31]
” + BAO	$\Lambda\text{CDM}+N_{\text{eff}}+\sum m_\nu$	$3.2 \pm 0.5$	[29]
Pl15[TT,TE,EE+lowP+lensing]	$\Lambda\text{CDM}+N_{\text{eff}}+5\text{-params.}$	$2.93^{+0.51}_{-0.48}$	[34]

**25.3.2. Are they really neutrinos, as expected?** : While a value of  $N_{\text{eff}}$  significantly different from zero (at more than  $10\sigma$ ) and consistent with the expected number 3.045 yields a powerful indirect confirmation of the  $\text{C}\nu\text{B}$ , departures from standard  $N_{\text{eff}}$  could be caused by any ingredient affecting the early-time expansion rate of the Universe. Extra relativistic particles (either decoupled, self-interacting, or interacting with a dark sector), a background of gravitational waves, an oscillating scalar field with quartic potential, departures from Einstein gravity, or large extra dimensions are some of the possibilities for such ingredients. In principle one could even assume that the cosmic neutrino background never existed or has decayed (like in the “neutrinoless universe” model of [37]) while another dark radiation component is responsible for  $N_{\text{eff}}$ . At least, cosmological data allow to narrow the range of possible interpretations of  $N_{\text{eff}} \simeq 3$  to the presence of decoupled relativistic relics like standard neutrinos. Indeed, free-streaming particles leave specific signatures, especially in the CMB, because their density and pressure perturbations, bulk velocities and anisotropic stress also source the metric perturbations. These signatures can be tested in several ways.

A first approach consists of introducing a self-interaction term in the neutrino equations. Refs [38,39] find that the Pl15+BAO data are compatible with no self-interactions. The upper limits to the effective coupling constant  $G_{\text{eff}}$  for a Fermi-like

four-fermions interaction at 95% confidence is  $\log_{10}(G_{\text{eff}}\text{MeV}^2) < -3.5(-2.7)$  for Planck CMB temperature data only [38]( +BAO [39]) .

A second approach consists of introducing two phenomenological parameters,  $c_{\text{eff}}$  and  $c_{\text{vis}}$  (see *e.g.*, [40,41,42]):  $c_{\text{eff}}^2$  generalizes the linear relation between isotropic pressure perturbations and density perturbations, while  $c_{\text{vis}}^2$  modifies the neutrino anisotropic stress equation. While relativistic free-streaming species have  $(c_{\text{eff}}^2, c_{\text{vis}}^2) = (1/3, 1/3)$ , a perfect relativistic fluid would have  $(c_{\text{eff}}^2, c_{\text{vis}}^2) = (1/3, 0)$ . Other values do not necessarily refer to a concrete model, but make it possible to interpolate between these limits. The latest *Planck* data strongly suggests  $(c_{\text{eff}}^2, c_{\text{vis}}^2) = (1/3, 1/3)$  [43,29]. Finally, Ref. [11] shows that *Planck* data are precise enough to detect the “neutrino drag” effect mentioned in Sec. 25.2, caused by gravitational interactions between neutrino and photon perturbations, and shifting the CMB peaks towards larger angular scales. These findings show that current cosmological data are able to detect not just the average density of some relativistic relics, but also their anisotropies.

### 25.3.3. *Neutrino masses* :

Table 25.2 shows a list of constraints on  $\sum m_\nu$  obtained with several combinations of data sets. The acronyms “Pl15”, “BAO”, “JLA”, and “HST” have been described in the previous subsection. “Pl16” refers to Planck intermediate results from 2016 in which the high- $\ell$  and lensing likelihood are identical to the 2015 version, but the low- $\ell$  temperature+polarization likelihood based on the Low Frequency Instrument (lowP) data is replaced by a newer version based on the High Frequency Instrument (SimLow) [32]. There “P(k)” refers to the several measurements of the matter power spectrum shape (and hence of the growth rate and of the clustering amplitude as function of scale) for “WZ”, by the WiggleZ survey [44], for “DR7”, from the Data Release 7 of the Sloan Digital Sky Survey [45], and for “DR12”, from the Data Release 12 of the Baryon Oscillation Spectroscopic Survey (BOSS) [46,47]. “Ly $\alpha$ ” refers to the BOSS measurement of the Lyman- $\alpha$  flux power spectrum in quasar spectra [48].

Given that most determinations of  $N_{\text{eff}}$  are compatible with the standard prediction,  $N_{\text{eff}} = 3.045$ , it is reasonable to adopt this value as a theoretical prior and to investigate neutrino mass constraints in the context of a minimal 7-parameter model,  $\Lambda\text{CDM} + \sum m_\nu$ . Under this assumption, the most robust constraints come from *Planck* temperature data and large-angle polarization information:  $\sum m_\nu < 0.72\text{ eV}$  (95%CL) using the 2015 low- $\ell$  polarization likelihood [29], or  $\sum m_\nu < 0.59\text{ eV}$  (95%CL) using the one from 2016 [32] (see also Ref. [49]). The high- $\ell$  polarization likelihood from *Planck* 2015, which should be used with caution, pushes the bound to  $\sum m_\nu < 0.34\text{ eV}$  (95%CL). Among the four effects of neutrino masses on the CMB spectra described before, current bounds are dominated by the first and the third effects (modified late background evolution, and distortions of the temperature and polarisation spectra through weak lensing).

Adding data on BAO scales is crucial, since the measurement of the angular diameter distance at small redshift allows us to break parameter degeneracies, for instance between  $\sum m_\nu$  and  $h$ . Combined with conservative Planck 2015 data, BAO experiments give  $\sum m_\nu < 0.21\text{ eV}$  (95%CL). Supernovae data are less constraining than BAO data for the neutrino mass determination. Because the parameter correlation between  $\sum m_\nu$  and  $H_0$

**Table 25.2:** Summary of  $\sum m_\nu$  constraints.

	Model	95% CL (eV)	Ref.
<b>CMB alone</b>			
Pl15[TT+lowP]	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.72$	[29]
Pl15[TT+lowP]	$\Lambda\text{CDM}+\sum m_\nu+N_{\text{eff}}$	$< 0.73$	[35]
Pl16[TT+SimLow]	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.59$	[32]
<b>CMB + probes of background evolution</b>			
Pl15[TT+lowP] + BAO	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.21$	[29]
Pl15[TT+lowP] + JLA	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.33$	[35]
Pl15[TT+lowP] + BAO	$\Lambda\text{CDM}+\sum m_\nu+N_{\text{eff}}$	$< 0.27$	[35]
<b>CMB + probes of background evolution + LSS</b>			
Pl15[TT+lowP+lensing]	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.68$	[29]
Pl15[TT+lowP+lensing] + BAO	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.25$	[35]
Pl15[TT+lowP] + $P(k)_{\text{DR12}}$	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.30$	[50]
Pl15[TT,TE,EE+lowP] + BAO+ $P(k)_{\text{WZ}}$	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.14$	[52]
Pl15[TT,TE,EE+lowP] + BAO+ $P(k)_{\text{DR7}}$	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.13$	[52]
Pl15[TT+lowP+lensing] + $\text{Ly}\alpha$	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.12$	[48]
Pl16[TT+SimLow+lensing] + BAO	$\Lambda\text{CDM}+\sum m_\nu$	$< 0.17$	[48]
Pl15[TT+lowP+lensing] + BAO	$\Lambda\text{CDM}+\sum m_\nu+\Omega_k$	$< 0.37$	[35]
Pl15[TT+lowP+lensing] + BAO	$\Lambda\text{CDM}+\sum m_\nu+w$	$< 0.37$	[35]
Pl15[TT+lowP+lensing] + BAO	$\Lambda\text{CDM}+\sum m_\nu+N_{\text{eff}}$	$< 0.32$	[29]
Pl15[TT,TE,EE+lowP+lensing]	$\Lambda\text{CDM}+\sum m_\nu+5\text{-params.}$	$< 0.66$	[34]

is negative, the inclusion of HST data provides stronger bounds on neutrinos masses, down to  $\sum m_\nu < 0.11$  eV (95% CL) when including LSS [52], but such bounds are subject to caution, since they come from a combination of slightly discrepant data sets (at the  $3.2\sigma$  level).

It is interesting to add LSS data sets, sensitive to the small-scale suppression of the matter power spectrum due to neutrino free-streaming. The inclusion of the *Planck* 2015 CMB lensing likelihood is not very constraining. Overall, adding CMB lensing to conservative *Planck* 2015 data gives stronger bounds, but only marginally (from  $\sum m_\nu < 0.72$  eV to  $\sum m_\nu < 0.68$  eV at 95%CL). The inclusion of several matter power spectrum determinations, listed in Table 25.2, also provides rather marginal improvements: the constraint  $\sum m_\nu < 0.17$  eV (95%CL) from Pl15[TT,TE,EE+lowP]+BAO is only pushed down to 0.14 eV (0.13 eV, 0.16 eV) when adding matter power spectrum data from WiggleZ (blue galaxies) [50]( SDSS-DR7 [52], BOSS-DR12 [52], red galaxies). The Lyman- $\alpha$  power spectrum data from BOSS are more constraining, since this leads to

$\sum m_\nu < 0.12 \text{ eV}$  (95% CL) in the absence of BAO or high- $\ell$  polarization data, but only with P115[TT+lowP+lensing] [48]. However, it is often stressed that the bounds coming from Lyman- $\alpha$  data involve more modelling of non-linear effects than the other techniques presented in this summary.

Upper bounds on neutrino masses become weaker when the data are analysed in the context of extended cosmological models, but not considerably weaker. For instance, one can see in Table 25.2 that bounds from P115[TT+lowP+lensing] + BAO tend to degrade from 0.25 eV to 0.37 eV (95% CL) when introducing an 8th free parameter accounting for spatial curvature or dynamical dark energy. In the extreme case considered by Ref. [34], with 12 free cosmological parameters, the bound from P115[TT,TE,EE+lowP+lensing] increases from 0.59 eV to 0.66 eV (95% CL). This shows that current cosmological data are precise enough to disentangle the effect of several extended cosmological parameters, and that neutrino mass bounds are becoming increasingly robust.

## 25.4. Future prospects and outlook

The cosmic neutrino background has been detected indirectly at very high statistical significance. Direct detection experiments are now being planned, *e.g.*, at the Princeton Tritium Observatory for Light, Early-universe, Massive-neutrino Yield (PTOLEMY) [53]. The detection prospects crucially depend on the exact value of neutrino masses and on the enhancement of their density at the location of the Earth through gravitational clustering in the Milky Way and its sub-halos – an effect however expected to be small [54,55,56].

Over the past few years the upper limit on the sum of neutrino masses has become increasingly stringent, first indicating that the mass ordering is hierarchical and recently putting the inverted hierarchy under pressure and favouring the normal hierarchy (although quantitative estimates of how disfavoured the inverted hierarchy is vary depending on assumptions, see *e.g.* [57,58]) which has consequences for planning future double beta decay experiments.

Neutrino mass and density bounds are expected to keep improving significantly over the next years, thanks to new LSS experiments like DES [59], Euclid [60], LSST [61], and SKA [62], or possible new CMB experiments like CMB-S4 [63], Pixie [65], CMBPol or CORE [64]. If the  $\Lambda$ CDM model is confirmed, and if neutrinos have standard properties, the total neutrino mass should be detected at the level of at least  $3\text{--}4\sigma$  even at the minimum level allowed by oscillations. This is the conclusion reached by several independent studies, using different dataset combinations (see *e.g.*, [66,67,68,69,70,71]). One should note that at the minimum level allowed by oscillations  $\sum m_\nu \sim 0.06$ , neutrinos constitute  $\sim 0.5\%$  of the Universe matter density, and their effects on the matter power spectrum is only at the 5% level, implying that exquisite control of systematic errors will be crucial to achieve the required accuracy. At this level, the information coming from the power spectrum shape is more powerful than that coming from geometrical measurements (*e.g.*, BAO). But exploiting the shape information requires improved understanding of the non-linear regime, and of galaxy bias for galaxy surveys. The fact that different surveys and different data set combinations have enough statistical power to reach this level, offers a much needed redundancy and the possibility to perform consistency checks

which in turns helps immensely with the control of systematic errors and in making the measurement robust. Using the entire Universe as a particle detector, the on-going and future observational efforts hold the exciting prospect to provide a measurement of the sum of neutrino masses and possibly indication of their mass hierarchy.

### References:

1. A. D. Dolgov, Phys. Reports **370**, 333 (2002).
2. J. Lesgourgues and S. Pastor, Phys. Reports **429**, 307 (2006).
3. S. Hannestad, Prog. in Part. Nucl. Phys. **65**, 185 (2010).
4. J. Lesgourgues *et al.*, *Neutrino cosmology* (Cambridge University Press, 2013).
5. J. Birrell, C.-T. Yang, and J. Rafelski, Nucl. Phys. **B890**, 481 (2014).
6. G. Mangano *et al.*, Nucl. Phys. **B729**, 221 (2005).
7. E. Grohs *et al.*, Phys. Rev. **D93**, 083522 (2016).
8. P. F. de Salas and S. Pastor, JCAP **1607**, 051 (2016).
9. S. Bashinsky and U. Seljak, Phys. Rev. **D69**, 083002 (2004).
10. Z. Hou *et al.*, Phys. Rev. **D87**, 083008 (2013).
11. B. Follin *et al.*, Phys. Rev. Lett. **115**, 091301 (2015).
12. A. Lewis and A. Challinor, Phys. Reports **429**, 1 (2006).
13. J. Lesgourgues and S. Pastor, Adv. High Energy Phys. **2012**, 608515 (2012).
14. Z. Hou *et al.*, Astrophys. J. **782**, 74 (2014).
15. J. R. Bond, G. Efstathiou and J. Silk, Phys. Rev. Lett. **45**, 1980 (1980).
16. W. Hu, D. J. Eisenstein and M. Tegmark, Phys. Rev. Lett. **80**, 5255 (1998).
17. S. Bird, M. Viel and M. G. Haehnelt, Mon. Not. R. Astron. Soc **420**, 2551 (2012).
18. C. Wagner, L. Verde and R. Jimenez, Astrophys. J. **752**, L31 (2012).
19. E. Castorina *et al.*, JCAP **1507**, 042 (2015).
20. J. Brandbyge and S. Hannestad, arXiv:1706.00025 [astro-ph.CO].
21. J. Adamek, R. Durrer and M. Kunz, arXiv:1707.06938 [astro-ph.CO].
22. M. Archidiacono *et al.*, JCAP **1702**, 052 (2017).
23. A. Liu *et al.*, Phys. Rev. **D93**, 043013 (2016).
24. R. Allison *et al.*, Phys. Rev. **D92**, 123535 (2015).
25. J. Lesgourgues, S. Pastor and L. Perotto, Phys. Rev. **D70**, 045016 (2004).
26. A. Slosar, Phys. Rev. **D73**, 123501 (2006).
27. R. Jimenez *et al.*, JCAP **1005**, 035 (2010).
28. R. Jimenez, C. Pena-Garay and L. Verde, Phys. Dark Univ. **15**, 31 (2017).
29. P. A. R. Ade *et al.*, [Planck Collaboration], Astron. & Astrophys. **594**, A13 (2016).
30. M. Betoule *et al.*, [SDSS Collaboration], Astron. & Astrophys. **568**, A22 (2014).
31. A. G. Riess *et al.*, Astrophys. J. **826**, 56 (2016).
32. N. Aghanim *et al.*, [Planck Collaboration], Astron. & Astrophys. **596**, A107 (2016).
33. J. L. Bernal, L. Verde and A. G. Riess, JCAP **1610**, 019 (2016).
34. E. Di Valentino, A. Melchiorri and J. Silk, Phys. Rev. **D92**, 12,1302 (2015).
35. Planck 2015 explanatory supplement, [wiki.cosmos.esa.int/planckpla2015/](http://wiki.cosmos.esa.int/planckpla2015/), “Mission Products: Cosmological parameters: Parameter Tables”.
36. M. Archidiacono *et al.*, JCAP **1608**, 067 (2016).
37. J. F. Beacom, N. F. Bell and S. Dodelson, Phys. Rev. Lett. **93**, 121302 (2004).
38. L. Lancaster *et al.*, JCAP **1707**, 033 (2017).

39. I. M. Oldengott *et al.*, [arXiv:1706.02123](#) [astro-ph.CO].
40. W. Hu, *Astrophys. J.* **506**, 485 (1998).
41. W. Hu *et al.*, *Phys. Rev.* **D59**, 023512 (1999).
42. M. Gerbino, E. Di Valentino and N. Said, *Phys. Rev.* **D88**, 063538 (2013).
43. B. Audren *et al.*, *JCAP* **1503**, 036 (2015).
44. S. Riemer-Sorensen, D. Parkinson and T. M. Davis, *Phys. Rev. D* **89**, 103505 (2014).
45. B. A. Reid *et al.*, *Mon. Not. R. Astron. Soc* **404**, 60 (2010).
46. S. Alam *et al.*, [BOSS Collaboration], *Mon. Not. R. Astron. Soc* **470**, 2617 (2017).
47. H. Gil-Marín *et al.*, *Mon. Not. R. Astron. Soc* **460**, 4188 (2016).
48. N. Palanque-Delabrouille *et al.*, *JCAP* **1511**, 011 (2015).
49. F. Capozzi *et al.*, *Phys. Rev. D* **95**, 096014 (2017).
50. S. Vagnozzi *et al.*, [arXiv:1701.08172](#) [astro-ph.CO].
51. C. Yeche *et al.*, *JCAP* **1706**, 047 (2017).
52. A. J. Cuesta, V. Niro and L. Verde, *Phys. Dark Univ.* **13**, 77 (2016).
53. S. Betts *et al.*, [arXiv:1307.4738](#) [astro-ph.IM]..
54. A. Ringwald and Y. Y. Y. Wong, *JCAP* **0412**, 005 (2004).
55. F. Villaescusa-Navarro *et al.*, *JCAP* **03**, 019 (2013).
56. P. F. de Salas *et al.*, *JCAP* **1709**, 034 (2017).
57. F. Simpson *et al.*, *JCAP* **06**, 029 (2017).
58. S. Hannestad, T. Schwetz, *JCAP* **1611**, 035 (2016).
59. Dark Energy Survey Collaboration *et al.*, *Mon. Not. R. Astron. Soc* **460**, 1270 (2016).
60. R. Laureijs *et al.*, (2011) [arXiv:1110.3193](#).
61. Paul A. Abell *et al.*; LSST Science and LSST Project Collaborations, <http://lss.fnal.gov/archive/test-tm/2000/fermilab-tm-2495-a.pdf>, (2009) [arXiv:0912.0201](#).
62. <http://www.skatelescope.org>.
63. K. Abajiazan *et al.*, 2016, [arXiv:1610.02743](#).
64. J. Delabrouille *et al.*, 2017, [arXiv:1706.04516](#).
65. A. Kogut, *JCAP* **1107**, 025 (2011).
66. C. Carbone *et al.*, *JCAP* **1103**, 030 (2011).
67. J. Hamann, S. Hannestad, Y. Wong, *JCAP* **1211**, 052 (2012).
68. B. Audren *et al.*, *JCAP* **1301**, 026 (2013).
69. R. Pearson, O. Zhan, *Phys. Rev.* **D89**, 043516 (2014).
70. R. Alison *et al.*, *Phys. Rev.* **D92**, 123535 (2015).
71. F. Villaescusa-Navarro, P. Bull, M. Viel, *Astrophys. J.* **814**, 146 (2015).