

# QUARKS

The  $u$ -,  $d$ -, and  $s$ -quark masses are estimates of so-called “current-quark masses,” in a mass-independent subtraction scheme such as  $\overline{\text{MS}}$  at a scale  $\mu \approx 2 \text{ GeV}$ . The  $c$ - and  $b$ -quark masses are the “running” masses in the  $\overline{\text{MS}}$  scheme. For the  $b$ -quark we also quote the 1S mass. These can be different from the heavy quark masses obtained in potential models.

**u**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_u = 2.2^{+0.5}_{-0.4} \text{ MeV} \quad \text{Charge} = \frac{2}{3} e \quad I_z = +\frac{1}{2}$$

$$m_u/m_d = 0.48^{+0.07}_{-0.08}$$

**d**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_d = 4.7^{+0.5}_{-0.3} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad I_z = -\frac{1}{2}$$

$$m_s/m_d = 17-22$$

$$\overline{m} = (m_u + m_d)/2 = 3.5^{+0.5}_{-0.2} \text{ MeV}$$

**s**

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_s = 95^{+9}_{-3} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Strangeness} = -1$$

$$m_s / ((m_u + m_d)/2) = 27.3 \pm 0.7$$

**c**

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_c = 1.275^{+0.025}_{-0.035} \text{ GeV} \quad \text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

$$m_c/m_s = 11.72 \pm 0.25$$

$$m_b/m_c = 4.53 \pm 0.05$$

$$m_b - m_c = 3.45 \pm 0.05 \text{ GeV}$$

**b**

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Mass } m = 4.18^{+0.04}_{-0.03} \text{ GeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

**t**

$I(J^P) = 0(\frac{1}{2}^+)$

Charge =  $\frac{2}{3} e$       Top = +1

Mass (direct measurements)  $m = 173.0 \pm 0.4 \text{ GeV}^{[a,b]}$  (S = 1.3)  
Mass (from cross-section measurements)  $m = 160^{+5}_{-4} \text{ GeV}^{[a]}$   
Mass (Pole from cross-section measurements)  $m = 173.1 \pm 0.9 \text{ GeV}$   
 $m_t - m_{\bar{t}} = -0.16 \pm 0.19 \text{ GeV}$   
Full width  $\Gamma = 1.41^{+0.19}_{-0.15} \text{ GeV}$  (S = 1.4)  
 $\Gamma(Wb)/\Gamma(Wq(q = b, s, d)) = 0.957 \pm 0.034$  (S = 1.5)

**t-quark EW Couplings**

$F_0 = 0.687 \pm 0.018$   
 $F_- = 0.320 \pm 0.013$   
 $F_+ = 0.002 \pm 0.011$   
 $F_{V+A} < 0.29, \text{ CL} = 95\%$

<b>t DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$\frac{P}{(\text{MeV}/c)}$
$t \rightarrow Wq(q = b, s, d)$			—
$t \rightarrow Wb$			—
$t \rightarrow e\nu_e b$	(13.3±0.6) %		—
$t \rightarrow \mu\nu_\mu b$	(13.4±0.6) %		—
$t \rightarrow \tau\nu_\tau b$	( 7.1±0.6) %		—
$t \rightarrow q\bar{q}b$	(66.5±1.4) %		—

**$\Delta T = 1$  weak neutral current (T1) modes**

$t \rightarrow Zq(q=u,c)$	T1	[c] < 5	$\times 10^{-4}$	95%	—
$t \rightarrow Hu$	T1	< 2.4	$\times 10^{-3}$	95%	—
$t \rightarrow Hc$	T1	< 2.2	$\times 10^{-3}$	95%	—
$t \rightarrow \ell^+ \bar{q} \bar{q}' (q=d,s,b; q'=u,c)$	T1	< 1.6	$\times 10^{-3}$	95%	—

**$b'$  (4<sup>th</sup> Generation) Quark, Searches for**

Mass  $m > 190 \text{ GeV}$ , CL = 95% ( $p\bar{p}$ , quasi-stable  $b'$ )  
Mass  $m > 755 \text{ GeV}$ , CL = 95% ( $pp$ , neutral-current decays)  
Mass  $m > 880 \text{ GeV}$ , CL = 95% ( $pp$ , charged-current decays)  
Mass  $m > 46.0 \text{ GeV}$ , CL = 95% ( $e^+e^-$ , all decays)

## **$t'$ ( $4^{th}$ Generation) Quark, Searches for**

$$m(t'(2/3)) > 1160 \text{ GeV, CL} = 95\% \quad (\text{neutral-current decays})$$

$$m(t'(2/3)) > 770 \text{ GeV, CL} = 95\% \quad (\text{charged-current decays})$$

$$m(t'(5/3)) > 990 \text{ GeV, CL} = 95\%$$

## **Free Quark Searches**

All searches since 1977 have had negative results.

### NOTES

- [a] A discussion of the definition of the top quark mass in these measurements can be found in the review “The Top Quark.”
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II and LHC at  $\sqrt{s} = 7 \text{ TeV}$ . Including the most recent unpublished results from Tevatron Run-II, the Tevatron Electroweak Working Group reports a top mass of  $173.2 \pm 0.9 \text{ GeV}$ . See the note “The Top Quark” in the Quark Particle Listings of this *Review*.
- [c] This limit is for  $\Gamma(t \rightarrow Zq)/\Gamma(t \rightarrow Wb)$ .