

QUARKS

The u -, d -, and s -quark masses are estimates of so-called “current-quark masses,” in a mass-independent subtraction scheme such as $\overline{\text{MS}}$ at a scale $\mu \approx 2 \text{ GeV}$. The c - and b -quark masses are the “running” masses in the $\overline{\text{MS}}$ scheme. This can be different from the heavy quark masses obtained in potential models.

u

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_u = 2.16^{+0.49}_{-0.26} \text{ MeV} \quad \text{Charge} = \frac{2}{3} e \quad I_z = +\frac{1}{2}$$

$$m_u/m_d = 0.47^{+0.06}_{-0.07}$$

d

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_d = 4.67^{+0.48}_{-0.17} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad I_z = -\frac{1}{2}$$

$$m_s/m_d = 17\text{--}22$$

$$\bar{m} = (m_u + m_d)/2 = 3.45^{+0.55}_{-0.15} \text{ MeV}$$

s

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_s = 93^{+11}_{-5} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Strangeness} = -1$$

$$m_s / ((m_u + m_d)/2) = 27.3^{+0.7}_{-1.3}$$

c

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_c = 1.27 \pm 0.02 \text{ GeV} \quad \text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

$$m_c/m_s = 11.72 \pm 0.25$$

$$m_b/m_c = 4.577 \pm 0.008$$

$$m_b - m_c = 3.45 \pm 0.05 \text{ GeV}$$

b

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_b = 4.18^{+0.03}_{-0.02} \text{ GeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

t

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Top} = +1$$

Mass (direct measurements) $m = 172.9 \pm 0.4 \text{ GeV}^{[a,b]}$ ($S = 1.3$)
 Mass (from cross-section measurements) $m = 160_{-4}^{+5} \text{ GeV}^{[a]}$
 Mass (Pole from cross-section measurements) $m = 173.1 \pm 0.9 \text{ GeV}$
 $m_t - m_{\bar{t}} = -0.16 \pm 0.19 \text{ GeV}$
 Full width $\Gamma = 1.42_{-0.15}^{+0.19} \text{ GeV}$ ($S = 1.4$)
 $\Gamma(Wb)/\Gamma(Wq(q = b, s, d)) = 0.957 \pm 0.034$ ($S = 1.5$)

t-quark EW Couplings

$F_0 = 0.687 \pm 0.018$
 $F_- = 0.320 \pm 0.013$
 $F_+ = 0.002 \pm 0.011$
 $F_{V+A} < 0.29, \text{ CL} = 95\%$

t DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$t \rightarrow Wq(q = b, s, d)$			—
$t \rightarrow Wb$			—
$t \rightarrow e\nu_e b$	(13.3±0.6) %		—
$t \rightarrow \mu\nu_\mu b$	(13.4±0.6) %		—
$t \rightarrow \tau\nu_\tau b$	(7.1±0.6) %		—
$t \rightarrow q\bar{q}b$	(66.5±1.4) %		—

$\Delta T = 1$ weak neutral current ($T1$) modes

$t \rightarrow Zq(q=u,c)$	$T1$	$[c] < 5$	$\times 10^{-4}$	95%	—
$t \rightarrow Hu$	$T1$	< 1.9	$\times 10^{-3}$	95%	—
$t \rightarrow Hc$	$T1$	< 1.6	$\times 10^{-3}$	95%	—
$t \rightarrow \ell^+ \bar{q}\bar{q}' (q=d,s,b; q'=u,c)$	$T1$	< 1.6	$\times 10^{-3}$	95%	—

b' (4th Generation) Quark, Searches for

Mass $m > 190 \text{ GeV}, \text{ CL} = 95\%$ ($p\bar{p}$, quasi-stable b')
 Mass $m > 1350 \text{ GeV}, \text{ CL} = 95\%$ (pp , charged-current decays)
 Mass $m > 46.0 \text{ GeV}, \text{ CL} = 95\%$ (e^+e^- , all decays)

t' (4th Generation) Quark, Searches for

$m(t'(2/3)) > 1160 \text{ GeV}, \text{ CL} = 95\%$ (neutral-current decays)
 $m(t'(2/3)) > 1295 \text{ GeV}, \text{ CL} = 95\%$ (charged-current decays)
 $m(t'(5/3)) > 1.350 \times 10^3 \text{ GeV}, \text{ CL} = 95\%$

Free Quark Searches

All searches since 1977 have had negative results.

NOTES

- [a] A discussion of the definition of the top quark mass in these measurements can be found in the review “The Top Quark.”
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II and LHC at $\sqrt{s} = 7$ TeV. Including the most recent unpublished results from Tevatron Run-II, the Tevatron Electroweak Working Group reports a top mass of 173.2 ± 0.9 GeV. See the note “The Top Quark’ in the Quark Particle Listings of this *Review*.
- [c] This limit is for $\Gamma(t \rightarrow Z q)/\Gamma(t \rightarrow W b)$.