

$c\bar{c}$ MESONS (including possibly non- $q\bar{q}$ states)

 $\eta_c(1S)$

$$J^{PC} = 0^+(0^-+)$$

 Mass $m = 2983.9 \pm 0.5$ MeV ($S = 1.3$)

 Full width $\Gamma = 31.9 \pm 0.7$ MeV

$\eta_c(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
--	--------------------------------	------------------	----------------

Decays involving hadronic resonances

$\eta'(958)\pi\pi$	(4.1 \pm 1.7) %		1323
$\rho\rho$	(1.8 \pm 0.5) %		1275
$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(2.0 \pm 0.7) %		1278
$K^*(892)\bar{K}^*(892)$	(7.1 \pm 1.3) $\times 10^{-3}$		1196
$K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-$	(1.1 \pm 0.5) %		1073
$\phi K^+ K^-$	(2.9 \pm 1.4) $\times 10^{-3}$		1104
$\phi\phi$	(1.79 \pm 0.20) $\times 10^{-3}$		1089
$\phi 2(\pi^+ \pi^-)$	< 4 $\times 10^{-3}$	90%	1251
$a_0(980)\pi$	< 2 %	90%	1327
$a_2(1320)\pi$	< 2 %	90%	1197
$K^*(892)\bar{K} + \text{c.c.}$	< 1.28 %	90%	1310
$f_2(1270)\eta$	< 1.1 %	90%	1145
$\omega\omega$	< 3.1 $\times 10^{-3}$	90%	1270
$\omega\phi$	< 2.5 $\times 10^{-4}$	90%	1185
$f_2(1270)f_2(1270)$	(9.8 \pm 2.5) $\times 10^{-3}$		774
$f_2(1270)f_2'(1525)$	(9.8 \pm 3.2) $\times 10^{-3}$		513
$f_0(980)\eta$	seen		1264
$f_0(1500)\eta$	seen		1025
$f_0(2200)\eta$	seen		498
$a_0(980)\pi$	seen		1327
$a_0(1320)\pi$	seen		—
$a_0(1450)\pi$	seen		1123
$a_0(1950)\pi$	seen		860
$K_0^*(1430)\bar{K}$	seen		—
$K_2^*(1430)\bar{K}$	seen		—
$K_0^*(1950)\bar{K}$	seen		—

Decays into stable hadrons

$K\bar{K}\pi$	(7.3 \pm 0.5) %	1381
$K\bar{K}\eta$	(1.36 \pm 0.16) %	1265
$\eta\pi^+\pi^-$	(1.7 \pm 0.5) %	1428

$\eta 2(\pi^+ \pi^-)$	$(4.4 \pm 1.3) \%$		1386
$K^+ K^- \pi^+ \pi^-$	$(6.9 \pm 1.1) \times 10^{-3}$		1345
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.5 \pm 0.6) \%$		1304
$K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}$	$(5.6 \pm 1.5) \%$		—
$K^+ K^- 2(\pi^+ \pi^-)$	$(7.5 \pm 2.4) \times 10^{-3}$		1254
$2(K^+ K^-)$	$(1.47 \pm 0.31) \times 10^{-3}$		1055
$\pi^+ \pi^- \pi^0$	$< 5 \times 10^{-4}$	90%	1476
$\pi^+ \pi^- \pi^0 \pi^0$	$(4.7 \pm 1.0) \%$		1460
$2(\pi^+ \pi^-)$	$(9.7 \pm 1.2) \times 10^{-3}$		1459
$2(\pi^+ \pi^- \pi^0)$	$(17.4 \pm 3.3) \%$		1409
$3(\pi^+ \pi^-)$	$(1.8 \pm 0.4) \%$		1407
$\rho \bar{\rho}$	$(1.51 \pm 0.16) \times 10^{-3}$		1160
$\rho \bar{\rho} \pi^0$	$(3.6 \pm 1.3) \times 10^{-3}$		1101
$\Lambda \bar{\Lambda}$	$(1.09 \pm 0.24) \times 10^{-3}$		991
$K^+ \bar{p} \Lambda + \text{c.c.}$	$(2.5 \pm 0.4) \times 10^{-3}$		772
$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	$(3.1 \pm 1.3) \times 10^{-3}$		693
$\Sigma^+ \bar{\Sigma}^-$	$(2.1 \pm 0.6) \times 10^{-3}$		901
$\Xi^- \bar{\Xi}^+$	$(9.0 \pm 2.6) \times 10^{-4}$		692
$\pi^+ \pi^- \rho \bar{\rho}$	$(5.3 \pm 1.8) \times 10^{-3}$		1027

Radiative decays

$\gamma \gamma$	$(1.57 \pm 0.12) \times 10^{-4}$		1492
-----------------	----------------------------------	--	------

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes

$\pi^+ \pi^-$	P, CP	$< 1.1 \times 10^{-4}$	90%	1485
$\pi^0 \pi^0$	P, CP	$< 4 \times 10^{-5}$	90%	1486
$K^+ K^-$	P, CP	$< 6 \times 10^{-4}$	90%	1408
$K_S^0 K_S^0$	P, CP	$< 3.1 \times 10^{-4}$	90%	1407

$J/\psi(1S)$

$$J^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 3096.900 \pm 0.006$ MeV

Full width $\Gamma = 92.9 \pm 2.8$ keV ($S = 1.1$)

$\Gamma_{ee} = 5.53 \pm 0.10$ keV

$\Gamma_{ee} < 5.4$ eV, CL = 90%

$J/\psi(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level (MeV/c)	p
hadrons	$(87.7 \pm 0.5) \%$		—
virtual $\gamma \rightarrow$ hadrons	$(13.50 \pm 0.30) \%$		—
ggg	$(64.1 \pm 1.0) \%$		—
γgg	$(8.8 \pm 1.1) \%$		—
$e^+ e^-$	$(5.971 \pm 0.032) \%$		1548
$e^+ e^- \gamma$	[a] $(8.8 \pm 1.4) \times 10^{-3}$		1548
$\mu^+ \mu^-$	$(5.961 \pm 0.033) \%$		1545

Decays involving hadronic resonances

$\rho\pi$	$(1.69 \pm 0.15) \%$	S=2.4	1448
$\rho^0\pi^0$	$(5.6 \pm 0.7) \times 10^{-3}$		1448
$\rho(770)^\mp K^\pm K_S^0$	$(1.9 \pm 0.4) \times 10^{-3}$		—
$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	$(2.3 \pm 0.7) \times 10^{-3}$		—
$\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.5 \pm 0.6) \times 10^{-4}$		—
$\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	$(2.0 \pm 0.5) \times 10^{-4}$		—
$\rho(1450)\eta'(958) \rightarrow$ $\pi^+\pi^-\eta'(958)$	$(3.3 \pm 0.7) \times 10^{-6}$		—
$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	$(1.7 \pm 1.1) \times 10^{-4}$		—
$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	$(8 \pm 40) \times 10^{-6}$		—
$a_2(1320)\rho$	$(1.09 \pm 0.22) \%$		1124
$\omega\pi^+\pi^+\pi^-\pi^-$	$(8.5 \pm 3.4) \times 10^{-3}$		1392
$\omega\pi^+\pi^-\pi^0$	$(4.0 \pm 0.7) \times 10^{-3}$		1418
$\omega\pi^+\pi^-$	$(7.2 \pm 1.0) \times 10^{-3}$		1435
$\omega f_2(1270)$	$(4.3 \pm 0.6) \times 10^{-3}$		1142
$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$		1266
$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.22 \\ 0.40 \end{smallmatrix}) \times 10^{-3}$		1266
$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.0 \\ 0.6 \end{smallmatrix}) \times 10^{-3}$		—
$K_S^0 \pi^- K^*(892)^+ + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$		1342
$K_S^0 \pi^- K^*(892)^+ + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$		—
$K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.6 \\ 0.5 \end{smallmatrix}) \times 10^{-6}$		—
$\eta K^*(892)^0 \bar{K}^*(892)^0$	$(1.15 \pm 0.26) \times 10^{-3}$		1003
$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$		—
$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$		1000
$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$		—
$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$		—
$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(4.9 \pm 2.8) \times 10^{-5}$		—
$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(8 \pm 6) \times 10^{-5}$		—
$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(7.5 \pm 3.5) \times 10^{-5}$		—
$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(4.0 \pm 1.0) \times 10^{-4}$		—
$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(4.66 \pm 0.31) \times 10^{-3}$		1012
$K^*(892)^+ K_2^*(1430)^- + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$		1012
$K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow$ $K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(4 \pm 4) \times 10^{-4}$		—

$K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow$	$(6.9 \pm 0.9) \times 10^{-4}$	–
$K^*(892)^0 K^- \pi^+ + \text{c.c.}$		
$\omega K^*(892) \bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	1097
$\bar{K} K^*(892) + \text{c.c.} \rightarrow$	$(5.0 \pm 0.5) \times 10^{-3}$	–
$K_S^0 K^\pm \pi^\mp$		
$K^+ K^*(892)^- + \text{c.c.}$	$(5.0 \pm 0.4) \times 10^{-3}$	1373
$K^+ K^*(892)^- + \text{c.c.} \rightarrow$	$(1.98 \pm 0.21) \times 10^{-3}$	–
$K^+ K^- \pi^0$		
$K^+ K^*(892)^- + \text{c.c.} \rightarrow$	$(3.0 \pm 0.4) \times 10^{-3}$	–
$K^0 K^\pm \pi^\mp + \text{c.c.}$		
$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	1373
$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$	$(3.2 \pm 0.4) \times 10^{-3}$	–
$K^0 K^\pm \pi^\mp + \text{c.c.}$		
$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	1170
$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(7.7 \pm 1.6) \times 10^{-3}$	1343
$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	1344
$K^*(892)^0 K_S^0 \pi^0$	$(6 \pm 4) \times 10^{-4}$	1343
$\omega \pi^0 \pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$	1436
$\omega \pi^0 \eta$	$(3.4 \pm 1.7) \times 10^{-4}$	1363
$b_1(1235)^\pm \pi^\mp$	[b] $(3.0 \pm 0.5) \times 10^{-3}$	1300
$\omega K^\pm K_S^0 \pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$	1210
$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$	1300
$\eta K^\pm K_S^0 \pi^\mp$	[b] $(2.2 \pm 0.4) \times 10^{-3}$	1278
$\phi K^*(892) \bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$	969
$\omega K \bar{K}$	$(1.9 \pm 0.4) \times 10^{-3}$	1268
$\omega f_0(1710) \rightarrow \omega K \bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$	878
$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$	1318
$\Delta(1232)^{++} \bar{p} \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	1030
$\omega \eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6 1394
$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3 1179
$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$	1176
$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$	875
$\phi K^+ K^-$	$(8.3 \pm 1.2) \times 10^{-4}$	1179
$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$	1036
$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$	938
$\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.})$	[b] $(1.16 \pm 0.05) \times 10^{-3}$	697
$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(1.07 \pm 0.08) \times 10^{-3}$	697
$K^+ K^- f_2'(1525)$	$(1.04 \pm 0.35) \times 10^{-3}$	892
$\phi f_2'(1525)$	$(8 \pm 4) \times 10^{-4}$	S=2.7 871
$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7 1365
$\phi \pi^0 \pi^0$	$(5.0 \pm 1.0) \times 10^{-4}$	1366
$\phi K^\pm K_S^0 \pi^\mp$	[b] $(7.2 \pm 0.8) \times 10^{-4}$	1114
$\omega f_1(1420)$	$(6.8 \pm 2.4) \times 10^{-4}$	1062
$\phi \eta$	$(7.4 \pm 0.8) \times 10^{-4}$	S=1.5 1320

$\Xi^0 \Xi^0$		$(1.17 \pm 0.04) \times 10^{-3}$		818
$\Xi(1530)^- \Xi^+$		$(5.9 \pm 1.5) \times 10^{-4}$		600
$p K^- \bar{\Sigma}(1385)^0$		$(5.1 \pm 3.2) \times 10^{-4}$		646
$\omega \pi^0$		$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4	1446
$\omega \pi^0 \rightarrow \pi^+ \pi^- \pi^0$		$(1.7 \pm 0.8) \times 10^{-5}$		—
$\phi \eta'(958)$		$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2	1192
$\phi f_0(980)$		$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9	1178
$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$		$(2.59 \pm 0.34) \times 10^{-4}$		—
$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$		$(1.8 \pm 0.5) \times 10^{-4}$		—
$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$		$(4.5 \pm 1.0) \times 10^{-6}$		—
$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 p^0 \pi^0$		$(1.7 \pm 0.6) \times 10^{-6}$		1045
$\eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-$		$(3.2 \pm 1.0) \times 10^{-4}$		—
$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$		$(4.4 \pm 1.4) \times 10^{-6}$		—
$\Xi(1530)^0 \Xi^0$		$(3.2 \pm 1.4) \times 10^{-4}$		608
$\Sigma(1385)^- \bar{\Sigma}^+$ (or c.c.)	[b]	$(3.1 \pm 0.5) \times 10^{-4}$		855
$\phi f_1(1285)$		$(2.6 \pm 0.5) \times 10^{-4}$		1032
$\phi f_1(1285) \rightarrow$		$(9.4 \pm 2.8) \times 10^{-7}$		952
$\phi \pi^0 f_0(980) \rightarrow$				
$\phi \pi^0 \pi^+ \pi^-$				
$\phi f_1(1285) \rightarrow$		$(2.1 \pm 2.2) \times 10^{-7}$		955
$\phi \pi^0 f_0(980) \rightarrow$				
$\phi \pi^0 \pi^0 \pi^0$				
$\eta \pi^+ \pi^-$		$(4.2 \pm 0.8) \times 10^{-4}$		1487
$\eta \rho$		$(1.93 \pm 0.23) \times 10^{-4}$		1396
$\omega \eta'(958)$		$(1.89 \pm 0.18) \times 10^{-4}$		1279
$\omega f_0(980)$		$(1.4 \pm 0.5) \times 10^{-4}$		1267
$\rho \eta'(958)$		$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6	1281
$a_2(1320)^\pm \pi^\mp$	[b]	$< 4.3 \times 10^{-3}$	CL=90%	1264
$K \bar{K}_2^*(1430) + \text{c.c.}$		$< 4.0 \times 10^{-3}$	CL=90%	1159
$K_1(1270)^\pm K^\mp$		$< 3.0 \times 10^{-3}$	CL=90%	1231
$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$		$(8.5 \pm 2.5) \times 10^{-7}$		—
$K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}$		$(3.6 \pm 1.8) \times 10^{-3}$		1117
$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$		$< 2.9 \times 10^{-3}$	CL=90%	604
$\phi \pi^0$		3×10^{-6} or 1×10^{-7}		1377
$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$		$(2.0 \pm 1.0) \times 10^{-5}$		946
$\omega f_2'(1525)$		$< 2.2 \times 10^{-4}$	CL=90%	1003
$\omega X(1835) \rightarrow \omega p \bar{p}$		$< 3.9 \times 10^{-6}$	CL=95%	—
$\phi X(1835) \rightarrow \phi p \bar{p}$		$< 2.1 \times 10^{-7}$	CL=90%	—
$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$		$< 2.8 \times 10^{-4}$	CL=90%	578
$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$		$< 6.13 \times 10^{-5}$	CL=90%	—
$\eta \phi(2170) \rightarrow \eta \phi f_0(980) \rightarrow$		$(1.2 \pm 0.4) \times 10^{-4}$		628
$\eta \phi \pi^+ \pi^-$				
$\eta \phi(2170) \rightarrow$		$< 2.52 \times 10^{-4}$	CL=90%	—
$\eta K^*(892)^0 \bar{K}^*(892)^0$				

$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	< 8.2	$\times 10^{-6}$	CL=90%	912
$\Delta(1232)^+ \bar{p}$	< 1	$\times 10^{-4}$	CL=90%	1100
$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	< 4.1	$\times 10^{-6}$	CL=90%	—
$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	< 1.80	$\times 10^{-3}$	CL=90%	806
$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	< 1.1	$\times 10^{-5}$	CL=90%	—
$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 2.1	$\times 10^{-5}$	CL=90%	—
$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 1.6	$\times 10^{-5}$	CL=90%	—
$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 5.6	$\times 10^{-5}$	CL=90%	—
$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 1.1	$\times 10^{-5}$	CL=90%	—

Decays into stable hadrons

$2(\pi^+ \pi^-) \pi^0$	$(3.37 \pm 0.26) \%$			1496
$3(\pi^+ \pi^-) \pi^0$	$(2.9 \pm 0.6) \%$			1433
$\pi^+ \pi^- \pi^0$	$(2.10 \pm 0.08) \%$		S=1.6	1533
$\pi^+ \pi^- \pi^0 \pi^0 \pi^0$	$(2.71 \pm 0.29) \%$			1497
$\rho^\pm \pi^\mp \pi^0 \pi^0$	$(1.41 \pm 0.22) \%$			1421
$\rho^+ \rho^- \pi^0$	$(6.0 \pm 1.1) \times 10^{-3}$			1298
$\pi^+ \pi^- \pi^0 K^+ K^-$	$(1.20 \pm 0.30) \%$			1368
$4(\pi^+ \pi^-) \pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$			1345
$\pi^+ \pi^- K^+ K^-$	$(6.84 \pm 0.32) \times 10^{-3}$			1407
$\pi^+ \pi^- K_S^0 K_L^0$	$(3.8 \pm 0.6) \times 10^{-3}$			1406
$\pi^+ \pi^- K_S^0 K_S^0$	$(1.68 \pm 0.19) \times 10^{-3}$			1406
$\pi^\pm \pi^0 K^\mp K_S^0$	$(5.7 \pm 0.5) \times 10^{-3}$			1408
$K^+ K^- K_S^0 K_S^0$	$(4.1 \pm 0.8) \times 10^{-4}$			1127
$\pi^+ \pi^- K^+ K^- \eta$	$(4.7 \pm 0.7) \times 10^{-3}$			1221
$\pi^0 \pi^0 K^+ K^-$	$(2.12 \pm 0.23) \times 10^{-3}$			1410
$\pi^0 \pi^0 K_S^0 K_L^0$	$(1.9 \pm 0.4) \times 10^{-3}$			1408
$K \bar{K} \pi$	$(6.1 \pm 1.0) \times 10^{-3}$			1442
$K^+ K^- \pi^0$	$(2.14 \pm 0.24) \times 10^{-3}$			1442
$K_S^0 K^\pm \pi^\mp$	$(5.6 \pm 0.5) \times 10^{-3}$			1440
$K_S^0 K_L^0 \pi^0$	$(2.06 \pm 0.27) \times 10^{-3}$			1440
$K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K_L^0 \pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$			—
$K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K_L^0 \pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$			—
$K_S^0 K_L^0 \eta$	$(1.44 \pm 0.34) \times 10^{-3}$			1328
$2(\pi^+ \pi^-)$	$(3.57 \pm 0.30) \times 10^{-3}$			1517
$3(\pi^+ \pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$			1466
$2(\pi^+ \pi^- \pi^0)$	$(1.61 \pm 0.21) \%$			1468
$2(\pi^+ \pi^-) \eta$	$(2.26 \pm 0.28) \times 10^{-3}$			1446
$3(\pi^+ \pi^-) \eta$	$(7.2 \pm 1.5) \times 10^{-4}$			1379
$\pi^+ \pi^- \pi^0 \pi^0 \eta$	$(2.3 \pm 0.5) \times 10^{-3}$			1448

$\rho^\pm \pi^\mp \pi^0 \eta$		$(1.9 \pm 0.8) \times 10^{-3}$		1326
$\rho \bar{p}$		$(2.121 \pm 0.029) \times 10^{-3}$		1232
$\rho \bar{p} \pi^0$		$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1	1176
$\rho \bar{p} \pi^+ \pi^-$		$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3	1107
$\rho \bar{p} \pi^+ \pi^- \pi^0$	[c]	$(2.3 \pm 0.9) \times 10^{-3}$	S=1.9	1033
$\rho \bar{p} \eta$		$(2.00 \pm 0.12) \times 10^{-3}$		948
$\rho \bar{p} \rho$	<	3.1×10^{-4}	CL=90%	774
$\rho \bar{p} \omega$		$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3	768
$\rho \bar{p} \eta'(958)$		$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0	596
$\rho \bar{p} a_0(980) \rightarrow \rho \bar{p} \pi^0 \eta$		$(6.8 \pm 1.8) \times 10^{-5}$		—
$\rho \bar{p} \phi$		$(5.19 \pm 0.33) \times 10^{-5}$		527
$n \bar{n}$		$(2.09 \pm 0.16) \times 10^{-3}$		1231
$n \bar{n} \pi^+ \pi^-$		$(4 \pm 4) \times 10^{-3}$		1106
$\Sigma^+ \bar{\Sigma}^-$		$(1.50 \pm 0.24) \times 10^{-3}$		992
$\Sigma^0 \bar{\Sigma}^0$		$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4	988
$2(\pi^+ \pi^-) K^+ K^-$		$(3.1 \pm 1.3) \times 10^{-3}$		1320
$\rho \bar{n} \pi^-$		$(2.12 \pm 0.09) \times 10^{-3}$		1174
$n N(1440)$	seen			978
$n N(1520)$	seen			928
$n N(1535)$	seen			917
$\Xi^- \bar{\Xi}^+$		$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4	807
$\Lambda \bar{\Lambda}$		$(1.89 \pm 0.09) \times 10^{-3}$	S=2.8	1074
$\Lambda \bar{\Sigma}^- \pi^+$ (or c.c.)	[b]	$(8.3 \pm 0.7) \times 10^{-4}$	S=1.2	950
$\rho K^- \bar{\Lambda} + \text{c.c.}$		$(8.7 \pm 1.1) \times 10^{-4}$		876
$2(K^+ K^-)$		$(7.2 \pm 0.8) \times 10^{-4}$		1131
$\rho K^- \bar{\Sigma}^0$		$(2.9 \pm 0.8) \times 10^{-4}$		819
$K^+ K^-$		$(2.86 \pm 0.21) \times 10^{-4}$		1468
$K_S^0 K_L^0$		$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4	1466
$\Lambda \bar{\Lambda} \pi^+ \pi^-$		$(4.3 \pm 1.0) \times 10^{-3}$		903
$\Lambda \bar{\Lambda} \eta$		$(1.62 \pm 0.17) \times 10^{-4}$		672
$\Lambda \bar{\Lambda} \pi^0$		$(3.8 \pm 0.4) \times 10^{-5}$		998
$\bar{\Lambda} n K_S^0 + \text{c.c.}$		$(6.5 \pm 1.1) \times 10^{-4}$		872
$\pi^+ \pi^-$		$(1.47 \pm 0.14) \times 10^{-4}$		1542
$\Lambda \bar{\Sigma} + \text{c.c.}$		$(2.83 \pm 0.23) \times 10^{-5}$		1034
$K_S^0 K_S^0$	<	1.4×10^{-8}	CL=95%	1466

Radiative decays

3γ		$(1.16 \pm 0.22) \times 10^{-5}$		1548
4γ	<	9×10^{-6}	CL=90%	1548
5γ	<	1.5×10^{-5}	CL=90%	1548
$\gamma \pi^0 \pi^0$		$(1.15 \pm 0.05) \times 10^{-3}$		1543
$\gamma \eta \pi^0$		$(2.14 \pm 0.31) \times 10^{-5}$		1497
$\gamma a_0(980)^0 \rightarrow \gamma \eta \pi^0$	<	2.5×10^{-6}	CL=95%	—
$\gamma a_2(1320)^0 \rightarrow \gamma \eta \pi^0$	<	6.6×10^{-6}	CL=95%	—

$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$		1466
$\gamma \eta_c(1S)$	$(1.7 \pm 0.4) \%$	S=1.5	111
$\gamma \eta_c(1S) \rightarrow 3\gamma$	$(3.8 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.3 \\ 1.0 \end{smallmatrix}) \times 10^{-6}$	S=1.1	—
$\gamma \pi^+ \pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$		1518
$\gamma \eta \pi \pi$	$(6.1 \pm 1.0) \times 10^{-3}$		1487
$\gamma \eta_2(1870) \rightarrow \gamma \eta \pi^+ \pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$		—
$\gamma \eta(1405/1475) \rightarrow \gamma K \bar{K} \pi$	[d] $(2.8 \pm 0.6) \times 10^{-3}$	S=1.6	1223
$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8	1223
$\gamma \eta(1405/1475) \rightarrow \gamma \eta \pi^+ \pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$		—
$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \phi$	$< 8.2 \times 10^{-5}$	CL=95%	—
$\gamma \eta(1405) \rightarrow \gamma \gamma \gamma$	$< 2.63 \times 10^{-6}$	CL=90%	—
$\gamma \eta(1475) \rightarrow \gamma \gamma \gamma$	$< 1.86 \times 10^{-6}$	CL=90%	—
$\gamma \rho \rho$	$(4.5 \pm 0.8) \times 10^{-3}$		1340
$\gamma \rho \omega$	$< 5.4 \times 10^{-4}$	CL=90%	1338
$\gamma \rho \phi$	$< 8.8 \times 10^{-5}$	CL=90%	1258
$\gamma \eta'(958)$	$(5.21 \pm 0.17) \times 10^{-3}$	S=1.4	1400
$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9	1517
$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$		878
$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$		—
$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$		1407
$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$		891
$\gamma \omega \omega$	$(1.61 \pm 0.33) \times 10^{-3}$		1336
$\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3	1223
$\gamma f_2(1270)$	$(1.64 \pm 0.12) \times 10^{-3}$	S=1.3	1286
$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.60 \\ 0.22 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$		—
$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$		—
$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.24 \\ 0.60 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(9.5 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.0 \\ 0.5 \end{smallmatrix}) \times 10^{-4}$	S=1.5	1075
$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.8 \pm 0.5) \times 10^{-4}$		—
$\gamma f_0(1710) \rightarrow \gamma \omega \omega$	$(3.1 \pm 1.0) \times 10^{-4}$		—
$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$(2.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.2 \\ 0.7 \end{smallmatrix}) \times 10^{-4}$		—
$\gamma \eta$	$(1.108 \pm 0.027) \times 10^{-3}$		1500
$\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi$	$(7.9 \pm 1.3) \times 10^{-4}$		1220
$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$		1283
$\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$		—
$\gamma f_2'(1525)$	$(5.7 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.8 \\ 0.5 \end{smallmatrix}) \times 10^{-4}$	S=1.5	1173
$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.7 \\ 0.5 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_2'(1525) \rightarrow \gamma \eta \eta$	$(3.4 \pm 1.4) \times 10^{-5}$		—

$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$(2.8 \pm 1.8) \times 10^{-4}$		—
$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$		—
$\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.20 \\ 0.33 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_0(1800) \rightarrow \gamma \omega \phi$	$(2.5 \pm 0.6) \times 10^{-4}$		—
$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	$(5.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 3.5 \\ 2.4 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_2(1950) \rightarrow$	$(7.0 \pm 2.2) \times 10^{-4}$		—
$\gamma K^*(892) \bar{K}^*(892)$			
$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$		1266
$\gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1	1166
$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$		1232
$\gamma \eta(2225)$	$(3.14 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.50 \\ 0.19 \end{smallmatrix}) \times 10^{-4}$		752
$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$		1048
$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$		—
$\gamma \eta(1760) \rightarrow \gamma \gamma \gamma$	$< 4.80 \times 10^{-6}$	CL=90%	—
$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.77 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.34 \\ 0.40 \end{smallmatrix}) \times 10^{-4}$	S=1.1	1006
$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.5 \\ 0.9 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2.0 \\ 1.3 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	$< 3.56 \times 10^{-6}$	CL=90%	—
$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.7 \\ 0.8 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma (K \bar{K} \pi) [J^{PC} = 0^{-+}]$	$(7 \pm 4) \times 10^{-4}$	S=2.1	1442
$\gamma \pi^0$	$(3.56 \pm 0.17) \times 10^{-5}$		1546
$\gamma p \bar{p} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%	1107
$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%	1074
$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.60 \\ 0.30 \end{smallmatrix}) \times 10^{-4}$		—
$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$		—
$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$		—
$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.19 \\ 0.50 \end{smallmatrix}) \times 10^{-4}$		—
$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$	CL=90%	—
$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$	CL=90%	—
$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$		—
$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$		—
$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2.4 \\ 2.2 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 4.0 \\ 1.5 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma f_0(1500) \rightarrow \gamma \pi \pi$	$(1.09 \pm 0.24) \times 10^{-4}$		1183
$\gamma f_0(1500) \rightarrow \gamma \eta \eta$	$(1.7 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.6 \\ 1.4 \end{smallmatrix}) \times 10^{-5}$		—
$\gamma A \rightarrow \gamma \text{invisible}$	$[e] < 6.3 \times 10^{-6}$	CL=90%	—
$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	$[f] < 5 \times 10^{-6}$	CL=90%	—

Dalitz decays

$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$	1546
$\eta e^+ e^-$	$(1.43 \pm 0.07) \times 10^{-5}$	1500
$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$	1400
$\eta U \rightarrow \eta e^+ e^-$	$< 9.11 \times 10^{-7}$	CL=90% —
$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	$< 2.0 \times 10^{-7}$	CL=90% —

Weak decays

$D^- e^+ \nu_e + \text{c.c.}$	$< 1.2 \times 10^{-5}$	CL=90% 984
$\overline{D}^0 e^+ e^- + \text{c.c.}$	$< 8.5 \times 10^{-8}$	CL=90% 987
$D_s^- e^+ \nu_e + \text{c.c.}$	$< 1.3 \times 10^{-6}$	CL=90% 923
$D_s^{*-} e^+ \nu_e + \text{c.c.}$	$< 1.8 \times 10^{-6}$	CL=90% 828
$D^- \pi^+ + \text{c.c.}$	$< 7.5 \times 10^{-5}$	CL=90% 977
$\overline{D}^0 \overline{K}^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90% 898
$\overline{D}^0 \overline{K}^{*0} + \text{c.c.}$	$< 2.5 \times 10^{-6}$	CL=90% 670
$D_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90% 915
$D_s^- \rho^+ + \text{c.c.}$	$< 1.3 \times 10^{-5}$	CL=90% 663

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

$\gamma\gamma$	C	$< 2.7 \times 10^{-7}$	CL=90%	1548
$\gamma\phi$	C	$< 1.4 \times 10^{-6}$	CL=90%	1381
$e^\pm \mu^\mp$	LF	$< 1.6 \times 10^{-7}$	CL=90%	1547
$e^\pm \tau^\mp$	LF	$< 8.3 \times 10^{-6}$	CL=90%	1039
$\mu^\pm \tau^\mp$	LF	$< 2.0 \times 10^{-6}$	CL=90%	1035

Other decays

invisible	$< 7 \times 10^{-4}$	CL=90%	—
-----------	----------------------	--------	---

 $\chi_{c0}(1P)$

$$J^{PC} = 0^+(0^{++})$$

 Mass $m = 3414.71 \pm 0.30$ MeV

 Full width $\Gamma = 10.8 \pm 0.6$ MeV

$\chi_{c0}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
---	--------------------------------	-----------------------------------	----------------

Hadronic decays

$2(\pi^+ \pi^-)$	$(2.34 \pm 0.18) \%$	1679
$\rho^0 \pi^+ \pi^-$	$(9.1 \pm 2.9) \times 10^{-3}$	1607
$f_0(980) f_0(980)$	$(6.6 \pm 2.1) \times 10^{-4}$	1391
$\pi^+ \pi^- \pi^0 \pi^0$	$(3.3 \pm 0.4) \%$	1680
$\rho^+ \pi^- \pi^0 + \text{c.c.}$	$(2.9 \pm 0.4) \%$	1607
$4\pi^0$	$(3.3 \pm 0.4) \times 10^{-3}$	1681
$\pi^+ \pi^- K^+ K^-$	$(1.81 \pm 0.14) \%$	1580

$K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-$	$(9.8^{+4.0}_{-2.8}) \times 10^{-4}$	–
$K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(8.0^{+2.0}_{-2.4}) \times 10^{-4}$	–
$K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(6.3 \pm 1.9) \times 10^{-3}$	–
$K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$< 2.7 \times 10^{-3}$	CL=90%
$f_0(980) f_0(980)$	$(1.6^{+1.0}_{-0.9}) \times 10^{-4}$	1391
$f_0(980) f_0(2200)$	$(7.9^{+2.0}_{-2.5}) \times 10^{-4}$	586
$f_0(1370) f_0(1370)$	$< 2.7 \times 10^{-4}$	CL=90%
$f_0(1370) f_0(1500)$	$< 1.7 \times 10^{-4}$	CL=90%
$f_0(1370) f_0(1710)$	$(6.7^{+3.5}_{-2.3}) \times 10^{-4}$	740
$f_0(1500) f_0(1370)$	$< 1.3 \times 10^{-4}$	CL=90%
$f_0(1500) f_0(1500)$	$< 5 \times 10^{-5}$	CL=90%
$f_0(1500) f_0(1710)$	$< 7 \times 10^{-5}$	CL=90%
$K^+ K^- \pi^+ \pi^- \pi^0$	$(8.6 \pm 0.9) \times 10^{-3}$	1545
$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(4.2 \pm 0.4) \times 10^{-3}$	1543
$K^+ K^- \pi^0 \pi^0$	$(5.6 \pm 0.9) \times 10^{-3}$	1582
$K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(2.49 \pm 0.33) \%$	1581
$\rho^+ K^- K^0 + \text{c.c.}$	$(1.21 \pm 0.21) \%$	1458
$K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(4.6 \pm 1.2) \times 10^{-3}$	–
$K_S^0 K_S^0 \pi^+ \pi^-$	$(5.7 \pm 1.1) \times 10^{-3}$	1579
$K^+ K^- \eta \pi^0$	$(3.0 \pm 0.7) \times 10^{-3}$	1468
$3(\pi^+ \pi^-)$	$(1.20 \pm 0.18) \%$	1633
$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(7.5 \pm 1.6) \times 10^{-3}$	1523
$K^*(892)^0 \bar{K}^*(892)^0$	$(1.7 \pm 0.6) \times 10^{-3}$	1456
$\pi \pi$	$(8.51 \pm 0.33) \times 10^{-3}$	1702
$\pi^0 \eta$	$< 1.8 \times 10^{-4}$	1661
$\pi^0 \eta'$	$< 1.1 \times 10^{-3}$	1570
$\pi^0 \eta_c$	$< 1.6 \times 10^{-3}$	CL=90%
$\eta \eta$	$(3.01 \pm 0.19) \times 10^{-3}$	1617
$\eta \eta'$	$(9.1 \pm 1.1) \times 10^{-5}$	1521
$\eta' \eta'$	$(2.17 \pm 0.12) \times 10^{-3}$	1413
$\omega \omega$	$(9.7 \pm 1.1) \times 10^{-4}$	1517
$\omega \phi$	$(1.41 \pm 0.13) \times 10^{-4}$	1447
$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$	1457
$K^+ K^-$	$(6.05 \pm 0.31) \times 10^{-3}$	1634
$K_S^0 K_S^0$	$(3.16 \pm 0.17) \times 10^{-3}$	1633
$\pi^+ \pi^- \eta$	$< 2.0 \times 10^{-4}$	CL=90%
$\pi^+ \pi^- \eta'$	$< 4 \times 10^{-4}$	CL=90%

$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	< 9	$\times 10^{-5}$	CL=90%	1610
$K^+ K^- \pi^0$	< 6	$\times 10^{-5}$	CL=90%	1611
$K^+ K^- \eta$	< 2.3	$\times 10^{-4}$	CL=90%	1512
$K^+ K^- K_S^0 K_S^0$	(1.4 ± 0.5)	$\times 10^{-3}$		1331
$K^+ K^- K^+ K^-$	(2.82 ± 0.29)	$\times 10^{-3}$		1333
$K^+ K^- \phi$	(9.7 ± 2.5)	$\times 10^{-4}$		1381
$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	(3.7 ± 0.6)	$\times 10^{-3}$		1326
$K^+ K^- \pi^0 \phi$	(1.90 ± 0.35)	$\times 10^{-3}$		1329
$\phi \pi^+ \pi^- \pi^0$	(1.18 ± 0.15)	$\times 10^{-3}$		1525
$\phi \phi$	(8.0 ± 0.7)	$\times 10^{-4}$		1370
$\rho \bar{\rho}$	(2.21 ± 0.08)	$\times 10^{-4}$		1426
$\rho \bar{\rho} \pi^0$	(7.0 ± 0.7)	$\times 10^{-4}$	S=1.3	1379
$\rho \bar{\rho} \eta$	(3.5 ± 0.4)	$\times 10^{-4}$		1187
$\rho \bar{\rho} \omega$	(5.2 ± 0.6)	$\times 10^{-4}$		1043
$\rho \bar{\rho} \phi$	(6.0 ± 1.4)	$\times 10^{-5}$		876
$\rho \bar{\rho} \pi^+ \pi^-$	(2.1 ± 0.7)	$\times 10^{-3}$	S=1.4	1320
$\rho \bar{\rho} \pi^0 \pi^0$	(1.04 ± 0.28)	$\times 10^{-3}$		1324
$\rho \bar{\rho} K^+ K^-$ (non-resonant)	(1.22 ± 0.26)	$\times 10^{-4}$		890
$\rho \bar{\rho} K_S^0 K_S^0$	< 8.8	$\times 10^{-4}$	CL=90%	884
$\rho \bar{n} \pi^-$	(1.27 ± 0.11)	$\times 10^{-3}$		1376
$\bar{\rho} n \pi^+$	(1.37 ± 0.12)	$\times 10^{-3}$		1376
$\rho \bar{n} \pi^- \pi^0$	(2.34 ± 0.21)	$\times 10^{-3}$		1321
$\bar{\rho} n \pi^+ \pi^0$	(2.21 ± 0.18)	$\times 10^{-3}$		1321
$\Lambda \bar{\Lambda}$	(3.27 ± 0.24)	$\times 10^{-4}$		1292
$\Lambda \bar{\Lambda} \pi^+ \pi^-$	(1.18 ± 0.13)	$\times 10^{-3}$		1153
$\Lambda \bar{\Lambda} \pi^+ \pi^-$ (non-resonant)	< 5	$\times 10^{-4}$	CL=90%	1153
$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	< 5	$\times 10^{-4}$	CL=90%	1083
$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	< 5	$\times 10^{-4}$	CL=90%	1083
$K^+ \bar{\rho} \Lambda + \text{c.c.}$	(1.25 ± 0.12)	$\times 10^{-3}$	S=1.3	1132
$K^+ \bar{\rho} \Lambda(1520) + \text{c.c.}$	(2.9 ± 0.7)	$\times 10^{-4}$		858
$\Lambda(1520) \bar{\Lambda}(1520)$	(3.1 ± 1.2)	$\times 10^{-4}$		779
$\Sigma^0 \bar{\Sigma}^0$	(4.68 ± 0.32)	$\times 10^{-4}$		1222
$\Sigma^+ \bar{\Sigma}^-$	(4.6 ± 0.8)	$\times 10^{-4}$	S=2.6	1225
$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	(1.6 ± 0.6)	$\times 10^{-4}$		1001
$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	(2.3 ± 0.7)	$\times 10^{-4}$		1001
$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	(1.94 ± 0.35)	$\times 10^{-4}$		873
$\Xi^0 \bar{\Xi}^0$	(3.1 ± 0.8)	$\times 10^{-4}$		1089
$\Xi^- \bar{\Xi}^+$	(4.8 ± 0.7)	$\times 10^{-4}$		1081
$\eta_c \pi^+ \pi^-$	< 7	$\times 10^{-4}$	CL=90%	307

Radiative decays

$\gamma J/\psi(1S)$	(1.40 ± 0.05)	%		303
$\gamma \rho^0$	< 9	$\times 10^{-6}$	CL=90%	1619
$\gamma \omega$	< 8	$\times 10^{-6}$	CL=90%	1618
$\gamma \phi$	< 6	$\times 10^{-6}$	CL=90%	1555

$\gamma\gamma$	$(2.04 \pm 0.09) \times 10^{-4}$	1707
$e^+ e^- J/\psi(1S)$	$(1.54 \pm 0.33) \times 10^{-4}$	303

 $\chi_{c1}(1P)$

$$J^G(J^{PC}) = 0^+(1^{++})$$

Mass $m = 3510.67 \pm 0.05$ MeV ($S = 1.2$)

Full width $\Gamma = 0.84 \pm 0.04$ MeV

$\chi_{c1}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Hadronic decays			
$3(\pi^+ \pi^-)$	$(5.8 \pm 1.4) \times 10^{-3}$	S=1.2	1683
$2(\pi^+ \pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$		1728
$\pi^+ \pi^- \pi^0 \pi^0$	$(1.19 \pm 0.15) \%$		1729
$\rho^+ \pi^- \pi^0 + \text{c.c.}$	$(1.45 \pm 0.24) \%$		1658
$\rho^0 \pi^+ \pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$		1657
$4\pi^0$	$(5.4 \pm 0.8) \times 10^{-4}$		1729
$\pi^+ \pi^- K^+ K^-$	$(4.5 \pm 1.0) \times 10^{-3}$		1632
$K^+ K^- \pi^0 \pi^0$	$(1.12 \pm 0.27) \times 10^{-3}$		1634
$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.15 \pm 0.13) \%$		1598
$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(7.5 \pm 0.8) \times 10^{-3}$		1596
$K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(8.6 \pm 1.4) \times 10^{-3}$		1632
$\rho^- K^+ \bar{K}^0 + \text{c.c.}$	$(5.0 \pm 1.2) \times 10^{-3}$		1514
$K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(2.3 \pm 0.6) \times 10^{-3}$		—
$K^+ K^- \eta \pi^0$	$(1.12 \pm 0.34) \times 10^{-3}$		1523
$\pi^+ \pi^- K_S^0 K_S^0$	$(6.9 \pm 2.9) \times 10^{-4}$		1630
$K^+ K^- \eta$	$(3.2 \pm 1.0) \times 10^{-4}$		1566
$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$(7.0 \pm 0.6) \times 10^{-3}$		1661
$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(10 \pm 4) \times 10^{-4}$		1602
$K^*(892)^+ K^- + \text{c.c.}$	$(1.4 \pm 0.6) \times 10^{-3}$		1602
$K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K^+ \pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$	CL=90%	—
$K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow$ $K_S^0 K^+ \pi^- + \text{c.c.}$	$< 2.1 \times 10^{-3}$	CL=90%	—
$K^+ K^- \pi^0$	$(1.81 \pm 0.24) \times 10^{-3}$		1662
$\eta \pi^+ \pi^-$	$(4.62 \pm 0.23) \times 10^{-3}$		1701
$a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-$	$(3.2 \pm 0.4) \times 10^{-3}$	S=2.2	—
$a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-$	$(1.76 \pm 0.24) \times 10^{-4}$		—
$a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-$	$(4.6 \pm 0.7) \times 10^{-5}$		—
$f_2(1270) \eta \rightarrow \eta \pi^+ \pi^-$	$(3.5 \pm 0.6) \times 10^{-4}$		—
$f_4(2050) \eta \rightarrow \eta \pi^+ \pi^-$	$(2.5 \pm 0.9) \times 10^{-5}$		—

$\pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow$ $\eta \pi^+ \pi^-$	< 5	$\times 10^{-5}$	CL=90%	—
$\pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow$ $\eta \pi^+ \pi^-$	< 1.5	$\times 10^{-5}$	CL=90%	—
$\pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow$ $\eta \pi^+ \pi^-$	< 8	$\times 10^{-6}$	CL=90%	—
$f_2(1270) \eta$	(6.7 ± 1.1)	$\times 10^{-4}$		1467
$\pi^+ \pi^- \eta'$	(2.2 ± 0.4)	$\times 10^{-3}$		1612
$K^+ K^- \eta'(958)$	(8.8 ± 0.9)	$\times 10^{-4}$		1461
$K_0^*(1430)^+ K^- + \text{c.c.}$	(6.4 ± 2.2)	$\times 10^{-4}$		—
$f_0(980) \eta'(958)$	(1.6 ± 1.4)	$\times 10^{-4}$		1460
$f_0(1710) \eta'(958)$	(7 ± 7)	$\times 10^{-5}$		1118
$f_2'(1525) \eta'(958)$	(9 ± 6)	$\times 10^{-5}$		1225
$\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-$	(3.5 ± 0.9)	$\times 10^{-7}$		—
$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	(3.2 ± 2.1)	$\times 10^{-3}$		1577
$K^*(892)^0 \bar{K}^*(892)^0$	(1.4 ± 0.4)	$\times 10^{-3}$		1512
$K^+ K^- K_S^0 K_S^0$	< 4	$\times 10^{-4}$	CL=90%	1390
$K^+ K^- K^+ K^-$	(5.4 ± 1.1)	$\times 10^{-4}$		1393
$K^+ K^- \phi$	(4.1 ± 1.5)	$\times 10^{-4}$		1440
$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	(3.3 ± 0.5)	$\times 10^{-3}$		1387
$K^+ K^- \pi^0 \phi$	(1.62 ± 0.30)	$\times 10^{-3}$		1390
$\phi \pi^+ \pi^- \pi^0$	(7.5 ± 1.0)	$\times 10^{-4}$		1578
$\omega \omega$	(5.7 ± 0.7)	$\times 10^{-4}$		1571
$\omega K^+ K^-$	(7.8 ± 0.9)	$\times 10^{-4}$		1513
$\omega \phi$	(2.7 ± 0.4)	$\times 10^{-5}$		1503
$\phi \phi$	(4.2 ± 0.5)	$\times 10^{-4}$		1429
$\rho \bar{\rho}$	(7.60 ± 0.34)	$\times 10^{-5}$		1484
$\rho \bar{\rho} \pi^0$	(1.55 ± 0.18)	$\times 10^{-4}$		1438
$\rho \bar{\rho} \eta$	(1.45 ± 0.25)	$\times 10^{-4}$		1254
$\rho \bar{\rho} \omega$	(2.12 ± 0.31)	$\times 10^{-4}$		1117
$\rho \bar{\rho} \phi$	< 1.7	$\times 10^{-5}$	CL=90%	962
$\rho \bar{\rho} \pi^+ \pi^-$	(5.0 ± 1.9)	$\times 10^{-4}$		1381
$\rho \bar{\rho} \pi^0 \pi^0$	< 5	$\times 10^{-4}$	CL=90%	1385
$\rho \bar{\rho} K^+ K^-$ (non-resonant)	(1.27 ± 0.22)	$\times 10^{-4}$		974
$\rho \bar{\rho} K_S^0 K_S^0$	< 4.5	$\times 10^{-4}$	CL=90%	968
$\rho \bar{n} \pi^-$	(3.8 ± 0.5)	$\times 10^{-4}$		1435
$\bar{\rho} n \pi^+$	(3.9 ± 0.5)	$\times 10^{-4}$		1435
$\rho \bar{n} \pi^- \pi^0$	(1.03 ± 0.12)	$\times 10^{-3}$		1383
$\bar{\rho} n \pi^+ \pi^0$	(1.01 ± 0.12)	$\times 10^{-3}$		1383
$\Lambda \bar{\Lambda}$	(1.14 ± 0.11)	$\times 10^{-4}$		1355
$\Lambda \bar{\Lambda} \pi^+ \pi^-$	(2.9 ± 0.5)	$\times 10^{-4}$		1223
$\Lambda \bar{\Lambda} \pi^+ \pi^-$ (non-resonant)	(2.5 ± 0.6)	$\times 10^{-4}$		1223

$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	< 1.3	$\times 10^{-4}$	CL=90%	1157
$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	< 1.3	$\times 10^{-4}$	CL=90%	1157
$K^+ \bar{p} \Lambda + \text{c.c.}$	(4.2 ± 0.4)	$\times 10^{-4}$	S=1.2	1203
$K^+ \bar{p} \Lambda(1520) + \text{c.c.}$	(1.7 ± 0.4)	$\times 10^{-4}$		950
$\Lambda(1520) \bar{\Lambda}(1520)$	< 9	$\times 10^{-5}$	CL=90%	879
$\Sigma^0 \bar{\Sigma}^0$	(4.2 ± 0.6)	$\times 10^{-5}$		1288
$\Sigma^+ \bar{\Sigma}^-$	(3.6 ± 0.7)	$\times 10^{-5}$		1291
$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	< 9	$\times 10^{-5}$	CL=90%	1081
$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	< 5	$\times 10^{-5}$	CL=90%	1081
$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	(1.35 ± 0.24)	$\times 10^{-4}$		963
$\Xi^0 \bar{\Xi}^0$	< 6	$\times 10^{-5}$	CL=90%	1163
$\Xi^- \bar{\Xi}^+$	(8.0 ± 2.1)	$\times 10^{-5}$		1155
$\pi^+ \pi^- + K^+ K^-$	< 2.1	$\times 10^{-3}$		–
$K_S^0 K_S^0$	< 6	$\times 10^{-5}$	CL=90%	1683
$\eta_c \pi^+ \pi^-$	< 3.2	$\times 10^{-3}$	CL=90%	413

Radiative decays

$\gamma J/\psi(1S)$	(34.3 ± 1.0)	%		389
$\gamma \rho^0$	(2.16 ± 0.17)	$\times 10^{-4}$		1670
$\gamma \omega$	(6.8 ± 0.8)	$\times 10^{-5}$		1668
$\gamma \phi$	(2.4 ± 0.5)	$\times 10^{-5}$		1607
$\gamma \gamma$	< 6.3	$\times 10^{-6}$	CL=90%	1755
$e^+ e^- J/\psi(1S)$	(3.65 ± 0.25)	$\times 10^{-3}$		389

$h_c(1P)$

$$J^G(J^{PC}) = 0^-(1^{+-})$$

Mass $m = 3525.38 \pm 0.11$ MeV

Full width $\Gamma = 0.7 \pm 0.4$ MeV

$h_c(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$J/\psi(1S) \pi \pi$	not seen		312
$J/\psi(1S) \pi^+ \pi^-$	< 2.3	$\times 10^{-3}$	90% 305
$\rho \bar{p}$	< 1.5	$\times 10^{-4}$	90% 1492
$\pi^+ \pi^- \pi^0$	< 2.2	$\times 10^{-3}$	1749
$2\pi^+ 2\pi^- \pi^0$	$(2.2^{+0.8}_{-0.7})$	%	1716
$3\pi^+ 3\pi^- \pi^0$	< 2.9	%	1661

Radiative decays

$\gamma \eta$	(4.7 ± 2.1)	$\times 10^{-4}$		1720
$\gamma \eta'(958)$	(1.5 ± 0.4)	$\times 10^{-3}$		1633
$\gamma \eta_c(1S)$	(51 ± 6)	%		500

$\chi_{c2}(1P)$

$$J^{PC} = 0^+(2^{++})$$

 Mass $m = 3556.17 \pm 0.07$ MeV

 Full width $\Gamma = 1.97 \pm 0.09$ MeV

$\chi_{c2}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level (MeV/c)	P
Hadronic decays			
$2(\pi^+\pi^-)$	(1.02±0.09) %		1751
$\pi^+\pi^-\pi^0\pi^0$	(1.83±0.23) %		1752
$\rho^+\pi^-\pi^0 + \text{c.c.}$	(2.19±0.34) %		1682
$4\pi^0$	(1.11±0.15) $\times 10^{-3}$		1752
$K^+K^-\pi^0\pi^0$	(2.1 ±0.4) $\times 10^{-3}$		1658
$K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(1.38±0.20) %		1657
$\rho^-K^+\bar{K}^0 + \text{c.c.}$	(4.1 ±1.2) $\times 10^{-3}$		1540
$K^*(892)^0K^-\pi^+ \rightarrow$ $K^-\pi^+K^0\pi^0 + \text{c.c.}$	(2.9 ±0.8) $\times 10^{-3}$		–
$K^*(892)^0\bar{K}^0\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(3.8 ±0.9) $\times 10^{-3}$		–
$K^*(892)^-K^+\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(3.7 ±0.8) $\times 10^{-3}$		–
$K^*(892)^+\bar{K}^0\pi^- \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(2.9 ±0.8) $\times 10^{-3}$		–
$K^+K^-\eta\pi^0$	(1.3 ±0.4) $\times 10^{-3}$		1549
$K^+K^-\pi^+\pi^-$	(8.4 ±0.9) $\times 10^{-3}$		1656
$K^+K^-\pi^+\pi^-\pi^0$	(1.17±0.13) %		1623
$K_S^0K^\pm\pi^\mp\pi^+\pi^-$	(7.3 ±0.8) $\times 10^{-3}$		1621
$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	(2.1 ±1.1) $\times 10^{-3}$		1602
$K^*(892)^0K^*(892)^0$	(2.3 ±0.4) $\times 10^{-3}$		1538
$3(\pi^+\pi^-)$	(8.6 ±1.8) $\times 10^{-3}$		1707
$\phi\phi$	(1.06±0.09) $\times 10^{-3}$		1457
$\omega\omega$	(8.4 ±1.0) $\times 10^{-4}$		1597
ωK^+K^-	(7.3 ±0.9) $\times 10^{-4}$		1540
$\omega\phi$	(9.6 ±2.7) $\times 10^{-6}$		1529
$\pi\pi$	(2.23±0.09) $\times 10^{-3}$		1773
$\rho^0\pi^+\pi^-$	(3.7 ±1.6) $\times 10^{-3}$		1682
$\pi^+\pi^-\pi^0$ (non-resonant)	(2.0 ±0.4) $\times 10^{-5}$		1765
$\rho(770)^\pm\pi^\mp$	(6 ±4) $\times 10^{-6}$		–
$\pi^+\pi^-\eta$	(4.8 ±1.3) $\times 10^{-4}$		1724
$\pi^+\pi^-\eta'$	(5.0 ±1.8) $\times 10^{-4}$		1636
$\eta\eta$	(5.4 ±0.4) $\times 10^{-4}$		1692
K^+K^-	(1.01±0.06) $\times 10^{-3}$		1708
$K_S^0K_S^0$	(5.2 ±0.4) $\times 10^{-4}$		1707
$K^*(892)^\pm K^\mp$	(1.44±0.21) $\times 10^{-4}$		1627

$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.24 \pm 0.27) \times 10^{-4}$		1627
$K_2^*(1430)^\pm K^\mp$	$(1.48 \pm 0.12) \times 10^{-3}$		—
$K_2^*(1430)^0 \bar{K}^0 + \text{c.c.}$	$(1.24 \pm 0.17) \times 10^{-3}$		1444
$K_3^*(1780)^\pm K^\mp$	$(5.2 \pm 0.8) \times 10^{-4}$		—
$K_3^*(1780)^0 \bar{K}^0 + \text{c.c.}$	$(5.6 \pm 2.1) \times 10^{-4}$		1276
$a_2(1320)^0 \pi^0$	$(1.29 \pm 0.34) \times 10^{-3}$		—
$a_2(1320)^\pm \pi^\mp$	$(1.8 \pm 0.6) \times 10^{-3}$		1531
$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$(1.28 \pm 0.18) \times 10^{-3}$		1685
$K^+ K^- \pi^0$	$(3.0 \pm 0.8) \times 10^{-4}$		1686
$K^+ K^- \eta$	$< 3.2 \times 10^{-4}$	90%	1592
$K^+ K^- \eta'(958)$	$(1.94 \pm 0.34) \times 10^{-4}$		1488
$\eta \eta'$	$(2.2 \pm 0.5) \times 10^{-5}$		1600
$\eta' \eta'$	$(4.6 \pm 0.6) \times 10^{-5}$		1498
$\pi^+ \pi^- K_S^0 K_S^0$	$(2.2 \pm 0.5) \times 10^{-3}$		1655
$K^+ K^- K_S^0 K_S^0$	$< 4 \times 10^{-4}$	90%	1418
$K^+ K^- K^+ K^-$	$(1.65 \pm 0.20) \times 10^{-3}$		1421
$K^+ K^- \phi$	$(1.42 \pm 0.29) \times 10^{-3}$		1468
$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(4.8 \pm 0.7) \times 10^{-3}$		1416
$K^+ K^- \pi^0 \phi$	$(2.7 \pm 0.5) \times 10^{-3}$		1419
$\phi \pi^+ \pi^- \pi^0$	$(9.3 \pm 1.2) \times 10^{-4}$		1603
$\rho \bar{\rho}$	$(7.33 \pm 0.33) \times 10^{-5}$		1510
$\rho \bar{\rho} \pi^0$	$(4.7 \pm 0.4) \times 10^{-4}$		1465
$\rho \bar{\rho} \eta$	$(1.74 \pm 0.25) \times 10^{-4}$		1285
$\rho \bar{\rho} \omega$	$(3.6 \pm 0.4) \times 10^{-4}$		1152
$\rho \bar{\rho} \phi$	$(2.8 \pm 0.9) \times 10^{-5}$		1002
$\rho \bar{\rho} \pi^+ \pi^-$	$(1.32 \pm 0.34) \times 10^{-3}$		1410
$\rho \bar{\rho} \pi^0 \pi^0$	$(7.8 \pm 2.3) \times 10^{-4}$		1414
$\rho \bar{\rho} K^+ K^-$ (non-resonant)	$(1.91 \pm 0.32) \times 10^{-4}$		1013
$\rho \bar{\rho} K_S^0 K_S^0$	$< 7.9 \times 10^{-4}$	90%	1007
$\rho \bar{n} \pi^-$	$(8.5 \pm 0.9) \times 10^{-4}$		1463
$\bar{\rho} n \pi^+$	$(8.9 \pm 0.8) \times 10^{-4}$		1463
$\rho \bar{n} \pi^- \pi^0$	$(2.17 \pm 0.18) \times 10^{-3}$		1411
$\bar{\rho} n \pi^+ \pi^0$	$(2.11 \pm 0.18) \times 10^{-3}$		1411
$\Lambda \bar{\Lambda}$	$(1.84 \pm 0.15) \times 10^{-4}$		1384
$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(1.25 \pm 0.15) \times 10^{-3}$		1255
$\Lambda \bar{\Lambda} \pi^+ \pi^-$ (non-resonant)	$(6.6 \pm 1.5) \times 10^{-4}$		1255
$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	$< 4 \times 10^{-4}$	90%	1192
$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	$< 6 \times 10^{-4}$	90%	1192
$K^+ \bar{\rho} \Lambda + \text{c.c.}$	$(7.8 \pm 0.5) \times 10^{-4}$		1236
$K^+ \bar{\rho} \Lambda(1520) + \text{c.c.}$	$(2.8 \pm 0.7) \times 10^{-4}$		992
$\Lambda(1520) \bar{\Lambda}(1520)$	$(4.6 \pm 1.5) \times 10^{-4}$		923
$\Sigma^0 \bar{\Sigma}^0$	$(3.7 \pm 0.6) \times 10^{-5}$		1319
$\Sigma^+ \bar{\Sigma}^-$	$(3.4 \pm 0.7) \times 10^{-5}$		1322
$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$< 1.6 \times 10^{-4}$	90%	1118

$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	< 8	$\times 10^{-5}$	90%	1118
$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	(1.76 ± 0.32)	$\times 10^{-4}$		1004
$\Xi^0 \bar{\Xi}^0$	< 1.0	$\times 10^{-4}$	90%	1197
$\Xi^- \bar{\Xi}^+$	(1.42 ± 0.32)	$\times 10^{-4}$		1189
$J/\psi(1S) \pi^+ \pi^- \pi^0$	< 1.5	%	90%	185
$\pi^0 \eta_c$	< 3.2	$\times 10^{-3}$	90%	511
$\eta_c(1S) \pi^+ \pi^-$	< 5.4	$\times 10^{-3}$	90%	459

Radiative decays

$\gamma J/\psi(1S)$	(19.0 ± 0.5)	%		430
$\gamma \rho^0$	< 1.9	$\times 10^{-5}$	90%	1694
$\gamma \omega$	< 6	$\times 10^{-6}$	90%	1692
$\gamma \phi$	< 7	$\times 10^{-6}$	90%	1632
$\gamma \gamma$	(2.85 ± 0.10)	$\times 10^{-4}$		1778
$e^+ e^- J/\psi(1S)$	(2.37 ± 0.16)	$\times 10^{-3}$		430

$\eta_c(2S)$

$$J^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.

$$\text{Mass } m = 3637.5 \pm 1.1 \text{ MeV} \quad (S = 1.2)$$

$$\text{Full width } \Gamma = 11.3_{-2.9}^{+3.2} \text{ MeV}$$

$\eta_c(2S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
hadrons	not seen		—
$K \bar{K} \pi$	(1.9 ± 1.2) %		1729
$K \bar{K} \eta$	(5 ± 4)	$\times 10^{-3}$	1637
$2\pi^+ 2\pi^-$	not seen		1792
$\rho^0 \rho^0$	not seen		1645
$3\pi^+ 3\pi^-$	not seen		1749
$K^+ K^- \pi^+ \pi^-$	not seen		1700
$K^{*0} \bar{K}^{*0}$	not seen		1585
$K^+ K^- \pi^+ \pi^- \pi^0$	(1.4 ± 1.0) %		1667
$K^+ K^- 2\pi^+ 2\pi^-$	not seen		1627
$K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	seen		1666
$2K^+ 2K^-$	not seen		1470
$\phi \phi$	not seen		1506
$p \bar{p}$	seen		1558
$\gamma \gamma$	(1.9 ± 1.3)	$\times 10^{-4}$	1819
$\gamma J/\psi(1S)$	< 1.4	%	90% 500
$\pi^+ \pi^- \eta$	not seen		1766
$\pi^+ \pi^- \eta'$	not seen		1680
$\pi^+ \pi^- \eta_c(1S)$	< 25	%	90% 537

$\psi(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 Mass $m = 3686.097 \pm 0.025$ MeV ($S = 2.6$)

 Full width $\Gamma = 294 \pm 8$ keV

 $\Gamma_{ee} = 2.33 \pm 0.04$ keV

$\psi(2S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
hadrons	(97.85 \pm 0.13) %		—
virtual $\gamma \rightarrow$ hadrons	(1.73 \pm 0.14) %	S=1.5	—
ggg	(10.6 \pm 1.6) %		—
γgg	(1.03 \pm 0.29) %		—
light hadrons	(15.4 \pm 1.5) %		—
e^+e^-	(7.93 \pm 0.17) $\times 10^{-3}$		1843
$\mu^+\mu^-$	(8.0 \pm 0.6) $\times 10^{-3}$		1840
$\tau^+\tau^-$	(3.1 \pm 0.4) $\times 10^{-3}$		489

Decays into $J/\psi(1S)$ and anything

$J/\psi(1S)$ anything	(61.4 \pm 0.6) %		—
$J/\psi(1S)$ neutrals	(25.38 \pm 0.32) %		—
$J/\psi(1S)\pi^+\pi^-$	(34.68 \pm 0.30) %		477
$J/\psi(1S)\pi^0\pi^0$	(18.24 \pm 0.31) %		481
$J/\psi(1S)\eta$	(3.37 \pm 0.05) %		199
$J/\psi(1S)\pi^0$	(1.268 \pm 0.032) $\times 10^{-3}$		528

Hadronic decays

$\pi^0 h_c(1P)$	(8.6 \pm 1.3) $\times 10^{-4}$		85
$3(\pi^+\pi^-)\pi^0$	(3.5 \pm 1.6) $\times 10^{-3}$		1746
$2(\pi^+\pi^-)\pi^0$	(2.9 \pm 1.0) $\times 10^{-3}$	S=4.7	1799
$\rho a_2(1320)$	(2.6 \pm 0.9) $\times 10^{-4}$		1501
$\pi^+\pi^-\pi^0\pi^0\pi^0$	(5.3 \pm 0.9) $\times 10^{-3}$		1800
$\rho^\pm\pi^\mp\pi^0\pi^0$	< 2.7 $\times 10^{-3}$	CL=90%	1737
$p\bar{p}$	(2.94 \pm 0.08) $\times 10^{-4}$		1586
$n\bar{n}$	(3.06 \pm 0.15) $\times 10^{-4}$		1586
$\Delta^{++}\bar{\Delta}^{--}$	(1.28 \pm 0.35) $\times 10^{-4}$		1371
$\Lambda\bar{\Lambda}\pi^0$	< 2.9 $\times 10^{-6}$	CL=90%	1412
$\Lambda\bar{\Lambda}\eta$	(2.5 \pm 0.4) $\times 10^{-5}$		1197
$\Lambda\bar{p}K^+$	(1.00 \pm 0.14) $\times 10^{-4}$		1327
$\Lambda\bar{p}K^+\pi^+\pi^-$	(1.8 \pm 0.4) $\times 10^{-4}$		1167
$\Lambda\bar{\Lambda}\pi^+\pi^-$	(2.8 \pm 0.6) $\times 10^{-4}$		1346
$\Lambda\bar{\Lambda}$	(3.81 \pm 0.13) $\times 10^{-4}$	S=1.4	1467
$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	(1.40 \pm 0.13) $\times 10^{-4}$		1376
$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	(1.54 \pm 0.14) $\times 10^{-4}$		1379
$\Lambda\bar{\Sigma}^0$	(1.23 \pm 0.24) $\times 10^{-5}$		1437
$\Sigma^0\bar{p}K^+ + \text{c.c.}$	(1.67 \pm 0.18) $\times 10^{-5}$		1291

$\Sigma^+ \bar{\Sigma}^-$	$(2.32 \pm 0.12) \times 10^{-4}$		1408
$\Sigma^0 \bar{\Sigma}^0$	$(2.35 \pm 0.09) \times 10^{-4}$	S=1.1	1405
$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$(8.5 \pm 0.7) \times 10^{-5}$		1218
$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$(8.5 \pm 0.8) \times 10^{-5}$		1218
$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(6.9 \pm 0.7) \times 10^{-5}$		1218
$\Xi^- \bar{\Xi}^+$	$(2.87 \pm 0.11) \times 10^{-4}$	S=1.1	1284
$\Xi^0 \bar{\Xi}^0$	$(2.3 \pm 0.4) \times 10^{-4}$	S=4.2	1291
$\Xi(1530)^0 \bar{\Xi}(1530)^0$	$(5.2 \begin{smallmatrix} +3.2 \\ -1.2 \end{smallmatrix}) \times 10^{-5}$		1025
$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(3.9 \pm 0.4) \times 10^{-5}$		1114
$\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$	$(5.2 \pm 1.6) \times 10^{-6}$		–
c.c.			
$\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$	$(1.20 \pm 0.32) \times 10^{-5}$		–
$K^- \Sigma^0 \bar{\Xi}^+ + \text{c.c.}$	$(3.7 \pm 0.4) \times 10^{-5}$		1060
$\Omega^- \bar{\Omega}^+$	$(5.2 \pm 0.4) \times 10^{-5}$		774
$\pi^0 p \bar{p}$	$(1.53 \pm 0.07) \times 10^{-4}$		1543
$N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(6.4 \begin{smallmatrix} +1.8 \\ -1.3 \end{smallmatrix}) \times 10^{-5}$		–
$N(1440) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(7.3 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}) \times 10^{-5}$	S=2.5	–
$N(1520) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(6.4 \begin{smallmatrix} +2.3 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$		–
$N(1535) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(2.5 \pm 1.0) \times 10^{-5}$		–
$N(1650) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(3.8 \begin{smallmatrix} +1.4 \\ -1.7 \end{smallmatrix}) \times 10^{-5}$		–
$N(1720) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(1.79 \begin{smallmatrix} +0.26 \\ -0.70 \end{smallmatrix}) \times 10^{-5}$		–
$N(2300) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(2.6 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-5}$		–
$N(2570) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(2.13 \begin{smallmatrix} +0.40 \\ -0.31 \end{smallmatrix}) \times 10^{-5}$		–
$\pi^0 f_0(2100) \rightarrow \pi^0 p \bar{p}$	$(1.1 \pm 0.4) \times 10^{-5}$		–
$\eta p \bar{p}$	$(6.0 \pm 0.4) \times 10^{-5}$		1373
$\eta f_0(2100) \rightarrow \eta p \bar{p}$	$(1.2 \pm 0.4) \times 10^{-5}$		–
$N(1535) \bar{p} \rightarrow \eta p \bar{p}$	$(4.4 \pm 0.7) \times 10^{-5}$		–
$\omega p \bar{p}$	$(6.9 \pm 2.1) \times 10^{-5}$		1247
$\eta' p \bar{p}$	$(1.10 \pm 0.13) \times 10^{-5}$		1141
$\phi p \bar{p}$	$< 2.4 \times 10^{-5}$	CL=90%	1109
$\pi^+ \pi^- p \bar{p}$	$(6.0 \pm 0.4) \times 10^{-4}$		1491
$p \bar{n} \pi^- \text{ or c.c.}$	$(2.48 \pm 0.17) \times 10^{-4}$		–
$p \bar{n} \pi^- \pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$		1492
$2(\pi^+ \pi^- \pi^0)$	$(4.8 \pm 1.5) \times 10^{-3}$		1776
$\eta \pi^+ \pi^-$	$< 1.6 \times 10^{-4}$	CL=90%	1791
$\eta \pi^+ \pi^- \pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$		1778
$2(\pi^+ \pi^-) \eta$	$(1.2 \pm 0.6) \times 10^{-3}$		1758
$\pi^+ \pi^- \pi^0 \pi^0 \eta$	$< 4 \times 10^{-4}$	CL=90%	1760
$\eta' \pi^+ \pi^- \pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$		1692

$\omega \pi^+ \pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1	1748
$b_1^\pm \pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1	1635
$b_1^0 \pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$		–
$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$		1515
$\omega \pi^0 \pi^0$	$(1.11 \pm 0.35) \times 10^{-3}$		1749
$\pi^0 \pi^0 K^+ K^-$	$(2.6 \pm 1.3) \times 10^{-4}$		1728
$\pi^+ \pi^- K^+ K^-$	$(7.3 \pm 0.5) \times 10^{-4}$		1726
$\pi^0 \pi^0 K_S^0 K_L^0$	$(1.3 \pm 0.6) \times 10^{-3}$		1726
$\rho^0 K^+ K^-$	$(2.2 \pm 0.4) \times 10^{-4}$		1616
$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$		1418
$K^+ K^- \pi^+ \pi^- \eta$	$(1.3 \pm 0.7) \times 10^{-3}$		1574
$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$		1611
$K^+ K^- 2(\pi^+ \pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$		1654
$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$		1581
$K_S^0 K_S^0 \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$		1724
$\rho^0 p \bar{p}$	$(5.0 \pm 2.2) \times 10^{-5}$		1252
$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$		1674
$2(\pi^+ \pi^-)$	$(2.4 \pm 0.6) \times 10^{-4}$	S=2.2	1817
$\rho^0 \pi^+ \pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	S=1.4	1750
$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$		1694
$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$(5.9 \pm 2.2) \times 10^{-5}$		–
$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$		–
$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$		–
$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$		–
$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$		–
$\eta K^+ K^-$, no $\eta \phi$	$(3.1 \pm 0.4) \times 10^{-5}$		1664
$\omega K^+ K^-$	$(1.62 \pm 0.11) \times 10^{-4}$	S=1.1	1614
$\omega K^*(892)^+ K^- + \text{c.c.}$	$(2.07 \pm 0.26) \times 10^{-4}$		1482
$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$(6.1 \pm 1.2) \times 10^{-5}$		1253
$\omega \bar{K}^*(892)^0 K^0$	$(1.68 \pm 0.30) \times 10^{-4}$		1481
$\omega \bar{K}_2^*(1430)^0 K^0$	$(5.8 \pm 2.2) \times 10^{-5}$		1251
$\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$		–
$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	$(1.09 \pm 0.26) \times 10^{-5}$		–
$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(3.0 \pm 1.0) \times 10^{-6}$		–
$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	$(1.2 \pm 0.7) \times 10^{-6}$		–
$3(\pi^+ \pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8	1774
$\rho \bar{p} \pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$		1435
$K^+ K^-$	$(7.5 \pm 0.5) \times 10^{-5}$		1776
$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$		1775
$\pi^+ \pi^- \pi^0$	$(2.01 \pm 0.17) \times 10^{-4}$	S=1.7	1830
$\rho(2150) \pi \rightarrow \pi^+ \pi^- \pi^0$	$(1.9 \pm 1.2) \times 10^{-4}$		–

$\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$	$(3.2 \pm 1.2) \times 10^{-5}$	S=1.8	—
$\pi^+\pi^-$	$(7.8 \pm 2.6) \times 10^{-6}$		1838
$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%	1532
$K_2^*(1430)^\pm K^\mp$	$(7.1 \pm_{-0.9}^{+1.3}) \times 10^{-5}$		—
$K^+K^-\pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$		1754
$K_S^0 K_L^0 \pi^0$	$< 3.0 \times 10^{-4}$	CL=90%	1753
$K_S^0 K_L^0 \eta$	$(1.3 \pm 0.5) \times 10^{-3}$		1661
$K^+K^*(892)^- + \text{c.c.}$	$(2.9 \pm 0.4) \times 10^{-5}$	S=1.2	1698
$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$		1697
$\phi\pi^+\pi^-$	$(1.18 \pm 0.26) \times 10^{-4}$	S=1.5	1690
$\phi f_0(980) \rightarrow \pi^+\pi^-$	$(7.5 \pm 3.3) \times 10^{-5}$	S=1.6	—
$2(K^+K^-)$	$(6.3 \pm 1.3) \times 10^{-5}$		1499
ϕK^+K^-	$(7.0 \pm 1.6) \times 10^{-5}$		1546
$2(K^+K^-)\pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$		1440
$\phi\eta$	$(3.10 \pm 0.31) \times 10^{-5}$		1654
$\eta\phi(2170), \phi(2170) \rightarrow$	$< 2.2 \times 10^{-6}$	CL=90%	—
$\phi f_0(980), f_0 \rightarrow \pi^+\pi^-$			
$\phi\eta'$	$(3.1 \pm 1.6) \times 10^{-5}$		1555
$\omega\eta'$	$(3.2 \pm_{-2.1}^{+2.5}) \times 10^{-5}$		1623
$\omega\pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$		1757
$\rho\eta'$	$(1.9 \pm_{-1.2}^{+1.7}) \times 10^{-5}$		1625
$\rho\eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1	1717
$\omega\eta$	$< 1.1 \times 10^{-5}$	CL=90%	1715
$\phi\pi^0$	$< 4 \times 10^{-7}$	CL=90%	1699
$\eta_c\pi^+\pi^-\pi^0$	$< 1.0 \times 10^{-3}$	CL=90%	512
$p\bar{p}K^+K^-$	$(2.7 \pm 0.7) \times 10^{-5}$		1118
$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$		1324
$\phi f_2'(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$		1321
$\Theta(1540)\bar{\Theta}(1540) \rightarrow$	$< 8.8 \times 10^{-6}$	CL=90%	—
$K_S^0 p K^- \bar{n} + \text{c.c.}$			
$\Theta(1540)K^-\bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 1.0 \times 10^{-5}$	CL=90%	—
$\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 7.0 \times 10^{-6}$	CL=90%	—
$\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 2.6 \times 10^{-5}$	CL=90%	—
$\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 6.0 \times 10^{-6}$	CL=90%	—
$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$		1775
$\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.}$	$< 1.7 \times 10^{-6}$	CL=90%	830

Radiative decays

$\gamma\chi_{c0}(1P)$	$(9.79 \pm 0.20) \%$	261
$\gamma\chi_{c1}(1P)$	$(9.75 \pm 0.24) \%$	171
$\gamma\chi_{c2}(1P)$	$(9.52 \pm 0.20) \%$	128
$\gamma\eta_c(1S)$	$(3.4 \pm 0.5) \times 10^{-3}$	S=1.3 635

$\gamma\eta_c(2S)$	$(7 \pm 5) \times 10^{-4}$		48
$\gamma\pi^0$	$(1.04 \pm 0.22) \times 10^{-6}$	S=1.4	1841
$\gamma\eta'(958)$	$(1.24 \pm 0.04) \times 10^{-4}$		1719
$\gamma f_2(1270)$	$(2.73 \begin{smallmatrix} +0.29 \\ -0.25 \end{smallmatrix}) \times 10^{-4}$	S=1.8	1622
$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(3.1 \pm 1.7) \times 10^{-5}$		1588
$\gamma f_0(1500)$	$(9.3 \pm 1.9) \times 10^{-5}$		1535
$\gamma f_2'(1525)$	$(3.3 \pm 0.8) \times 10^{-5}$		1528
$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.5 \pm 0.6) \times 10^{-5}$		–
$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(6.6 \pm 0.7) \times 10^{-5}$		–
$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(4.8 \pm 1.0) \times 10^{-6}$		1244
$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(3.2 \pm 1.0) \times 10^{-6}$		1193
$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 5.8 \times 10^{-6}$	CL=90%	1168
$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 9.5 \times 10^{-6}$	CL=90%	1168
$\gamma\gamma$	$< 1.5 \times 10^{-4}$	CL=90%	1843
$\gamma\eta$	$(9.2 \pm 1.8) \times 10^{-7}$		1802
$\gamma\eta\pi^+\pi^-$	$(8.7 \pm 2.1) \times 10^{-4}$		1791
$\gamma\eta(1405) \rightarrow \gamma K \bar{K} \pi$	$< 9 \times 10^{-5}$	CL=90%	1569
$\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$		–
$\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow$ $\gamma\pi^+\pi^-\pi^0$	$< 5.0 \times 10^{-7}$	CL=90%	–
$\gamma\eta(1475) \rightarrow K \bar{K} \pi$	$< 1.4 \times 10^{-4}$	CL=90%	–
$\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-$	$< 8.8 \times 10^{-5}$	CL=90%	–
$\gamma 2(\pi^+\pi^-)$	$(4.0 \pm 0.6) \times 10^{-4}$		1817
$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	$(3.7 \pm 0.9) \times 10^{-4}$		1674
$\gamma K^{*0} \bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$		1613
$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	$(2.6 \pm 0.5) \times 10^{-4}$		1753
$\gamma K^+ K^- \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$		1726
$\gamma p \bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0	1586
$\gamma f_2(1950) \rightarrow \gamma p \bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$		–
$\gamma f_2(2150) \rightarrow \gamma p \bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$		–
$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(4.6 \begin{smallmatrix} +1.8 \\ -4.0 \end{smallmatrix}) \times 10^{-6}$		–
$\gamma X \rightarrow \gamma p \bar{p}$	$[g] < 2 \times 10^{-6}$	CL=90%	–
$\gamma\pi^+\pi^- p \bar{p}$	$(2.8 \pm 1.4) \times 10^{-5}$		1491
$\gamma 2(\pi^+\pi^-) K^+ K^-$	$< 2.2 \times 10^{-4}$	CL=90%	1654
$\gamma 3(\pi^+\pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%	1774
$\gamma K^+ K^- K^+ K^-$	$< 4 \times 10^{-5}$	CL=90%	1499
$\gamma\gamma J/\psi$	$(3.1 \begin{smallmatrix} +1.0 \\ -1.2 \end{smallmatrix}) \times 10^{-4}$		542
$e^+ e^- \eta'$	$(1.90 \pm 0.26) \times 10^{-6}$		1719
$e^+ e^- \chi_{c0}(1P)$	$(1.06 \pm 0.24) \times 10^{-3}$		261
$e^+ e^- \chi_{c1}(1P)$	$(8.5 \pm 0.6) \times 10^{-4}$		171
$e^+ e^- \chi_{c2}(1P)$	$(7.0 \pm 0.8) \times 10^{-4}$		128

Weak decays

$$D^0 e^+ e^- + \text{c.c.} \quad < 1.4 \quad \times 10^{-7} \quad \text{CL}=90\% \quad 1371$$

Other decays

$$\text{invisible} \quad < 1.6 \quad \% \quad \text{CL}=90\% \quad -$$

 $\psi(3770)$

$$J^G(J^{PC}) = 0^-(1^--)$$

$$\text{Mass } m = 3773.13 \pm 0.35 \text{ MeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma = 27.2 \pm 1.0 \text{ MeV}$$

$$\Gamma_{ee} = 0.262 \pm 0.018 \text{ keV} \quad (S = 1.4)$$

In addition to the dominant decay mode to $D\bar{D}$, $\psi(3770)$ was found to decay into the final states containing the J/ψ (BAI 05, ADAM 06). ADAMS 06 and HUANG 06A searched for various decay modes with light hadrons and found a statistically significant signal for the decay to $\phi\eta$ only (ADAMS 06).

$\psi(3770)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$D\bar{D}$	(93 $^{+8}_{-9}$) %	S=2.0	286
$D^0\bar{D}^0$	(52 $^{+4}_{-5}$) %	S=2.0	286
D^+D^-	(41 ± 4) %	S=2.0	252
$J/\psi\pi^+\pi^-$	(1.93 ± 0.28) $\times 10^{-3}$		560
$J/\psi\pi^0\pi^0$	(8.0 ± 3.0) $\times 10^{-4}$		564
$J/\psi\eta$	(9 ± 4) $\times 10^{-4}$		360
$J/\psi\pi^0$	< 2.8 $\times 10^{-4}$	CL=90%	603
e^+e^-	(9.6 ± 0.7) $\times 10^{-6}$	S=1.3	1887

Decays to light hadrons

$b_1(1235)\pi$	< 1.4	$\times 10^{-5}$	CL=90%	1683
$\phi\eta'$	< 7	$\times 10^{-4}$	CL=90%	1607
$\omega\eta'$	< 4	$\times 10^{-4}$	CL=90%	1672
$\rho^0\eta'$	< 6	$\times 10^{-4}$	CL=90%	1674
$\phi\eta$	(3.1 ± 0.7) $\times 10^{-4}$			1703
$\omega\eta$	< 1.4	$\times 10^{-5}$	CL=90%	1762
$\rho^0\eta$	< 5	$\times 10^{-4}$	CL=90%	1764
$\phi\pi^0$	< 3	$\times 10^{-5}$	CL=90%	1746
$\omega\pi^0$	< 6	$\times 10^{-4}$	CL=90%	1803
$\pi^+\pi^-\pi^0$	< 5	$\times 10^{-6}$	CL=90%	1874
$\rho\pi$	< 5	$\times 10^{-6}$	CL=90%	1804
$K^*(892)^+K^- + \text{c.c.}$	< 1.4	$\times 10^{-5}$	CL=90%	1745
$K^*(892)^0\bar{K}^0 + \text{c.c.}$	< 1.2	$\times 10^{-3}$	CL=90%	1744
$K_S^0K_L^0$	< 1.2	$\times 10^{-5}$	CL=90%	1820
$2(\pi^+\pi^-)$	< 1.12	$\times 10^{-3}$	CL=90%	1861

$2(\pi^+\pi^-\pi^0)$	< 1.06	$\times 10^{-3}$	CL=90%	1843
$2(\pi^+\pi^-\pi^0)$	< 5.85	%	CL=90%	1821
$\omega\pi^+\pi^-$	< 6.0	$\times 10^{-4}$	CL=90%	1794
$3(\pi^+\pi^-)$	< 9.1	$\times 10^{-3}$	CL=90%	1819
$3(\pi^+\pi^-)\pi^0$	< 1.37	%	CL=90%	1792
$3(\pi^+\pi^-)2\pi^0$	< 11.74	%	CL=90%	1760
$\eta\pi^+\pi^-$	< 1.24	$\times 10^{-3}$	CL=90%	1836
$\pi^+\pi^-2\pi^0$	< 8.9	$\times 10^{-3}$	CL=90%	1862
$\rho^0\pi^+\pi^-$	< 6.9	$\times 10^{-3}$	CL=90%	1796
$\eta3\pi$	< 1.34	$\times 10^{-3}$	CL=90%	1824
$\eta2(\pi^+\pi^-)$	< 2.43	%	CL=90%	1804
$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%	1708
$\eta'3\pi$	< 2.44	$\times 10^{-3}$	CL=90%	1740
$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%	1772
$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%	1737
$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%	1774
$4(\pi^+\pi^-)$	< 1.67	%	CL=90%	1757
$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%	1720
$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%	1597
$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%	1741
$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%	1624
$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%	1622
ωK^+K^-	< 3.4	$\times 10^{-4}$	CL=90%	1664
$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%	1722
$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%	1693
$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%	1692
$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%	1705
$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%	1702
$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%	1660
ηK^+K^-	< 4.1	$\times 10^{-4}$	CL=90%	1712
$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%	1624
$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%	1665
$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%	1552
ϕK^+K^-	< 7.5	$\times 10^{-4}$	CL=90%	1598
$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%	1493
$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%	1425
$K_S^0 K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%	1799
$K_S^0 K^-\pi^+\pi^0$	< 1.33	%	CL=90%	1773
$K_S^0 K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%	1664
$K_S^0 K^-\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%	1739
$K_S^0 K^-\pi^+\rho^0$	< 1.6	%	CL=90%	1621
$K_S^0 K^-\pi^+\eta$	< 1.3	%	CL=90%	1669
$K_S^0 K^-\pi^+\pi^-\pi^0$	< 4.18	%	CL=90%	1703
$K_S^0 K^-\pi^+\pi^-\eta$	< 4.8	%	CL=90%	1570

$K_S^0 K^- \pi^+ 2(\pi^+ \pi^-)$	< 1.22	%	CL=90%	1658
$K_S^0 K^- \pi^+ 2\pi^0$	< 2.65	%	CL=90%	1742
$K_S^0 K^- K^+ K^- \pi^+$	< 4.9	$\times 10^{-3}$	CL=90%	1490
$K_S^0 K^- K^+ K^- \pi^+ \pi^0$	< 3.0	%	CL=90%	1427
$K_S^0 K^- K^+ K^- \pi^+ \eta$	< 2.2	%	CL=90%	1214
$K^{*0} K^- \pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%	1722
$\rho \bar{\rho} \pi^0$	< 4	$\times 10^{-5}$	CL=90%	1595
$\rho \bar{\rho} \pi^+ \pi^-$	< 5.8	$\times 10^{-4}$	CL=90%	1544
$\Lambda \bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%	1521
$\rho \bar{\rho} \pi^+ \pi^- \pi^0$	< 1.85	$\times 10^{-3}$	CL=90%	1490
$\omega \rho \bar{\rho}$	< 2.9	$\times 10^{-4}$	CL=90%	1309
$\Lambda \bar{\Lambda} \pi^0$	< 7	$\times 10^{-5}$	CL=90%	1468
$\rho \bar{\rho} 2(\pi^+ \pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%	1425
$\eta \rho \bar{\rho}$	< 5.4	$\times 10^{-4}$	CL=90%	1430
$\eta \rho \bar{\rho} \pi^+ \pi^-$	< 3.3	$\times 10^{-3}$	CL=90%	1284
$\rho^0 \rho \bar{\rho}$	< 1.7	$\times 10^{-3}$	CL=90%	1313
$\rho \bar{\rho} K^+ K^-$	< 3.2	$\times 10^{-4}$	CL=90%	1185
$\eta \rho \bar{\rho} K^+ K^-$	< 6.9	$\times 10^{-3}$	CL=90%	736
$\pi^0 \rho \bar{\rho} K^+ K^-$	< 1.2	$\times 10^{-3}$	CL=90%	1093
$\phi \rho \bar{\rho}$	< 1.3	$\times 10^{-4}$	CL=90%	1178
$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.5	$\times 10^{-4}$	CL=90%	1404
$\Lambda \bar{\rho} K^+$	< 2.8	$\times 10^{-4}$	CL=90%	1387
$\Lambda \bar{\rho} K^+ \pi^+ \pi^-$	< 6.3	$\times 10^{-4}$	CL=90%	1234
$\Lambda \bar{\Lambda} \eta$	< 1.9	$\times 10^{-4}$	CL=90%	1262
$\Sigma^+ \bar{\Sigma}^-$	< 1.0	$\times 10^{-4}$	CL=90%	1464
$\Sigma^0 \bar{\Sigma}^0$	< 4	$\times 10^{-5}$	CL=90%	1462
$\Xi^+ \bar{\Xi}^-$	< 1.5	$\times 10^{-4}$	CL=90%	1346
$\Xi^0 \bar{\Xi}^0$	< 1.4	$\times 10^{-4}$	CL=90%	1353

Radiative decays

$\gamma \chi_{c2}$	< 6.4	$\times 10^{-4}$	CL=90%	211
$\gamma \chi_{c1}$	$(2.49 \pm 0.23) \times 10^{-3}$			253
$\gamma \chi_{c0}$	$(6.9 \pm 0.6) \times 10^{-3}$			341
$\gamma \eta_c$	< 7	$\times 10^{-4}$	CL=90%	707
$\gamma \eta_c(2S)$	< 9	$\times 10^{-4}$	CL=90%	133
$\gamma \eta'$	< 1.8	$\times 10^{-4}$	CL=90%	1765
$\gamma \eta$	< 1.5	$\times 10^{-4}$	CL=90%	1847
$\gamma \pi^0$	< 2	$\times 10^{-4}$	CL=90%	1884

$\psi_2(3823)$

$$I^G(J^{PC}) = 0^-(2^{--})$$

I, J, P need confirmation.

Mass $m = 3822.2 \pm 1.2$ MeV

Full width $\Gamma < 16$ MeV, CL = 90%

$\psi_2(3823)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\chi_{c1}\gamma$	seen	299
$\chi_{c2}\gamma$	not seen	257

 $\chi_{c1}(3872)$

$$J^{PC} = 0^+(1^{++})$$

 Mass $m = 3871.69 \pm 0.17$ MeV

 $m_{\chi_{c1}(3872)} - m_{J/\psi} = 775 \pm 4$ MeV

 Full width $\Gamma < 1.2$ MeV, CL = 90%

$\chi_{c1}(3872)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi^+\pi^-J/\psi(1S)$	> 3.2 %	650
$\omega J/\psi(1S)$	> 2.3 %	†
$D^0\bar{D}^0\pi^0$	> 40 %	117
$\bar{D}^{*0}D^0$	> 30 %	4
$\gamma J/\psi$	$> 7 \times 10^{-3}$	697
$\gamma\psi(2S)$	> 4 %	181
$\pi^+\pi^-\eta_c(1S)$	not seen	746
$\pi^+\pi^-\chi_{c1}$	not seen	218
$p\bar{p}$	not seen	1693

 $Z_c(3900)$

$$J^{PC} = 1^+(1^{+-})$$

 Mass $m = 3887.2 \pm 2.3$ MeV ($S = 1.6$)

 Full width $\Gamma = 28.2 \pm 2.6$ MeV

$Z_c(3900)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$J/\psi\pi$	seen	699
$h_c\pi^\pm$	not seen	318
$\eta_c\pi^+\pi^-$	not seen	759
$(D\bar{D}^*)^\pm$	seen	—
$D^0D^{*-} + c.c.$	seen	153
$D^-D^{*0} + c.c.$	seen	144
$\omega\pi^\pm$	not seen	1862
$J/\psi\eta$	not seen	510
$D^+D^{*-} + c.c.$	seen	—
$D^0\bar{D}^{*0} + c.c.$	seen	—

X(3915)

$$I^G(J^{PC}) = 0^+(0 \text{ or } 2^{++})$$

Mass $m = 3918.4 \pm 1.9$ MeVFull width $\Gamma = 20 \pm 5$ MeV ($S = 1.1$)

X(3915) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\omega J/\psi$	seen	222
$\pi^+ \pi^- \eta_c(1S)$	not seen	785
$\eta_c \eta$	not seen	665
$\eta_c \pi^0$	not seen	814
$K \bar{K}$	not seen	1896
$\gamma \gamma$	seen	1959

 $\chi_{c2}(3930)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 3927.2 \pm 2.6$ MeVFull width $\Gamma = 24 \pm 6$ MeV

$\chi_{c2}(3930)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma \gamma$	seen	1964
$D \bar{D}$	seen	615
$D^+ D^-$	seen	600
$D^0 \bar{D}^0$	seen	615
$\pi^+ \pi^- \eta_c(1S)$	not seen	792
$K \bar{K}$	not seen	1901

X(4020)

$$I^G(J^{PC}) = 1^+(?^{? -})$$

Mass $m = 4024.1 \pm 1.9$ MeVFull width $\Gamma = 13 \pm 5$ MeV ($S = 1.7$)

X(4020) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$h_c(1P)\pi$	seen	450
$D^* \bar{D}^*$	seen	85
$D \bar{D}^* + \text{c.c.}$	not seen	542
$\eta_c \pi^+ \pi^-$	not seen	872
$J/\psi(1S)\pi^\pm$	not seen	811

$\psi(4040)$ [*h*]

$$I^G(J^{PC}) = 0^-(1^{--})$$

 Mass $m = 4039 \pm 1$ MeV

 Full width $\Gamma = 80 \pm 10$ MeV

 $\Gamma_{ee} = 0.86 \pm 0.07$ keV

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

$\psi(4040)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
e^+e^-	$(1.07 \pm 0.16) \times 10^{-5}$		2019
$D\bar{D}$	seen		775
$D^0\bar{D}^0$	seen		775
D^+D^-	seen		763
$D^*\bar{D} + \text{c.c.}$	seen		569
$D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen		575
$D^*(2010)^+D^- + \text{c.c.}$	seen		561
$D^*\bar{D}^*$	seen		193
$D^*(2007)^0\bar{D}^*(2007)^0$	seen		226
$D^*(2010)^+D^*(2010)^-$	seen		193
$D^0D^-\pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0\bar{D}^0 + \text{c.c.},$ $D^*(2010)^+D^- + \text{c.c.}$)	not seen		—
$D\bar{D}^*\pi$ (excl. $D^*\bar{D}^*$)	not seen		—
$D^0\bar{D}^{*-}\pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+D^*(2010)^-$)	seen		—
$D_s^+D_s^-$	seen		452
$J/\psi\pi^+\pi^-$	< 4	$\times 10^{-3}$	90% 794
$J/\psi\pi^0\pi^0$	< 2	$\times 10^{-3}$	90% 797
$J/\psi\eta$	(5.2 ± 0.7)	$\times 10^{-3}$	675
$J/\psi\pi^0$	< 2.8	$\times 10^{-4}$	90% 823
$J/\psi\pi^+\pi^-\pi^0$	< 2	$\times 10^{-3}$	90% 746
$\chi_{c1}\gamma$	< 3.4	$\times 10^{-3}$	90% 494
$\chi_{c2}\gamma$	< 5	$\times 10^{-3}$	90% 454
$\chi_{c1}\pi^+\pi^-\pi^0$	< 1.1	%	90% 306
$\chi_{c2}\pi^+\pi^-\pi^0$	< 3.2	%	90% 233
$h_c(1P)\pi^+\pi^-$	< 3	$\times 10^{-3}$	90% 403
$\phi\pi^+\pi^-$	< 3	$\times 10^{-3}$	90% 1880
$\Lambda\bar{\Lambda}\pi^+\pi^-$	< 2.9	$\times 10^{-4}$	90% 1578
$\Lambda\bar{\Lambda}\pi^0$	< 9	$\times 10^{-5}$	90% 1636
$\Lambda\bar{\Lambda}\eta$	< 3.0	$\times 10^{-4}$	90% 1452
$\Sigma^+\bar{\Sigma}^-$	< 1.3	$\times 10^{-4}$	90% 1632

$\Sigma^0 \bar{\Sigma}^0$	< 7	$\times 10^{-5}$	90%	1630
$\Xi^+ \bar{\Xi}^-$	< 1.6	$\times 10^{-4}$	90%	1527
$\Xi^0 \bar{\Xi}^0$	< 1.8	$\times 10^{-4}$	90%	1533

$\chi_{c1}(4140)$

$$J^{PC} = 0^+(1^{++})$$

Mass $m = 4146.8 \pm 2.4$ MeV ($S = 1.1$)

Full width $\Gamma = 22_{-7}^{+8}$ MeV ($S = 1.3$)

$\chi_{c1}(4140)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$J/\psi \phi$	seen	217
$\gamma\gamma$	not seen	2073

$\psi(4160)$ ^[h]

$$J^{PC} = 0^-(1^{--})$$

Mass $m = 4191 \pm 5$ MeV

Full width $\Gamma = 70 \pm 10$ MeV

$\Gamma_{ee} = 0.48 \pm 0.22$ keV

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

$\psi(4160)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$		2096
$\mu^+ \mu^-$	seen		2093
$D \bar{D}$	seen		956
$D^0 \bar{D}^0$	seen		956
$D^+ D^-$	seen		947
$D^* \bar{D} + \text{c.c.}$	seen		798
$D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen		802
$D^*(2010)^+ D^- + \text{c.c.}$	seen		792
$D^* \bar{D}^*$	seen		592
$D^*(2007)^0 \bar{D}^*(2007)^0$	seen		604
$D^*(2010)^+ D^*(2010)^-$	seen		592
$D^0 D^- \pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0 \bar{D}^0 + \text{c.c.},$ $D^*(2010)^+ D^- + \text{c.c.}$)	not seen		—
$D \bar{D}^* \pi + \text{c.c.}$ (excl. $D^* \bar{D}^*$)	seen		—
$D^0 D^{*-} \pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+ D^*(2010)^-$)	not seen		—

$D_s^+ D_s^-$	not seen			719
$D_s^{*+} D_s^- + \text{c.c.}$	seen			385
$J/\psi \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%	919
$J/\psi \pi^0 \pi^0$	< 3	$\times 10^{-3}$	90%	922
$J/\psi K^+ K^-$	< 2	$\times 10^{-3}$	90%	407
$J/\psi \eta$	< 8	$\times 10^{-3}$	90%	822
$J/\psi \pi^0$	< 1	$\times 10^{-3}$	90%	944
$J/\psi \eta'$	< 5	$\times 10^{-3}$	90%	457
$J/\psi \pi^+ \pi^- \pi^0$	< 1	$\times 10^{-3}$	90%	879
$\psi(2S) \pi^+ \pi^-$	< 4	$\times 10^{-3}$	90%	396
$\chi_{c1} \gamma$	< 5	$\times 10^{-3}$	90%	625
$\chi_{c2} \gamma$	< 1.3	%	90%	587
$\chi_{c1} \pi^+ \pi^- \pi^0$	< 2	$\times 10^{-3}$	90%	496
$\chi_{c2} \pi^+ \pi^- \pi^0$	< 8	$\times 10^{-3}$	90%	445
$h_c(1P) \pi^+ \pi^-$	< 5	$\times 10^{-3}$	90%	556
$h_c(1P) \pi^0 \pi^0$	< 2	$\times 10^{-3}$	90%	560
$h_c(1P) \eta$	< 2	$\times 10^{-3}$	90%	348
$h_c(1P) \pi^0$	< 4	$\times 10^{-4}$	90%	600
$\phi \pi^+ \pi^-$	< 2	$\times 10^{-3}$	90%	1961
$\gamma \chi_{c1}(3872) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 6.8	$\times 10^{-5}$	90%	—
$\gamma X(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.36	$\times 10^{-4}$	90%	—
$\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.18	$\times 10^{-4}$	90%	—
$\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.47	$\times 10^{-4}$	90%	—
$\gamma \chi_{c1}(3872) \rightarrow \gamma \gamma J/\psi$	< 1.05	$\times 10^{-4}$	90%	—
$\gamma X(3915) \rightarrow \gamma \gamma J/\psi$	< 1.26	$\times 10^{-4}$	90%	—
$\gamma X(3930) \rightarrow \gamma \gamma J/\psi$	< 8.8	$\times 10^{-5}$	90%	—
$\gamma X(3940) \rightarrow \gamma \gamma J/\psi$	< 1.79	$\times 10^{-4}$	90%	—

ψ(4260)

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 4230 \pm 8$ MeV (S = 2.9)

Full width $\Gamma = 55 \pm 19$ MeV (S = 4.4)

ψ(4260) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$J/\psi \pi^+ \pi^-$	seen	950
$J/\psi f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$	seen	—
$Z_c(3900)^\pm \pi^\mp, Z_c^\pm \rightarrow J/\psi \pi^\pm$	seen	—
$J/\psi \pi^0 \pi^0$	seen	952
$J/\psi K^+ K^-$	seen	477
$J/\psi K_S^0 K_S^0$	not seen	465
$J/\psi \eta$	not seen	857
$J/\psi \pi^0$	not seen	974

$J/\psi\eta'$	not seen	520
$J/\psi\pi^+\pi^-\pi^0$	not seen	912
$J/\psi\eta\pi^0$	not seen	780
$J/\psi\eta\eta$	not seen	247
$\psi(2S)\pi^+\pi^-$	not seen	437
$\psi(2S)\eta$	not seen	†
$\chi_{c0}\omega$	not seen	205
$\chi_{c1}\pi^+\pi^-\pi^0$	not seen	537
$\chi_{c2}\pi^+\pi^-\pi^0$	not seen	489
$h_c(1P)\pi^+\pi^-$	not seen	593
$\phi\pi^+\pi^-$	not seen	1982
$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$	not seen	—
$D\bar{D}$	not seen	998
$D^0\bar{D}^0$	not seen	998
D^+D^-	not seen	989
$D^*\bar{D} + \text{c.c.}$	not seen	887
$D^*(2007)^0\bar{D}^0 + \text{c.c.}$	not seen	—
$D^*(2010)^+D^- + \text{c.c.}$	not seen	—
$D^*\bar{D}^*$	not seen	657
$D^*(2007)^0\bar{D}^*(2007)^0$	not seen	668
$D^*(2010)^+D^*(2010)^-$	not seen	657
$D^0D^-\pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0\bar{D}^{*0} + \text{c.c.},$ $D^*(2010)^+D^- + \text{c.c.}$)	not seen	—
$D\bar{D}^*\pi + \text{c.c.}$ (excl. $D^*\bar{D}^*$)	not seen	723
$D^0D^{*-}\pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+D^*(2010)^-$)	not seen	—
$D^0D^*(2010)^-\pi^+ + \text{c.c.}$	not seen	716
$D^*\bar{D}^*\pi$	not seen	395
$D_s^+D_s^-$	not seen	774
$D_s^{*+}D_s^- + \text{c.c.}$	not seen	615
$D_s^{*+}D_s^{*-}$	not seen	109
$\rho\bar{\rho}$	not seen	1896
$\rho\bar{\rho}\pi^0$	not seen	1860
$K_S^0K^\pm\pi^\mp$	not seen	2037
$K^+K^-\pi^0$	not seen	2038

Radiative decays

$\eta_c(1S)\gamma$	possibly seen	1063
$\chi_{c1}\gamma$	not seen	658
$\chi_{c2}\gamma$	not seen	620
$\chi_{c1}(3872)\gamma$	seen	343

$\chi_{c1}(4274)$

$$J^{PC} = 0^+(1^{++})$$

$$\text{Mass } m = 4274^{+8}_{-6} \text{ MeV}$$

$$\text{Full width } \Gamma = 49 \pm 12 \text{ MeV}$$

$\chi_{c1}(4274)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$J/\psi \phi$	seen	503

 $\psi(4360)$

$$J^{PC} = 0^-(1^{--})$$

$$\psi(4360) \text{ MASS} = 4368 \pm 13 \text{ MeV} \quad (S = 3.7)$$

$$\psi(4360) \text{ WIDTH} = 96 \pm 7 \text{ MeV}$$

$\psi(4360)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\psi(2S)\pi^+\pi^-$	seen	573
$\psi_2(3823)\pi^+\pi^-$	possibly seen	440

 $\psi(4415)$ [h]

$$J^{PC} = 0^-(1^{--})$$

$$\text{Mass } m = 4421 \pm 4 \text{ MeV}$$

$$\text{Full width } \Gamma = 62 \pm 20 \text{ MeV}$$

$$\Gamma_{ee} = 0.58 \pm 0.07 \text{ keV}$$

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

$\psi(4415)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$D\bar{D}$	seen		1187
$D^0\bar{D}^0$	seen		1187
D^+D^-	seen		1179
$D^*\bar{D} + \text{c.c.}$	seen		1063
$D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen		1067
$D^*(2010)^+D^- + \text{c.c.}$	seen		1059
$D^*\bar{D}^*$	seen		919
$D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen		927
$D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen		919
$D^0D^-\pi^+$ (excl. $D^*(2007)^0\bar{D}^0$ +c.c., $D^*(2010)^+D^-$ +c.c.)	< 2.3	%	90%
			–

$D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.}$	$(10 \pm 4) \%$		–
$D^0 D^{*-} \pi^+ + \text{c.c.}$	$< 11 \%$	90%	926
$D_s^+ D_s^-$	not seen		1006
$\omega \chi_{c2}$	possibly seen		330
$D_s^{*+} D_s^- + \text{c.c.}$	seen		–
$D_s^{*+} D_s^{*-}$	not seen		652
$\psi_2(3823) \pi^+ \pi^-$	possibly seen		494
$J/\psi \eta$	$< 6 \times 10^{-3}$	90%	1022
$\chi_{c1} \gamma$	$< 8 \times 10^{-4}$	90%	817
$\chi_{c2} \gamma$	$< 4 \times 10^{-3}$	90%	780
$e^+ e^-$	$(9.4 \pm 3.2) \times 10^{-6}$		2210

$Z_c(4430)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

G, C need confirmation.

Quantum numbers not established.

$$\text{Mass } m = 4478_{-18}^{+15} \text{ MeV}$$

$$\text{Full width } \Gamma = 181 \pm 31 \text{ MeV}$$

$Z_c(4430)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi^+ \psi(2S)$	seen	711
$\pi^+ J/\psi$	seen	1162

$\psi(4660)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

I needs confirmation.

$$\psi(4660) \text{ MASS} = 4643 \pm 9 \text{ MeV} \quad (S = 1.2)$$

$$\psi(4660) \text{ WIDTH} = 72 \pm 11 \text{ MeV}$$

$\psi(4660)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\psi(2S) \pi^+ \pi^-$	seen	820

NOTES

- [a] For $E_\gamma > 100$ MeV.
- [b] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [c] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.
- [d] See the “Note on the $\eta(1405)$ ” in the $\eta(1405)$ Particle Listings.
- [e] For a narrow state A with mass less than 960 MeV.
- [f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.
- [g] For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.
- [h] J^{PC} known by production in e^+e^- via single photon annihilation. I^G is not known; interpretation of this state as a single resonance is unclear because of the expectation of substantial threshold effects in this energy region.