

CHARMED MESONS

($C = \pm 1$)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \quad \text{similarly for } D^{*'}\text{'s}$$

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1869.65 \pm 0.05 \text{ MeV}$$

$$\text{Mean life } \tau = (1040 \pm 7) \times 10^{-15} \text{ s}$$

$$c\tau = 311.8 \text{ } \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.096 \pm 0.004 \text{ [a]}$$

$$\Gamma(c \rightarrow D^*(2010)^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_L^0 e^\pm \nu) = (-0.6 \pm 1.6)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.41 \pm 0.09)\%$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.18 \pm 0.16)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (-0.3 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (-0.1 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.0 \pm 1.2)\%$$

$$A_{CP}(\pi^\pm \pi^0) = (2.4 \pm 1.2)\%$$

$$A_{CP}(\pi^\pm \eta) = (1.0 \pm 1.5)\% \quad (S = 1.4)$$

$$A_{CP}(\pi^\pm \eta'(958)) = (-0.6 \pm 0.7)\%$$

$$A_{CP}(\bar{K}^0 / K^0 K^\pm) = (0.11 \pm 0.17)\%$$

$$A_{CP}(K_S^0 K^\pm) = (-0.11 \pm 0.25)\%$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.37 \pm 0.29)\%$$

$$A_{CP}(K^\pm K^{*0}) = (-0.3 \pm 0.4)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.09 \pm 0.19)\% \quad (S = 1.2)$$

$$A_{CP}(K^\pm K_0^*(1430)^0) = (8_{-6}^{+7})\%$$

$$A_{CP}(K^\pm K_2^*(1430)^0) = (43_{-26}^{+20})\%$$

$$A_{CP}(K^\pm K_0^*(700)) = (-12_{-13}^{+18})\%$$

$$A_{CP}(a_0(1450)^0 \pi^\pm) = (-19_{-16}^{+14})\%$$

$$A_{CP}(\phi(1680) \pi^\pm) = (-9 \pm 26)\%$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = (-2 \pm 4)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) = (-4 \pm 7)\%$$

$$A_{CP}(K^\pm \pi^0) = (-4 \pm 11)\%$$

χ^2 tests of CP-violation (CPV)

Local CPV in $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm = 78.1\%$

Local CPV in $D^\pm \rightarrow K^+ K^- \pi^\pm = 31\%$

CP violating asymmetries of P-odd (T-odd) moments

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-12 \pm 11) \times 10^{-3} [b]$$

D^+ form factors

$$f_+(0)|V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell = 0.719 \pm 0.011 \quad (S = 1.6)$$

$$r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell = -2.13 \pm 0.14$$

$$r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell = -3 \pm 12 \quad (S = 1.5)$$

$$f_+(0)|V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell = 0.1407 \pm 0.0025$$

$$r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell = -2.00 \pm 0.13$$

$$r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell = -4 \pm 5$$

$$f_+(0)|V_{cd}| \text{ in } D^+ \rightarrow \eta e^+ \nu_e = (8.3 \pm 0.5) \times 10^{-2}$$

$$r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e = -5.3 \pm 2.7 \quad (S = 1.9)$$

$$r_\nu \equiv V(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e = 1.24 \pm 0.11$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e = 1.06 \pm 0.16$$

$$r_\nu \equiv V(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e = 1.48 \pm 0.16$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e = 0.83 \pm 0.12$$

$$r_\nu \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 1.49 \pm 0.05 \quad (S = 2.1)$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.802 \pm 0.021$$

$$r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.0 \pm 0.4$$

$$\Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 1.13 \pm 0.08$$

$$\Gamma_+/ \Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.22 \pm 0.06 \quad (S = 1.6)$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07 \pm 0.30) %		—
μ^+ anything	(17.6 \pm 3.2) %		—
K^- anything	(25.7 \pm 1.4) %		—
\bar{K}^0 anything + K^0 anything	(61 \pm 5) %		—
K^+ anything	(5.9 \pm 0.8) %		—
$K^*(892)^-$ anything	(6 \pm 5) %		—
$\bar{K}^*(892)^0$ anything	(23 \pm 5) %		—
$K^*(892)^0$ anything	< 6.6	%	CL=90%

η anything	(6.3 \pm 0.7)%	–
η' anything	(1.04 \pm 0.18)%	–
ϕ anything	(1.03 \pm 0.12)%	–

Leptonic and semileptonic modes

$e^+ \nu_e$	< 8.8	$\times 10^{-6}$ CL=90%	935
$\gamma e^+ \nu_e$	< 3.0	$\times 10^{-5}$ CL=90%	935
$\mu^+ \nu_\mu$	(3.74 \pm 0.17)	$\times 10^{-4}$	932
$\tau^+ \nu_\tau$	< 1.2	$\times 10^{-3}$ CL=90%	90
$\bar{K}^0 e^+ \nu_e$	(8.73 \pm 0.10)%		869
$\bar{K}^0 \mu^+ \nu_\mu$	(8.76 \pm 0.19)%		865
$K^- \pi^+ e^+ \nu_e$	(4.02 \pm 0.18)%	S=3.2	864
$\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow$ $K^- \pi^+$	(3.77 \pm 0.17)%		722
$(K^- \pi^+) [0.8-1.0] \text{GeV} e^+ \nu_e$	(3.39 \pm 0.09)%		864
$(K^- \pi^+)_{S\text{-wave}} e^+ \nu_e$	(2.28 \pm 0.11)	$\times 10^{-3}$	–
$\bar{K}^*(1410)^0 e^+ \nu_e,$ $\bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	< 6	$\times 10^{-3}$ CL=90%	–
$\bar{K}_2^*(1430)^0 e^+ \nu_e,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	< 5	$\times 10^{-4}$ CL=90%	–
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7	$\times 10^{-3}$ CL=90%	864
$\bar{K}^*(892)^0 e^+ \nu_e$	(5.40 \pm 0.10)%	S=1.1	722
$K^- \pi^+ \mu^+ \nu_\mu$	(3.65 \pm 0.34)%		851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.52 \pm 0.10)%		717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(1.9 \pm 0.5)	$\times 10^{-3}$	851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(5.27 \pm 0.15)%		717
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.5	$\times 10^{-3}$ CL=90%	825
$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.3	$\times 10^{-4}$ CL=90%	380
$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.5	$\times 10^{-3}$ CL=90%	105
$\pi^0 e^+ \nu_e$	(3.72 \pm 0.17)	$\times 10^{-3}$ S=2.0	930
$\pi^0 \mu^+ \nu_\mu$	(3.50 \pm 0.15)	$\times 10^{-3}$	927
$\eta e^+ \nu_e$	(1.11 \pm 0.07)	$\times 10^{-3}$	855
$\rho^0 e^+ \nu_e$	(2.18 \pm 0.17)	$\times 10^{-3}$	774
$\rho^0 \mu^+ \nu_\mu$	(2.4 \pm 0.4)	$\times 10^{-3}$	770
$\omega e^+ \nu_e$	(1.69 \pm 0.11)	$\times 10^{-3}$	771
$\eta'(958) e^+ \nu_e$	(2.0 \pm 0.4)	$\times 10^{-4}$	690
$a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0$	(1.7 \pm 0.8)	$\times 10^{-4}$	–
$\phi e^+ \nu_e$	< 1.3	$\times 10^{-5}$ CL=90%	657
$D^0 e^+ \nu_e$	< 1.0	$\times 10^{-4}$ CL=90%	5

Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$

$K_S^0 \pi^+$	(1.562 ± 0.031) %	S=1.7	863
$K_L^0 \pi^+$	(1.46 ± 0.05) %		863
$K^- 2\pi^+$	[c] (9.38 ± 0.16) %	S=1.6	846
$(K^- \pi^+)_{S\text{-wave}} \pi^+$	(7.52 ± 0.17) %		846
$\bar{K}_0^*(1430)^0 \pi^+$,	[d] (1.25 ± 0.06) %		382
$\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(892)^0 \pi^+$,	(1.04 ± 0.12) %		714
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(1410)^0 \pi^+$, $\bar{K}^{*0} \rightarrow$	not seen		381
$K^- \pi^+$			
$\bar{K}_2^*(1430)^0 \pi^+$,	[d] (2.3 ± 0.7) × 10 ⁻⁴		371
$\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(1680)^0 \pi^+$,	[d] (2.2 ± 1.1) × 10 ⁻⁴		58
$\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$			
$K^-(2\pi^+)_{I=2}$	(1.45 ± 0.26) %		–
$K_S^0 \pi^+ \pi^0$	[c] (7.36 ± 0.21) %		845
$K_S^0 \rho^+$	(6.14 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.60 \\ 0.35 \end{smallmatrix}$) %		677
$K_S^0 \rho(1450)^+$, $\rho^+ \rightarrow \pi^+ \pi^0$	(1.5 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.2 \\ 1.4 \end{smallmatrix}$) × 10 ⁻³		–
$\bar{K}^*(892)^0 \pi^+$,	(2.64 ± 0.32) × 10 ⁻³		714
$\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$			
$\bar{K}_0^*(1430)^0 \pi^+$, $\bar{K}_0^{*0} \rightarrow$	(2.7 ± 0.9) × 10 ⁻³		–
$K_S^0 \pi^0$			
$\bar{K}_0^*(1680)^0 \pi^+$, $\bar{K}_0^{*0} \rightarrow$	(10 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 7 \\ 10 \end{smallmatrix}$) × 10 ⁻⁴		–
$K_S^0 \pi^0$			
$\bar{K}^0 \pi^+$, $\bar{K}^0 \rightarrow K_S^0 \pi^0$	(6 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 5 \\ 4 \end{smallmatrix}$) × 10 ⁻³		–
$K_S^0 \pi^+ \pi^0$ nonresonant	(3 ± 4) × 10 ⁻³		845
$K_S^0 \pi^+ \pi^0$ nonresonant and	(1.37 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.21 \\ 0.40 \end{smallmatrix}$) %		–
$\bar{K}^0 \pi^+$			
$(K_S^0 \pi^0)_{S\text{-wave}} \pi^+$	(1.27 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.27 \\ 0.33 \end{smallmatrix}$) %		845
$K_S^0 \pi^+ \eta'(958)$	(1.90 ± 0.21) × 10 ⁻³		481
$K^- 2\pi^+ \pi^0$	[e] (6.25 ± 0.18) %		816
$K_S^0 2\pi^+ \pi^-$	[e] (3.10 ± 0.09) %		814
$K^- 3\pi^+ \pi^-$	[c] (5.7 ± 0.5) × 10 ⁻³	S=1.1	772
$\bar{K}^*(892)^0 2\pi^+ \pi^-$,	(1.2 ± 0.4) × 10 ⁻³		645
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(892)^0 \rho^0 \pi^+$,	(2.3 ± 0.4) × 10 ⁻³		239
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(892)^0 a_1(1260)^+$	[f] (9.3 ± 1.9) × 10 ⁻³		†

$K^- \rho^0 2\pi^+$	$(1.72 \pm 0.28) \times 10^{-3}$	524
$K^- 3\pi^+ \pi^-$ nonresonant	$(4.0 \pm 2.9) \times 10^{-4}$	772
$K^+ 2K_S^0$	$(2.54 \pm 0.13) \times 10^{-3}$	545
$K^+ K^- K_S^0 \pi^+$	$(2.4 \pm 0.5) \times 10^{-4}$	436

Pionic modes

$\pi^+ \pi^0$	$(1.247 \pm 0.033) \times 10^{-3}$	925
$2\pi^+ \pi^-$	$(3.27 \pm 0.18) \times 10^{-3}$	909
$\rho^0 \pi^+$	$(8.3 \pm 1.5) \times 10^{-4}$	767
$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	$(1.83 \pm 0.16) \times 10^{-3}$	909
$\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-$	$(1.38 \pm 0.12) \times 10^{-3}$	–
$f_0(980) \pi^+,$ $f_0(980) \rightarrow \pi^+ \pi^-$	$(1.56 \pm 0.33) \times 10^{-4}$	669
$f_0(1370) \pi^+,$ $f_0(1370) \rightarrow \pi^+ \pi^-$	$(8 \pm 4) \times 10^{-5}$	–
$f_2(1270) \pi^+,$ $f_2(1270) \rightarrow \pi^+ \pi^-$	$(5.0 \pm 0.9) \times 10^{-4}$	485
$\rho(1450)^0 \pi^+,$ $\rho(1450)^0 \rightarrow \pi^+ \pi^-$	$< 8 \times 10^{-5} \text{CL=95\%}$	338
$f_0(1500) \pi^+,$ $f_0(1500) \rightarrow \pi^+ \pi^-$	$(1.1 \pm 0.4) \times 10^{-4}$	–
$f_0(1710) \pi^+,$ $f_0(1710) \rightarrow \pi^+ \pi^-$	$< 5 \times 10^{-5} \text{CL=95\%}$	–
$f_0(1790) \pi^+,$ $f_0(1790) \rightarrow \pi^+ \pi^-$	$< 7 \times 10^{-5} \text{CL=95\%}$	–
$(\pi^+ \pi^+)_{S\text{-wave}} \pi^-$	$< 1.2 \times 10^{-4} \text{CL=95\%}$	909
$2\pi^+ \pi^-$ nonresonant	$< 1.1 \times 10^{-4} \text{CL=95\%}$	909
$\pi^+ 2\pi^0$	$(4.7 \pm 0.4) \times 10^{-3}$	910
$2\pi^+ \pi^- \pi^0$	$(1.16 \pm 0.08) \%$	883
$3\pi^+ 2\pi^-$	$(1.66 \pm 0.16) \times 10^{-3}$ S=1.1	845
$\eta \pi^+$	$(3.77 \pm 0.09) \times 10^{-3}$	848
$\eta \pi^+ \pi^0$	$(1.38 \pm 0.35) \times 10^{-3}$	831
$\omega \pi^+$	$(2.8 \pm 0.6) \times 10^{-4}$	764
$\eta'(958) \pi^+$	$(4.97 \pm 0.19) \times 10^{-3}$	681
$\eta'(958) \pi^+ \pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$	654

Hadronic modes with a $K\bar{K}$ pair

$K^+ K_S^0$	$(3.04 \pm 0.09) \times 10^{-3}$ S=2.2	793
$K^+ K^- \pi^+$	[c] $(9.93 \pm 0.24) \times 10^{-3}$	744
$\phi \pi^+, \phi \rightarrow K^+ K^-$	$(2.76 \pm_{-0.09}^{+0.08}) \times 10^{-3}$	647
$K^+ \bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(2.55 \pm_{-0.14}^{+0.09}) \times 10^{-3}$	613
$K^+ \bar{K}_0^*(1430)^0,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	$(1.9 \pm 0.4) \times 10^{-3}$	–

$K^+ \bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^- \pi^+$	$(1.7 \begin{smallmatrix} + 1.3 \\ - 0.8 \end{smallmatrix}) \times 10^{-4}$	—
$K^+ \bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^- \pi^+$	$(6.9 \begin{smallmatrix} + 4.0 \\ - 2.1 \end{smallmatrix}) \times 10^{-4}$	—
$a_0(1450)^0 \pi^+, a_0^0 \rightarrow K^+ K^-$	$(4.6 \begin{smallmatrix} + 7.0 \\ - 1.9 \end{smallmatrix}) \times 10^{-4}$	—
$\phi(1680) \pi^+, \phi \rightarrow K^+ K^-$	$(5.1 \begin{smallmatrix} + 4.0 \\ - 1.9 \end{smallmatrix}) \times 10^{-5}$	—
$K_S^0 K_S^0 \pi^+$	$(2.70 \pm 0.13) \times 10^{-3}$	741
$K^+ K_S^0 \pi^+ \pi^-$	$(1.74 \pm 0.18) \times 10^{-3}$	678
$K_S^0 K^- 2\pi^+$	$(2.38 \pm 0.17) \times 10^{-3}$	678
$K^+ K^- 2\pi^+ \pi^-$	$(2.3 \pm 1.2) \times 10^{-4}$	601

A few poorly measured branching fractions:

$\phi \pi^+ \pi^0$	$(2.3 \pm 1.0) \%$	619
$\phi \rho^+$	$< 1.5 \%$ CL=90%	260
$K^+ K^- \pi^+ \pi^0$ non- ϕ	$(1.5 \begin{smallmatrix} + 0.7 \\ - 0.6 \end{smallmatrix}) \%$	682
$K^*(892)^+ K_S^0$	$(1.7 \pm 0.8) \%$	612

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(2.08 \pm 0.21) \times 10^{-4}$	S=1.4	864
$K^+ \eta$	$(1.25 \pm 0.16) \times 10^{-4}$	S=1.1	776
$K^+ \eta'(958)$	$(1.85 \pm 0.20) \times 10^{-4}$		571
$K^+ \pi^+ \pi^-$	$(5.42 \pm 0.22) \times 10^{-4}$		846
$K^+ \rho^0$	$(2.1 \pm 0.5) \times 10^{-4}$		679
$K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow K^+ \pi^-$	$(2.5 \pm 0.4) \times 10^{-4}$		714
$K^+ f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$	$(4.8 \pm 2.9) \times 10^{-5}$		—
$K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow K^+ \pi^-$	$(4.3 \pm 2.9) \times 10^{-5}$		—
$K^+ \pi^+ \pi^-$ nonresonant	not seen		846
$2K^+ K^-$	$(8.9 \pm 2.1) \times 10^{-5}$		550

$\Delta C = 1$ weak neutral current ($C1$) modes, or Lepton Family number (LF) or Lepton number (L) violating modes

$\pi^+ e^+ e^-$	$C1$	$< 1.1 \times 10^{-6}$ CL=90%	930
$\pi^+ \pi^0 e^+ e^-$		$< 1.4 \times 10^{-5}$ CL=90%	925
$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[g]	$(1.7 \begin{smallmatrix} + 1.4 \\ - 0.9 \end{smallmatrix}) \times 10^{-6}$	—
$\pi^+ \mu^+ \mu^-$	$C1$	$< 7.3 \times 10^{-8}$ CL=90%	918
$\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$	[g]	$(1.8 \pm 0.8) \times 10^{-6}$	—
$\rho^+ \mu^+ \mu^-$	$C1$	$< 5.6 \times 10^{-4}$ CL=90%	757
$K^+ e^+ e^-$	[h]	$< 1.0 \times 10^{-6}$ CL=90%	870
$K^+ \pi^0 e^+ e^-$		$< 1.5 \times 10^{-5}$ CL=90%	864

$K_S^0 \pi^+ e^+ e^-$		< 2.6	$\times 10^{-5} \text{CL}=90\%$	—
$K_S^0 K^+ e^+ e^-$		< 1.1	$\times 10^{-5} \text{CL}=90\%$	—
$K^+ \mu^+ \mu^-$		$[h] < 4.3$	$\times 10^{-6} \text{CL}=90\%$	856
$\pi^+ e^+ \mu^-$	LF	< 2.9	$\times 10^{-6} \text{CL}=90\%$	927
$\pi^+ e^- \mu^+$	LF	< 3.6	$\times 10^{-6} \text{CL}=90\%$	927
$K^+ e^+ \mu^-$	LF	< 1.2	$\times 10^{-6} \text{CL}=90\%$	866
$K^+ e^- \mu^+$	LF	< 2.8	$\times 10^{-6} \text{CL}=90\%$	866
$\pi^- 2e^+$	L	< 1.1	$\times 10^{-6} \text{CL}=90\%$	930
$\pi^- 2\mu^+$	L	< 2.2	$\times 10^{-8} \text{CL}=90\%$	918
$\pi^- e^+ \mu^+$	L	< 2.0	$\times 10^{-6} \text{CL}=90\%$	927
$\rho^- 2\mu^+$	L	< 5.6	$\times 10^{-4} \text{CL}=90\%$	757
$K^- 2e^+$	L	< 9	$\times 10^{-7} \text{CL}=90\%$	870
$K^- 2\mu^+$	L	< 1.0	$\times 10^{-5} \text{CL}=90\%$	856
$K^- e^+ \mu^+$	L	< 1.9	$\times 10^{-6} \text{CL}=90\%$	866
$K^*(892)^- 2\mu^+$	L	< 8.5	$\times 10^{-4} \text{CL}=90\%$	703

 D^0

$$I(J^P) = \frac{1}{2}(0^-)$$

 Mass $m = 1864.83 \pm 0.05$ MeV

 $m_{D^\pm} - m_{D^0} = 4.822 \pm 0.015$ MeV

 Mean life $\tau = (410.1 \pm 1.5) \times 10^{-15}$ s

 $c\tau = 122.9$ μm

Mixing and related parameters

$$|m_{D_1^0} - m_{D_2^0}| = (0.95^{+0.41}_{-0.44}) \times 10^{10} \hbar \text{ s}^{-1}$$

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.29^{+0.14}_{-0.18}) \times 10^{-2}$$

$$|q/p| = 0.92^{+0.12}_{-0.09}$$

$$A_\Gamma = (-0.125 \pm 0.526) \times 10^{-3}$$

 $K^+ \pi^-$ relative strong phase: $\cos \delta = 0.97 \pm 0.11$
 $K^- \pi^+ \pi^0$ coherence factor $R_{K\pi\pi^0} = 0.82 \pm 0.06$
 $K^- \pi^+ \pi^0$ average relative strong phase $\delta^{K\pi\pi^0} = (199 \pm 14)^\circ$
 $K^- \pi^- 2\pi^+$ coherence factor $R_{K3\pi} = 0.53^{+0.18}_{-0.21}$
 $K^- \pi^- 2\pi^+$ average relative strong phase $\delta^{K3\pi} = (125^{+22}_{-14})^\circ$
 $D^0 \rightarrow K^- \pi^- 2\pi^+$, $R_{K3\pi} (y \cos \delta^{K3\pi} - x \sin \delta^{K3\pi}) = (-3.0 \pm 0.7) \times 10^{-3} \text{ TeV}^{-1}$
 $K_S^0 K^+ \pi^-$ coherence factor $R_{K_S^0 K\pi} = 0.70 \pm 0.08$
 $K_S^0 K^+ \pi^-$ average relative strong phase $\delta^{K_S^0 K\pi} = (0 \pm 16)^\circ$
 $K^* K$ coherence factor $R_{K^* K} = 0.94 \pm 0.12$
 $K^* K$ average relative strong phase $\delta^{K^* K} = (-17 \pm 18)^\circ$

CP-violation decay-rate asymmetries (labeled by the D^0 decay)

$$\begin{aligned}
A_{CP}(K^+ K^-) &= (-0.07 \pm 0.11)\% \\
A_{CP}(2K_S^0) &= (0.4 \pm 1.4)\% \\
A_{CP}(\pi^+ \pi^-) &= (0.13 \pm 0.14)\% \\
A_{CP}(\pi^0 \pi^0) &= (0.0 \pm 0.6)\% \\
A_{CP}(\rho \gamma) &= (6 \pm 15) \times 10^{-2} \\
A_{CP}(\phi \gamma) &= (-9 \pm 7) \times 10^{-2} \\
A_{CP}(\bar{K}^*(892)^0 \gamma) &= (-0.3 \pm 2.0) \times 10^{-2} \\
A_{CP}(\pi^+ \pi^- \pi^0) &= (0.3 \pm 0.4)\% \\
A_{CP}(\rho(770)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) &= (1.2 \pm 0.9)\% [i] \\
A_{CP}(\rho(770)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (-3.1 \pm 3.0)\% [i] \\
A_{CP}(\rho(770)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) &= (-1.0 \pm 1.7)\% [i] \\
A_{CP}(\rho(1450)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) &= (0 \pm 70)\% [i] \\
A_{CP}(\rho(1450)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (-20 \pm 40)\% [i] \\
A_{CP}(\rho(1450)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) &= (6 \pm 9)\% [i] \\
A_{CP}(\rho(1700)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) &= (-5 \pm 14)\% [i] \\
A_{CP}(\rho(1700)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (13 \pm 9)\% [i] \\
A_{CP}(\rho(1700)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) &= (8 \pm 11)\% [i] \\
A_{CP}(f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (0 \pm 35)\% [i] \\
A_{CP}(f_0(1370) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (25 \pm 18)\% [i] \\
A_{CP}(f_0(1500) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (0 \pm 18)\% [i] \\
A_{CP}(f_0(1710) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (0 \pm 24)\% [i] \\
A_{CP}(f_2(1270) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (-4 \pm 6)\% [i] \\
A_{CP}(\sigma(400) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) &= (6 \pm 8)\% [i] \\
A_{CP}(\text{nonresonant } \pi^+ \pi^- \pi^0) &= (-13 \pm 23)\% [i] \\
A_{CP}(a_1(1260)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) &= (5 \pm 6)\% \\
A_{CP}(a_1(1260)^- \pi^+ \rightarrow 2\pi^+ 2\pi^-) &= (14 \pm 18)\% \\
A_{CP}(\pi(1300)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) &= (-2 \pm 15)\% \\
A_{CP}(\pi(1300)^- \pi^+ \rightarrow 2\pi^+ 2\pi^-) &= (-6 \pm 30)\% \\
A_{CP}(a_1(1640)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) &= (9 \pm 26)\% \\
A_{CP}(\pi_2(1670)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) &= (7 \pm 18)\% \\
A_{CP}(\sigma f_0(1370) \rightarrow 2\pi^+ 2\pi^-) &= (-15 \pm 19)\% \\
A_{CP}(\sigma \rho(770)^0 \rightarrow 2\pi^+ 2\pi^-) &= (3 \pm 27)\% \\
A_{CP}(2\rho(770)^0 \rightarrow 2\pi^+ 2\pi^-) &= (-6 \pm 6)\% \\
A_{CP}(2f_2(1270) \rightarrow 2\pi^+ 2\pi^-) &= (-28 \pm 24)\% \\
A_{CP}(K^+ K^- \pi^0) &= (-1.0 \pm 1.7)\% \\
A_{CP}(K^*(892)^+ K^- \rightarrow K^+ K^- \pi^0) &= (-0.9 \pm 1.3)\% [i] \\
A_{CP}(K^*(1410)^+ K^- \rightarrow K^+ K^- \pi^0) &= (-21 \pm 24)\% [i] \\
A_{CP}((K^+ \pi^0)_{S\text{-wave}} K^- \rightarrow K^+ K^- \pi^0) &= (7 \pm 15)\% [i] \\
A_{CP}(\phi(1020) \pi^0 \rightarrow K^+ K^- \pi^0) &= (1.1 \pm 2.2)\% [i]
\end{aligned}$$

$$\begin{aligned}
A_{CP}(f_0(980)\pi^0 \rightarrow K^+K^-\pi^0) &= (-3 \pm 19)\% [i] \\
A_{CP}(a_0(980)^0\pi^0 \rightarrow K^+K^-\pi^0) &= (-5 \pm 16)\% [i] \\
A_{CP}(f'_2(1525)\pi^0 \rightarrow K^+K^-\pi^0) &= (0 \pm 160)\% [i] \\
A_{CP}(K^*(892)^-K^+ \rightarrow K^+K^-\pi^0) &= (-5 \pm 4)\% [i] \\
A_{CP}(K^*(1410)^-K^+ \rightarrow K^+K^-\pi^0) &= (-17 \pm 29)\% [i] \\
A_{CP}((K^-\pi^0)_{S\text{-wave}}K^+ \rightarrow K^+K^-\pi^0) &= (-10 \pm 40)\% [i] \\
A_{CP}(K_S^0\pi^0) &= (-0.20 \pm 0.17)\% \\
A_{CP}(K_S^0\eta) &= (0.5 \pm 0.5)\% \\
A_{CP}(K_S^0\eta') &= (1.0 \pm 0.7)\% \\
A_{CP}(K_S^0\phi) &= (-3 \pm 9)\% \\
A_{CP}(K^-\pi^+) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+\pi^-) &= (-0.9 \pm 1.4)\% \\
A_{CP}(D_{CP}(\pm 1) \rightarrow K^\mp\pi^\pm) &= (12.7 \pm 1.5)\% \\
A_{CP}(K^-\pi^+\pi^0) &= (0.1 \pm 0.5)\% \\
A_{CP}(K^+\pi^-\pi^0) &= (0 \pm 5)\% \\
A_{CP}(K_S^0\pi^+\pi^-) &= (-0.1 \pm 0.8)\% \\
A_{CP}(K^*(892)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (0.4 \pm 0.5)\% \\
A_{CP}(K^*(892)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (1 \pm 6)\% \\
A_{CP}(\bar{K}^0\rho^0 \rightarrow K_S^0\pi^+\pi^-) &= (-0.1 \pm 0.5)\% \\
A_{CP}(\bar{K}^0\omega \rightarrow K_S^0\pi^+\pi^-) &= (-13 \pm 7)\% \\
A_{CP}(\bar{K}^0f_0(980) \rightarrow K_S^0\pi^+\pi^-) &= (-0.4 \pm 2.7)\% \\
A_{CP}(\bar{K}^0f_2(1270) \rightarrow K_S^0\pi^+\pi^-) &= (-4 \pm 5)\% \\
A_{CP}(\bar{K}^0f_0(1370) \rightarrow K_S^0\pi^+\pi^-) &= (-1 \pm 9)\% \\
A_{CP}(\bar{K}^0\rho^0(1450) \rightarrow K_S^0\pi^+\pi^-) &= (-4 \pm 10)\% \\
A_{CP}(\bar{K}^0f_0(600) \rightarrow K_S^0\pi^+\pi^-) &= (-3 \pm 5)\% \\
A_{CP}(K^*(1410)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (-2 \pm 9)\% \\
A_{CP}(K_0^*(1430)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (4 \pm 4)\% \\
A_{CP}(K_0^*(1430)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (12 \pm 15)\% \\
A_{CP}(K_2^*(1430)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (3 \pm 6)\% \\
A_{CP}(K_2^*(1430)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (-10 \pm 32)\% \\
A_{CP}(K^-\pi^+\pi^+\pi^-) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+\pi^-\pi^+\pi^-) &= (-2 \pm 4)\% \\
A_{CP}(K^+K^-\pi^+\pi^-) &= (1.3 \pm 1.7)\% \\
A_{CP}(K_1^*(1270)^+K^- \rightarrow K^+K^-\pi^+\pi^-) &= (25 \pm 16)\% \\
A_{CP}(K_1^*(1270)^+K^- \rightarrow K^{*0}\pi^+K^-) &= (-1 \pm 10)\% \\
A_{CP}(K_1^*(1270)^-K^+ \rightarrow \bar{K}^{*0}\pi^-K^+) &= (-10 \pm 32)\% \\
A_{CP}(K_1^*(1270)^-K^+ \rightarrow K^+K^-\pi^+\pi^-) &= (-50 \pm 20)\% \\
A_{CP}(K_1^*(1270)^+K^- \rightarrow \rho^0K^+K^-) &= (-7 \pm 17)\% \\
A_{CP}(K_1^*(1270)^-K^+ \rightarrow \rho^0K^-K^+) &= (10 \pm 13)\% \\
A_{CP}(K_1^*(1400)^+K^- \rightarrow K^+K^-\pi^+\pi^-) &= (9 \pm 25)\% \\
A_{CP}(K^*(1410)^+K^- \rightarrow K^{*0}\pi^+K^-) &= (-20 \pm 17)\% \\
A_{CP}(K^*(1410)^-K^+ \rightarrow \bar{K}^{*0}\pi^-K^+) &= (-1 \pm 14)\%
\end{aligned}$$

$$A_{CP}(K^*(1680)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) = (-17 \pm 29)\%$$

$$A_{CP}(K^{*0} \bar{K}^{*0}) \text{ in } D^0, \bar{D}^0 \rightarrow K^{*0} \bar{K}^{*0} = (-5 \pm 14)\%$$

$$A_{CP}(K^{*0} \bar{K}^{*0} \text{ S-wave}) = (10 \pm 14)\%$$

$$A_{CP}(\phi \rho^0) \text{ in } D^0, \bar{D}^0 \rightarrow \phi \rho^0 = (1 \pm 9)\%$$

$$A_{CP}(\phi \rho^0 \text{ S-wave}) = (-3 \pm 5)\%$$

$$A_{CP}(\phi \rho^0 \text{ D-wave}) = (-37 \pm 19)\%$$

$$A_{CP}(\phi(\pi^+ \pi^-)_{\text{S-wave}}) = (0 \pm 50)\%$$

$$A_{CP}(K^*(892)^0 (K^- \pi^+)_{\text{S-wave}}) = (-10 \pm 40)\%$$

$$A_{CP}(K^+ K^- \pi^+ \pi^- \text{ non-resonant}) = (8 \pm 20)\%$$

$$A_{CP}((K^- \pi^+)_{\text{P-wave}} (K^+ \pi^-)_{\text{S-wave}}) = (3 \pm 11)\%$$

$$A_{CP}(K^+ K^- \mu^+ \mu^-) \text{ in } D^0, \bar{D}^0 \rightarrow K^+ K^- \mu^+ \mu^- = (0 \pm 11)\%$$

$$A_{CP}(\pi^+ \pi^- \mu^+ \mu^-) \text{ in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \mu^+ \mu^- = (5 \pm 4)\%$$

CP-even fractions (labeled by the D^0 decay)

$$\text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^0 \text{ decays} = (97.3 \pm 1.7)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow K^+ K^- \pi^0 \text{ decays} = (73 \pm 6)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- \text{ decays} = (76.9 \pm 2.3)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0 \text{ decays} = (23.8 \pm 1.7)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow K^+ K^- \pi^+ \pi^- \text{ decays} = (75 \pm 4)\%$$

CP-violation asymmetry difference

$$\Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-0.161 \pm 0.029)\%$$

χ^2 tests of CP-violation (CPV) p-values

$$\text{Local CPV in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \pi^0 = 4.9\%$$

$$\text{Local CPV in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- = (0.6 \pm 0.2)\%$$

$$\text{Local CPV in } D^0, \bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^- = 96\%$$

$$\text{Local CPV in } D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^0 = 16.6\%$$

$$\text{Local CPV in } D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^+ \pi^- = 9.1\%$$

T-violation decay-rate asymmetry

$$A_T(K^+ K^- \pi^+ \pi^-) = (1.7 \pm 2.7) \times 10^{-3} [b]$$

$$A_{T\text{viol}}(K_S \pi^+ \pi^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow K_S \pi^+ \pi^- \pi^0 = (-0.3_{-1.6}^{+1.4}) \times 10^{-3}$$

CPT-violation decay-rate asymmetry

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$r_V \equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 1.7 \pm 0.8$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 0.9 \pm 0.4$$

$$f_+(0) \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.736 \pm 0.004$$

$$f_+(0) |V_{cs}| \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.7166 \pm 0.0030$$

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = -2.40 \pm 0.16$$

$$\begin{aligned}
 r_2 \equiv a_2/a_0 \text{ in } D^0 &\rightarrow K^- \ell^+ \nu_\ell = 5 \pm 4 \\
 f_+(0) \text{ in } D^0 &\rightarrow \pi^- \ell^+ \nu_\ell = 0.637 \pm 0.009 \\
 f_+(0) |V_{cd}| \text{ in } D^0 &\rightarrow \pi^- \ell^+ \nu_\ell = 0.1436 \pm 0.0026 \quad (S = 1.5) \\
 r_1 \equiv a_1/a_0 \text{ in } D^0 &\rightarrow \pi^- \ell^+ \nu_\ell = -1.97 \pm 0.28 \quad (S = 1.4) \\
 r_2 \equiv a_1/a_0 \text{ in } D^0 &\rightarrow \pi^- \ell^+ \nu_\ell = -0.2 \pm 2.2 \quad (S = 1.7)
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level(MeV/c)	p
Topological modes			
0-prongs	[j] (15 ± 6) %		—
2-prongs	(71 ± 6) %		—
4-prongs	[k] (14.6 ± 0.5) %		—
6-prongs	[l] (6.5 ± 1.3) × 10 ⁻⁴		—
Inclusive modes			
e^+ anything	[n] (6.49 ± 0.11) %		—
μ^+ anything	(6.8 ± 0.6) %		—
K^- anything	(54.7 ± 2.8) %	S=1.3	—
\bar{K}^0 anything + K^0 anything	(47 ± 4) %		—
K^+ anything	(3.4 ± 0.4) %		—
$K^*(892)^-$ anything	(15 ± 9) %		—
$\bar{K}^*(892)^0$ anything	(9 ± 4) %		—
$K^*(892)^+$ anything	< 3.6 %	CL=90%	—
$K^*(892)^0$ anything	(2.8 ± 1.3) %		—
η anything	(9.5 ± 0.9) %		—
η' anything	(2.48 ± 0.27) %		—
ϕ anything	(1.05 ± 0.11) %		—
invisibles	< 9.4 × 10 ⁻⁵	CL=90%	—
Semileptonic modes			
$K^- e^+ \nu_e$	(3.542 ± 0.035) %	S=1.3	867
$K^- \mu^+ \nu_\mu$	(3.41 ± 0.04) %		864
$K^*(892)^- e^+ \nu_e$	(2.15 ± 0.16) %		719
$K^*(892)^- \mu^+ \nu_\mu$	(1.89 ± 0.24) %		714
$K^- \pi^0 e^+ \nu_e$	(1.6 ± 1.3 / 0.5) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$	(2.7 ± 0.9 / 0.7) %		860
$K^- \pi^+ \pi^- e^+ \nu_e$	(2.8 ± 1.4 / 1.1) × 10 ⁻⁴		843

$K_1(1270)^- e^+ \nu_e$	$(7.6 \begin{smallmatrix} + \\ - \end{smallmatrix} 4.0 \\ 3.1) \times 10^{-4}$		498
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.3	$\times 10^{-3}$	CL=90% 821
$(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$	< 1.5	$\times 10^{-3}$	CL=90% 692
$\pi^- e^+ \nu_e$	$(2.91 \pm 0.04) \times 10^{-3}$		927
$\pi^- \mu^+ \nu_\mu$	$(2.67 \pm 0.12) \times 10^{-3}$	S=1.3	924
$\rho^- e^+ \nu_e$	$(1.77 \pm 0.16) \times 10^{-3}$		771
$a(980)^- e^+ \nu_e, a^- \rightarrow \eta \pi^-$	$(1.33 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.34 \\ 0.30) \times 10^{-4}$		–
Hadronic modes with one \bar{K}			
$K^- \pi^+$	$(3.950 \pm 0.031) \%$	S=1.2	861
$K_S^0 \pi^0$	$(1.240 \pm 0.022) \%$		860
$K_L^0 \pi^0$	$(10.0 \pm 0.7) \times 10^{-3}$		860
$K_S^0 \pi^+ \pi^-$	[c] $(2.80 \pm 0.18) \%$	S=1.1	842
$K_S^0 \rho^0$	$(6.3 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.6 \\ 0.8) \times 10^{-3}$		674
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	$(2.0 \pm 0.6) \times 10^{-4}$		670
$K_S^0 (\pi^+ \pi^-)_{S\text{-wave}}$	$(3.3 \pm 0.8) \times 10^{-3}$		842
$K_S^0 f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$(1.20 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.40 \\ 0.23) \times 10^{-3}$		549
$K_S^0 f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	$(2.8 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.9 \\ 1.3) \times 10^{-3}$		†
$K_S^0 f_2(1270), f_2 \rightarrow \pi^+ \pi^-$	$(9 \begin{smallmatrix} + \\ - \end{smallmatrix} 10 \\ 6) \times 10^{-5}$		262
$K^*(892)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(1.64 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.14 \\ 0.17) \%$		711
$K_0^*(1430)^- \pi^+, K_0^{*-} \rightarrow K_S^0 \pi^-$	$(2.67 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.40 \\ 0.33) \times 10^{-3}$		378
$K_2^*(1430)^- \pi^+, K_2^{*-} \rightarrow K_S^0 \pi^-$	$(3.4 \begin{smallmatrix} + \\ - \end{smallmatrix} 1.9 \\ 1.0) \times 10^{-4}$		367
$K^*(1680)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(4.4 \pm 3.5) \times 10^{-4}$		46
$K^*(892)^+ \pi^-, K^{*+} \rightarrow K_S^0 \pi^+$	[o] $(1.13 \begin{smallmatrix} + \\ - \end{smallmatrix} 0.60 \\ 0.34) \times 10^{-4}$		711
$K_0^*(1430)^+ \pi^-, K_0^{*+} \rightarrow K_S^0 \pi^+$	[o] < 1.4	$\times 10^{-5}$	CL=95% –
$K_2^*(1430)^+ \pi^-, K_2^{*+} \rightarrow K_S^0 \pi^+$	[o] < 3.4	$\times 10^{-5}$	CL=95% –
$K_S^0 \pi^+ \pi^-$ nonresonant	$(2.5 \begin{smallmatrix} + \\ - \end{smallmatrix} 6.0 \\ 1.6) \times 10^{-4}$		842
$K^- \pi^+ \pi^0$	[c] $(14.4 \pm 0.5) \%$	S=2.0	844
$K^- \rho^+$	$(11.3 \pm 0.7) \%$		675
$K^- \rho(1700)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(8.2 \pm 1.8) \times 10^{-3}$		†

$K^*(892)^- \pi^+$, $K^*(892)^- \rightarrow$	$(2.31 \pm_{-0.20}^{+0.40}) \%$		711
$\bar{K}^*(892)^0 \pi^0$, $\bar{K}^*(892)^0 \rightarrow$	$(1.95 \pm 0.24) \%$		711
$K_0^*(1430)^- \pi^+$, $K_0^{*-} \rightarrow$	$(4.8 \pm 2.2) \times 10^{-3}$		378
$\bar{K}_0^*(1430)^0 \pi^0$, $\bar{K}_0^{*0} \rightarrow$	$(5.9 \pm_{-1.6}^{+5.0}) \times 10^{-3}$		379
$K^*(1680)^- \pi^+$, $K^{*-} \rightarrow$	$(1.9 \pm 0.7) \times 10^{-3}$		46
$K^- \pi^+ \pi^0$ nonresonant	$(1.15 \pm_{-0.20}^{+0.60}) \%$		844
$K_S^0 2\pi^0$	$(9.1 \pm 1.1) \times 10^{-3}$	S=2.2	843
$K_S^0 (2\pi^0)_{S-wave}$	$(2.6 \pm 0.7) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \pi^0$, $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(8.1 \pm 0.7) \times 10^{-3}$		711
$\bar{K}^*(1430)^0 \pi^0$, $\bar{K}^{*0} \rightarrow$	$(4 \pm_{23}) \times 10^{-5}$		—
$\bar{K}^*(1680)^0 \pi^0$, $\bar{K}^{*0} \rightarrow$	$(1.0 \pm 0.4) \times 10^{-3}$		—
$K_S^0 f_2(1270)$, $f_2 \rightarrow 2\pi^0$	$(2.3 \pm 1.1) \times 10^{-4}$		—
$2K_S^0$, one $K_S^0 \rightarrow 2\pi^0$	$(3.2 \pm 1.1) \times 10^{-4}$		—
$K^- 2\pi^+ \pi^-$	[c] $(8.23 \pm 0.14) \%$	S=1.1	813
$K^- \pi^+ \rho^0$ total	$(6.87 \pm 0.31) \%$		609
$K^- \pi^+ \rho^0$ 3-body	$(6.1 \pm 1.6) \times 10^{-3}$		609
$(K^- \pi^+)_{S-wave} \rho^0$	$(7.7 \pm 1.1) \times 10^{-4}$		609
$\bar{K}^*(892)^0 \rho^0$, $\bar{K}^{*0} \rightarrow$	$(1.01 \pm 0.05) \%$		416
$\bar{K}^*(892)^0 (\pi^+ \pi^-)_{S-wave}$,	$(1.9 \pm 0.4) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$(\bar{K}^*(892)^0 \rho^0)_{S-wave}$,	$(6.0 \pm 0.4) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$(\bar{K}^*(892)^0 \rho^0)_{P-wave}$,	$(4.96 \pm 0.25) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$(\bar{K}^*(892)^0 \rho^0)_{D-wave}$,	$(6.8 \pm 0.4) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(892)^0 \rho^0$ transverse,	$(1.2 \pm 0.4) \%$		417
$\bar{K}^{*0} \rightarrow K^- \pi^+$			
$(\bar{K}^*(892)^0 \rho(1450)^0)$	$(5.0 \pm 1.8) \times 10^{-4}$		—
$S-wave$, $\bar{K}^*(892)^0 \rightarrow$			
$K^- \pi^+$, $\rho(1450)^0 \rightarrow$			
$\pi^+ \pi^-$			

$(\bar{K}^*(892)^0 \rho(1450)^0)$	$(1.63 \pm 0.28) \times 10^{-3}$	—
ρ -wave, $\bar{K}^*(892)^0 \rightarrow$		
$K^- \pi^+$, $\rho(1450)^0 \rightarrow$		
$\pi^+ \pi^-$		
$(\bar{K}^*(892)^0 \rho(1450)^0)$	$(3.8 \pm 1.3) \times 10^{-4}$	—
D -wave, $\bar{K}^*(892)^0 \rightarrow$		
$K^- \pi^+$, $\rho(1450)^0 \rightarrow$		
$\pi^+ \pi^-$		
$K^- a_1(1260)^+$, $a_1^+ \rightarrow$	$(3.13 \pm 0.21) \%$	—
$2\pi^+ \pi^-$ total		
$K^- a_1(1260)^+$, $a_1^+ \rightarrow$	$(4.33 \pm 0.32) \%$	327
$\rho^0 \pi^+$		
$K^- a_1(1260)^+$,	$(4.4 \pm 0.4) \%$	—
$a_1(1260)^+ \rightarrow$		
$(\rho^0 \pi^+)_{S\text{-wave}}$		
$K^- a_1(1260)^+$,	$(2.5 \pm 1.2) \times 10^{-4}$	—
$a_1(1260)^+ \rightarrow$		
$(\rho^0 \pi^+)_{D\text{-wave}}$		
$K_1(1270)^- \pi^+$, $K_1^- \rightarrow$	$(3.9 \pm 0.4) \times 10^{-3}$	—
$\frac{K^- \pi^+ \pi^- \text{ total}}{\bar{K}^*(892)^0 \pi^+ \pi^-}$ 3-body,	$(6.0 \pm 0.5) \times 10^{-3}$	685
$\bar{K}^{*0} \rightarrow K^- \pi^+$		
$K_1(1270)^- \pi^+$, $K_1^- \rightarrow$	$(6.6 \pm 2.3) \times 10^{-4}$	484
$\bar{K}^*(892)^0 \pi^-$, $\bar{K}^{*0} \rightarrow$		
$K^- \pi^+$		
$K_1(1270)^- \pi^+$,	$(8 \pm 12) \times 10^{-5}$	—
$K_1(1270)^- \rightarrow$		
$(\bar{K}^{*0} \pi^-)_{S\text{-wave}}$,		
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K_1(1270)^- \pi^+$,	$(5.8 \pm 2.3) \times 10^{-4}$	—
$K_1(1270)^- \rightarrow$		
$(\bar{K}^{*0} \pi^-)_{D\text{-wave}}$,		
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K_1(1270)^- \pi^+$,	$(2.8 \pm 0.5) \times 10^{-3}$	—
$K_1(1270)^- \rightarrow$		
$(K^- \rho^0)_{S\text{-wave}}$		
$K_1(1400)^- \pi^+$,	$(9.5 \pm 1.9) \times 10^{-4}$	—
$K_1(1400)^- \rightarrow$		
$(\bar{K}^*(892)^0 \pi^-)_{S\text{-wave}}$,		
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K_2^*(1430)^- \pi^+$,	$(3.8 \pm 0.4) \times 10^{-4}$	—
$K_2^*(1430)^- \rightarrow$		
$\bar{K}^*(892)^0 \pi^-$,		
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		

$K(1460)^- \pi^+$, $K(1460)^- \rightarrow$	$(3.1 \pm 0.4) \times 10^{-3}$	–
$K^- \pi^+ \pi^-$ total		
$K^- 2\pi^+ \pi^-$ nonresonant	$(1.81 \pm 0.07) \%$	813
$K_S^0 \pi^+ \pi^- \pi^0$	[p] $(5.2 \pm 0.6) \%$	813
$K_S^0 \eta$, $\eta \rightarrow \pi^+ \pi^- \pi^0$	$(1.17 \pm 0.03) \times 10^{-3}$	772
$K_S^0 \omega$, $\omega \rightarrow \pi^+ \pi^- \pi^0$	$(9.9 \pm 0.6) \times 10^{-3}$	670
$K^- 2\pi^+ \pi^- \pi^0$	$(4.3 \pm 0.4) \%$	771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$, $\bar{K}^{*0} \rightarrow$	$(1.3 \pm 0.6) \%$	643
$K^- \pi^+$		
$K^- \pi^+ \omega$, $\omega \rightarrow \pi^+ \pi^- \pi^0$	$(2.8 \pm 0.5) \%$	605
$\bar{K}^*(892)^0 \omega$, $\bar{K}^{*0} \rightarrow$	$(6.5 \pm 3.0) \times 10^{-3}$	410
$K^- \pi^+$, $\omega \rightarrow$		
$\pi^+ \pi^- \pi^0$		
$K_S^0 \eta \pi^0$	$(5.7 \pm 1.1) \times 10^{-3}$	721
$K_S^0 a_0(980)$, $a_0 \rightarrow \eta \pi^0$	$(6.8 \pm 2.1) \times 10^{-3}$	–
$\bar{K}^*(892)^0 \eta$, $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(1.7 \pm 0.5) \times 10^{-3}$	–
$K_S^0 2\pi^+ 2\pi^-$	$(2.66 \pm 0.30) \times 10^{-3}$	768
$K_S^0 \rho^0 \pi^+ \pi^-$, no $K^*(892)^-$	$(1.1 \pm 0.7) \times 10^{-3}$	–
$K^*(892)^- 2\pi^+ \pi^-$,	$(5 \pm 7) \times 10^{-4}$	642
$K^*(892)^- \rightarrow K_S^0 \pi^-$,		
no ρ^0		
$K^*(892)^- \rho^0 \pi^+$,	$(1.6 \pm 0.6) \times 10^{-3}$	230
$K^*(892)^- \rightarrow K_S^0 \pi^-$		
$K_S^0 2\pi^+ 2\pi^-$ nonresonant	$< 1.2 \times 10^{-3}$	CL=90% 768
$K^- 3\pi^+ 2\pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	713

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. These nine modes below are all corrected for unseen decays of the resonances.

$K_S^0 \eta$	$(5.09 \pm 0.13) \times 10^{-3}$	772
$K_S^0 \omega$	$(1.11 \pm 0.06) \%$	670
$K_S^0 \eta'(958)$	$(9.49 \pm 0.32) \times 10^{-3}$	565
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	$(1.9 \pm 0.9) \%$	643
$K^- \pi^+ \omega$	$(3.1 \pm 0.6) \%$	605
$\bar{K}^*(892)^0 \omega$	$(1.1 \pm 0.5) \%$	410
$K^- \pi^+ \eta'(958)$	$(6.43 \pm 0.34) \times 10^{-3}$	479
$K_S^0 \eta'(958) \pi^0$	$(2.52 \pm 0.27) \times 10^{-3}$	479
$\bar{K}^*(892)^0 \eta'(958)$	$< 1.0 \times 10^{-3}$	CL=90% 119

Hadronic modes with three K 's

$K_S^0 K^+ K^-$	$(4.42 \pm 0.32) \times 10^{-3}$	544
$K_S^0 a_0(980)^0$, $a_0^0 \rightarrow K^+ K^-$	$(2.9 \pm 0.4) \times 10^{-3}$	–
$K^- a_0(980)^+$, $a_0^+ \rightarrow$	$(5.9 \pm 1.8) \times 10^{-4}$	–
$K^+ K_S^0$		

$K^+ a_0(980)^-, a_0^- \rightarrow$	< 1.1	$\times 10^{-4}$	CL=95%	–
$K^- K_S^0$				
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	< 9	$\times 10^{-5}$	CL=95%	–
$K_S^0 \phi, \phi \rightarrow K^+ K^-$	(2.03 ± 0.15)	$\times 10^{-3}$		520
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	(1.7 ± 1.1)	$\times 10^{-4}$		–
$3K_S^0$	(7.5 ± 0.7)	$\times 10^{-4}$	S=1.4	539
$K^+ 2K^- \pi^+$	(2.25 ± 0.32)	$\times 10^{-4}$		434
$K^+ K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$	(4.5 ± 1.8)	$\times 10^{-5}$		†
$K^- \pi^+$				
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	(4.1 ± 1.7)	$\times 10^{-5}$		422
$\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-,$	(1.08 ± 0.21)	$\times 10^{-4}$		†
$\bar{K}^{*0} \rightarrow K^- \pi^+$				
$K^+ 2K^- \pi^+$ nonresonant	(3.4 ± 1.5)	$\times 10^{-5}$		434
$2K_S^0 K^\pm \pi^\mp$	(5.9 ± 1.3)	$\times 10^{-4}$		427

Pionic modes

$\pi^+ \pi^-$	(1.455 ± 0.024)	$\times 10^{-3}$	S=1.3	922
$2\pi^0$	(8.26 ± 0.25)	$\times 10^{-4}$		923
$\pi^+ \pi^- \pi^0$	(1.49 ± 0.06)	%	S=2.1	907
$\rho^+ \pi^-$	(1.01 ± 0.04)	%		764
$\rho^0 \pi^0$	(3.86 ± 0.23)	$\times 10^{-3}$		764
$\rho^- \pi^+$	(5.15 ± 0.25)	$\times 10^{-3}$		764
$\rho(1450)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	(1.6 ± 2.1)	$\times 10^{-5}$		–
$\rho(1450)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	(4.5 ± 2.0)	$\times 10^{-5}$		–
$\rho(1450)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	(2.7 ± 0.4)	$\times 10^{-4}$		–
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	(6.1 ± 1.5)	$\times 10^{-4}$		–
$\rho(1700)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	(7.4 ± 1.8)	$\times 10^{-4}$		–
$\rho(1700)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	(4.8 ± 1.1)	$\times 10^{-4}$		–
$f_0(980) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	(3.7 ± 0.9)	$\times 10^{-5}$		–
$f_0(500) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	(1.22 ± 0.22)	$\times 10^{-4}$		–
$f_0(1370) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	(5.5 ± 2.1)	$\times 10^{-5}$		–
$f_0(1500) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	(5.8 ± 1.6)	$\times 10^{-5}$		–
$f_0(1710) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	(4.6 ± 1.6)	$\times 10^{-5}$		–
$f_2(1270) \pi^0, f_2 \rightarrow \pi^+ \pi^-$	(1.97 ± 0.21)	$\times 10^{-4}$		–
$\pi^+ \pi^- \pi^0$ nonresonant	(1.3 ± 0.4)	$\times 10^{-4}$		907
$3\pi^0$	(2.0 ± 0.5)	$\times 10^{-4}$		908
$2\pi^+ 2\pi^-$	(7.56 ± 0.20)	$\times 10^{-3}$		880
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	(4.54 ± 0.31)	$\times 10^{-3}$		–
$2\pi^+ \pi^-$ total				
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	(3.14 ± 0.21)	$\times 10^{-3}$		–
$\rho^0 \pi^+$ S-wave				
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	(1.9 ± 0.5)	$\times 10^{-4}$		–
$\rho^0 \pi^+$ D-wave				

$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	$(6.4 \pm 0.7) \times 10^{-4}$	—
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \rho^0 \pi^-$ <i>S-wave</i>	$(2.3 \pm 0.9) \times 10^{-4}$	—
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \sigma \pi^-$	$(6.1 \pm 3.4) \times 10^{-5}$	—
$\pi(1300)^+ \pi^-$, $\pi(1300)^+ \rightarrow \sigma \pi^+$	$(5.1 \pm 2.7) \times 10^{-4}$	—
$\pi(1300)^- \pi^+$, $\pi(1300)^- \rightarrow \sigma \pi^-$	$(2.3 \pm 2.2) \times 10^{-4}$	—
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+$ <i>D-wave</i>	$(3.2 \pm 1.6) \times 10^{-4}$	—
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	$(1.8 \pm 1.4) \times 10^{-4}$	—
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow f_2(1270)^0 \pi^+$, $f_2^0 \rightarrow \pi^+ \pi^-$	$(2.0 \pm 0.9) \times 10^{-4}$	—
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow \sigma \pi^+$	$(2.6 \pm 1.0) \times 10^{-4}$	—
$2\rho^0$ total	$(1.85 \pm 0.13) \times 10^{-3}$	518
$2\rho^0$, parallel helicities	$(8.3 \pm 3.2) \times 10^{-5}$	—
$2\rho^0$, perpendicular helicities	$(4.8 \pm 0.6) \times 10^{-4}$	—
$2\rho^0$, longitudinal helicities	$(1.27 \pm 0.10) \times 10^{-3}$	—
$2\rho(770)^0$, <i>S-wave</i>	$(1.8 \pm 1.3) \times 10^{-4}$	—
$2\rho(770)^0$, <i>P-wave</i>	$(5.3 \pm 1.3) \times 10^{-4}$	—
$2\rho(770)^0$, <i>D-wave</i>	$(6.2 \pm 3.0) \times 10^{-4}$	—
Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$ 3-body total	$(1.51 \pm 0.12) \times 10^{-3}$	—
$\sigma \pi^+ \pi^-$	$(6.2 \pm 0.9) \times 10^{-4}$	—
$\sigma \rho(770)^0$	$(5.0 \pm 2.5) \times 10^{-4}$	—
$f_0(980) \pi^+ \pi^-$, $f_0 \rightarrow \pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-4}$	—
$f_2(1270) \pi^+ \pi^-$, $f_2 \rightarrow \pi^+ \pi^-$	$(3.7 \pm 0.6) \times 10^{-4}$	—
$2f_2(1270)$, $f_2 \rightarrow \pi^+ \pi^-$	$(1.6 \pm 1.8) \times 10^{-4}$	—
$f_0(1370) \sigma$, $f_0 \rightarrow \pi^+ \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	—
$\pi^+ \pi^- 2\pi^0$	$(1.02 \pm 0.09) \%$	882
$\eta \pi^0$ [q]	$(6.3 \pm 0.6) \times 10^{-4}$	S=1.1 846
$\omega \pi^0$ [q]	$(1.17 \pm 0.35) \times 10^{-4}$	761
$\omega \eta$	$(1.98 \pm 0.18) \times 10^{-3}$	S=1.1 648
$2\pi^+ 2\pi^- \pi^0$	$(4.2 \pm 0.5) \times 10^{-3}$	844
$\eta \pi^+ \pi^-$ [q]	$(1.09 \pm 0.16) \times 10^{-3}$	827
$\omega \pi^+ \pi^-$ [q]	$(1.6 \pm 0.5) \times 10^{-3}$	738
$\eta 2\pi^0$	$(3.8 \pm 1.3) \times 10^{-4}$	829
$3\pi^+ 3\pi^-$	$(4.3 \pm 1.2) \times 10^{-4}$	795
$\eta'(958) \pi^0$	$(9.2 \pm 1.0) \times 10^{-4}$	678

$\eta'(958)\pi^+\pi^-$	$(4.5 \pm 1.7) \times 10^{-4}$		650
2η	$(2.11 \pm 0.19) \times 10^{-3}$	S=2.2	754
$2\eta\pi^0$	$(7.3 \pm 2.2) \times 10^{-4}$		699
3η	$< 1.3 \times 10^{-4}$	CL=90%	421
$\eta\eta'(958)$	$(1.01 \pm 0.19) \times 10^{-3}$		537

Hadronic modes with a $K\bar{K}$ pair

K^+K^-	$(4.08 \pm 0.06) \times 10^{-3}$	S=1.6	791
$2K_S^0$	$(1.41 \pm 0.05) \times 10^{-4}$	S=1.1	789
$K_S^0K^-\pi^+$	$(3.3 \pm 0.5) \times 10^{-3}$	S=1.1	739
$\bar{K}^*(892)^0K_S^0, \bar{K}^{*0} \rightarrow$ $K^-\pi^+$	$(8.2 \pm 1.6) \times 10^{-5}$		608
$K^*(892)^+K^-, K^{*+} \rightarrow$ $K_S^0\pi^+$	$(1.89 \pm 0.30) \times 10^{-3}$		—
$\bar{K}^*(1410)^0K_S^0, \bar{K}^{*0} \rightarrow$ $K^-\pi^+$	$(1.3 \pm 1.9) \times 10^{-4}$		—
$K^*(1410)^+K^-, K^{*+} \rightarrow$ $K_S^0\pi^+$	$(3.2 \pm 1.9) \times 10^{-4}$		—
$(K^-\pi^+)_{S\text{-wave}}K_S^0$	$(6.0 \pm 2.9) \times 10^{-4}$		739
$(K_S^0\pi^+)_{S\text{-wave}}K^-$	$(3.9 \pm 1.0) \times 10^{-4}$		739
$a_0(980)^-\pi^+, a_0^- \rightarrow K_S^0K^-$	$(1.3 \pm 1.4) \times 10^{-4}$		—
$a_0(1450)^-\pi^+, a_0^- \rightarrow$ $K_S^0K^-$	$(2.5 \pm 2.0) \times 10^{-5}$		—
$a_2(1320)^-\pi^+, a_2^- \rightarrow$ $K_S^0K^-$	$(5 \pm 5) \times 10^{-6}$		—
$\rho(1450)^-\pi^+, \rho^- \rightarrow K_S^0K^-$	$(4.6 \pm 2.5) \times 10^{-5}$		—
$K_S^0K^+\pi^-$	$(2.17 \pm 0.34) \times 10^{-3}$	S=1.1	739
$K^*(892)^0K_S^0, K^{*0} \rightarrow$ $K^+\pi^-$	$(1.12 \pm 0.21) \times 10^{-4}$		608
$K^*(892)^-K^+, K^{*-} \rightarrow$ $K_S^0\pi^-$	$(6.2 \pm 1.0) \times 10^{-4}$		—
$K^*(1410)^0K_S^0, K^{*0} \rightarrow$ $K^+\pi^+$	$(5 \pm 8) \times 10^{-5}$		—
$K^*(1410)^-K^+, K^{*-} \rightarrow$ $K_S^0\pi^-$	$(2.6 \pm 2.0) \times 10^{-4}$		—
$(K^+\pi^-)_{S\text{-wave}}K_S^0$	$(3.7 \pm 1.9) \times 10^{-4}$		739
$(K_S^0\pi^-)_{S\text{-wave}}K^+$	$(1.4 \pm 0.6) \times 10^{-4}$		739
$a_0(980)^+\pi^-, a_0^+ \rightarrow K_S^0K^+$	$(6 \pm 4) \times 10^{-4}$		—
$a_0(1450)^+\pi^-, a_0^+ \rightarrow$ $K_S^0K^+$	$(3.2 \pm 2.5) \times 10^{-5}$		—
$\rho(1700)^+\pi^-, \rho^+ \rightarrow K_S^0K^+$	$(1.1 \pm 0.6) \times 10^{-5}$		—
$K^+K^-\pi^0$	$(3.42 \pm 0.14) \times 10^{-3}$		743

$K^*(892)^+ K^-$, $K^*(892)^+ \rightarrow K^+ \pi^0$	$(1.52 \pm 0.07) \times 10^{-3}$	—
$K^*(892)^- K^+$, $K^*(892)^- \rightarrow K^- \pi^0$	$(5.4 \pm 0.4) \times 10^{-4}$	—
$(K^+ \pi^0)_{S\text{-wave}} K^-$	$(2.43 \pm 0.18) \times 10^{-3}$	743
$(K^- \pi^0)_{S\text{-wave}} K^+$	$(1.3 \pm 0.5) \times 10^{-4}$	743
$f_0(980) \pi^0$, $f_0 \rightarrow K^+ K^-$	$(3.6 \pm 0.6) \times 10^{-4}$	—
$\phi \pi^0$, $\phi \rightarrow K^+ K^-$	$(6.6 \pm 0.4) \times 10^{-4}$	—
$2K_S^0 \pi^0$	$< 5.9 \times 10^{-4}$	740
$K^+ K^- \pi^+ \pi^-$	$(2.47 \pm 0.11) \times 10^{-3}$	677
$\phi(\pi^+ \pi^-)_{S\text{-wave}}$, $\phi \rightarrow K^+ K^-$	$(10 \pm 5) \times 10^{-5}$	614
$(\phi \rho^0)_{S\text{-wave}}$, $\phi \rightarrow K^+ K^-$	$(6.9 \pm 0.6) \times 10^{-4}$	250
$(\phi \rho^0)_{P\text{-wave}}$, $\phi \rightarrow K^+ K^-$	$(4.0 \pm 1.9) \times 10^{-5}$	—
$(\phi \rho^0)_{D\text{-wave}}$, $\phi \rightarrow K^+ K^-$	$(4.2 \pm 1.4) \times 10^{-5}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{S\text{-wave}}$, $K^{*0} \rightarrow K^\pm \pi^\mp$	$(1.1 \pm 0.5) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{P\text{-wave}}$, $K^* \rightarrow K^\pm \pi^\mp$	$(9 \pm 4) \times 10^{-5}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{D\text{-wave}}$, $K^* \rightarrow K^\pm \pi^\mp$	$(9.9 \pm 2.3) \times 10^{-5}$	—
$K^*(892)^0 (K^- \pi^+)_{S\text{-wave}}$ 3-body, $K^{*0} \rightarrow K^+ \pi^-$	$(1.4 \pm 0.6) \times 10^{-4}$	—
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow K^{*0} \pi^+$	$(1.4 \pm 0.9) \times 10^{-4}$	—
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow K^*(1430)^0 \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(1.5 \pm 0.5) \times 10^{-4}$	—
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow \rho^0 K^+$	$(2.2 \pm 0.6) \times 10^{-4}$	—
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow \omega(782) K^+$, $\omega \rightarrow \pi^+ \pi^-$	$(1.5 \pm 1.2) \times 10^{-5}$	—
$K_1(1270)^- K^+$, $K_1^- \rightarrow \rho^0 K^-$	$(1.3 \pm 0.4) \times 10^{-4}$	—
$K_1(1400)^+ K^-$, $K_1^+ \rightarrow K^*(892)^0 \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(3.1 \pm 1.7) \times 10^{-4}$	—
$K_1(1680)^+ K^-$, $K_1^+ \rightarrow K^{*0} \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(8.9 \pm 3.2) \times 10^{-5}$	—
$K^+ K^- \pi^+ \pi^-$ non-resonant	$(2.7 \pm 0.6) \times 10^{-4}$	—
$2K_S^0 \pi^+ \pi^-$	$(1.22 \pm 0.23) \times 10^{-3}$	673
$K_S^0 K^- 2\pi^+ \pi^-$	$< 1.4 \times 10^{-4}$	CL=90% 595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$	600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi\eta$	$(1.5 \pm 0.5) \times 10^{-4}$		489
$\phi\omega$	< 2.1	$\times 10^{-3}$	CL=90% 238

Radiative modes

$\rho^0\gamma$	$(1.82 \pm 0.32) \times 10^{-5}$		771
$\omega\gamma$	< 2.4	$\times 10^{-4}$	CL=90% 768
$\phi\gamma$	$(2.81 \pm 0.19) \times 10^{-5}$		654
$\bar{K}^*(892)^0\gamma$	$(4.2 \pm 0.7) \times 10^{-4}$		719

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

$K^+ \ell^- \bar{\nu}_\ell$ via \bar{D}^0	< 2.2	$\times 10^{-5}$	CL=90%	–
K^+ or $K^*(892)^+$ $e^- \bar{\nu}_e$ via \bar{D}^0	< 6	$\times 10^{-5}$	CL=90%	–
$K^+ \pi^-$ DC	$(1.50 \pm 0.07) \times 10^{-4}$		S=3.0	861
$K^+ \pi^-$ via DCS	$(1.364 \pm 0.026) \times 10^{-4}$			–
$K^+ \pi^-$ via \bar{D}^0	< 1.6	$\times 10^{-5}$	CL=95%	861
$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \bar{D}^0$	< 1.8	$\times 10^{-4}$	CL=95%	–
$K^*(892)^+ \pi^-$, $K^{*+} \rightarrow K_S^0 \pi^+$ DC	$(1.13 \pm_{-0.34}^{+0.60}) \times 10^{-4}$			711
$K_0^*(1430)^+ \pi^-$, $K_0^{*+} \rightarrow K_S^0 \pi^+$ DC	< 1.4	$\times 10^{-5}$		–
$K_2^*(1430)^+ \pi^-$, $K_2^{*+} \rightarrow K_S^0 \pi^+$ DC	< 3.4	$\times 10^{-5}$		–
$K^+ \pi^- \pi^0$ DC	$(3.06 \pm 0.15) \times 10^{-4}$			844
$K^+ \pi^- \pi^0$ via \bar{D}^0	$(7.6 \pm_{-0.6}^{+0.5}) \times 10^{-4}$			–
$K^+ \pi^+ 2\pi^-$ via DCS	$(2.49 \pm 0.07) \times 10^{-4}$			–
$K^+ \pi^+ 2\pi^-$ DC	$(2.65 \pm 0.06) \times 10^{-4}$			813
$K^+ \pi^+ 2\pi^-$ via \bar{D}^0	$(7.9 \pm 3.0) \times 10^{-6}$			812
$(K^*(892)^0 \rho(770)^0)$ S-wave via DCS, $K^*(892)^0 \rightarrow K^+ \pi^-$	$(2.4 \pm 0.5) \times 10^{-5}$			–
$(K^*(892)^0 \rho(770)^0)$ P-wave via DCS, $K^*(892)^0 \rightarrow K^+ \pi^-$	$(2.10 \pm 0.26) \times 10^{-5}$			–
$(K^*(892)^0 \rho(770)^0)$ D-wave via DCS, $K^*(892)^0 \rightarrow K^+ \pi^-$	$(2.54 \pm 0.33) \times 10^{-5}$			–

$(K^*(892)^0 \rho(1450)^0)$	$(2.0 \pm 0.5) \times 10^{-5}$	–
<i>S-wave</i> via DCS, $K^*(892)^0 \rightarrow$ $K^+ \pi^-$, $\rho(1450)^0 \rightarrow$ $\pi^+ \pi^-$		
$(K_1(1270)^+ \pi^-)$	$(4.5 \pm 0.6) \times 10^{-5}$	–
<i>S-wave</i> via DCS, $K_1(1270)^+ \rightarrow$ $K^+ \pi^- \pi^+$ total		
$(K_1(1400)^+ \pi^-)$	$(6.6 \pm 0.7) \times 10^{-5}$	–
<i>S-wave</i> via DCS, $K_1(1400)^+ \rightarrow$ $(K^*(892)^0 \pi^+)$		
<i>S-wave</i> , $K^*(892)^0 \rightarrow$ $K^+ \pi^-$		
$K^+ \pi^+ 2\pi^-$ non-resonant via DCS	$(5.2 \pm 0.5) \times 10^{-5}$	–
μ^- anything via \bar{D}^0	$< 4 \times 10^{-4}$	CL=90% –

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton Family number (LF) violating modes,
Lepton (L) or Baryon (B) number violating modes**

$\gamma\gamma$	C1	< 8.5	$\times 10^{-7}$	CL=90%	932
$e^+ e^-$	C1	< 7.9	$\times 10^{-8}$	CL=90%	932
$\mu^+ \mu^-$	C1	< 6.2	$\times 10^{-9}$	CL=90%	926
$\pi^0 e^+ e^-$	C1	< 4	$\times 10^{-6}$	CL=90%	928
$\pi^0 \mu^+ \mu^-$	C1	< 1.8	$\times 10^{-4}$	CL=90%	915
$\eta e^+ e^-$	C1	< 3	$\times 10^{-6}$	CL=90%	852
$\eta \mu^+ \mu^-$	C1	< 5.3	$\times 10^{-4}$	CL=90%	838
$\pi^+ \pi^- e^+ e^-$	C1	< 7	$\times 10^{-6}$	CL=90%	922
$\rho^0 e^+ e^-$	C1	< 1.0	$\times 10^{-4}$	CL=90%	771
$\pi^+ \pi^- \mu^+ \mu^-$	C1	(9.6 ± 1.2)	$\times 10^{-7}$		894
$\pi^+ \pi^- \mu^+ \mu^-$ (non-res)		< 5.5	$\times 10^{-7}$	CL=90%	–
$\rho^0 \mu^+ \mu^-$	C1	< 2.2	$\times 10^{-5}$	CL=90%	754
$\omega e^+ e^-$	C1	< 6	$\times 10^{-6}$	CL=90%	768
$\omega \mu^+ \mu^-$	C1	< 8.3	$\times 10^{-4}$	CL=90%	751
$K^- K^+ e^+ e^-$	C1	< 1.1	$\times 10^{-5}$	CL=90%	791
$\phi e^+ e^-$	C1	< 5.2	$\times 10^{-5}$	CL=90%	654
$K^- K^+ \mu^+ \mu^-$	C1	(1.54 ± 0.32)	$\times 10^{-7}$		710
$K^- K^+ \mu^+ \mu^-$ (non-res)		< 3.3	$\times 10^{-5}$	CL=90%	–
$\phi \mu^+ \mu^-$	C1	< 3.1	$\times 10^{-5}$	CL=90%	631
$\bar{K}^0 e^+ e^-$	[h]	< 2.4	$\times 10^{-5}$	CL=90%	866
$\bar{K}^0 \mu^+ \mu^-$	[h]	< 2.6	$\times 10^{-4}$	CL=90%	852

$K^- \pi^+ e^+ e^-$	CI	< 4.1	$\times 10^{-5}$	CL=90%	861
$\bar{K}^*(892)^0 e^+ e^-$		$[h] < 4.7$	$\times 10^{-5}$	CL=90%	719
$K^- \pi^+ \mu^+ \mu^-$	CI	< 3.59	$\times 10^{-4}$	CL=90%	829
$K^- \pi^+ \mu^+ \mu^-, 675 < m_{\mu\mu} < 875 \text{ MeV}$		(4.2 ± 0.4)	$\times 10^{-6}$		–
$\bar{K}^*(892)^0 \mu^+ \mu^-$		$[h] < 2.4$	$\times 10^{-5}$	CL=90%	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	CI	< 8.1	$\times 10^{-4}$	CL=90%	863
$\mu^\pm e^\mp$	LF	$[r] < 1.3$	$\times 10^{-8}$	CL=90%	929
$\pi^0 e^\pm \mu^\mp$	LF	$[r] < 8.6$	$\times 10^{-5}$	CL=90%	924
$\eta e^\pm \mu^\mp$	LF	$[r] < 1.0$	$\times 10^{-4}$	CL=90%	848
$\pi^+ \pi^- e^\pm \mu^\mp$	LF	$[r] < 1.5$	$\times 10^{-5}$	CL=90%	911
$\rho^0 e^\pm \mu^\mp$	LF	$[r] < 4.9$	$\times 10^{-5}$	CL=90%	767
$\omega e^\pm \mu^\mp$	LF	$[r] < 1.2$	$\times 10^{-4}$	CL=90%	764
$K^- K^+ e^\pm \mu^\mp$	LF	$[r] < 1.8$	$\times 10^{-4}$	CL=90%	754
$\phi e^\pm \mu^\mp$	LF	$[r] < 3.4$	$\times 10^{-5}$	CL=90%	648
$\bar{K}^0 e^\pm \mu^\mp$	LF	$[r] < 1.0$	$\times 10^{-4}$	CL=90%	863
$K^- \pi^+ e^\pm \mu^\mp$	LF	$[r] < 5.53$	$\times 10^{-4}$	CL=90%	848
$\bar{K}^*(892)^0 e^\pm \mu^\mp$	LF	$[r] < 8.3$	$\times 10^{-5}$	CL=90%	714
$2\pi^- 2e^+ + \text{c.c.}$	L	< 1.12	$\times 10^{-4}$	CL=90%	922
$2\pi^- 2\mu^+ + \text{c.c.}$	L	< 2.9	$\times 10^{-5}$	CL=90%	894
$K^- \pi^- 2e^+ + \text{c.c.}$	L	< 2.06	$\times 10^{-4}$	CL=90%	861
$K^- \pi^- 2\mu^+ + \text{c.c.}$	L	< 3.9	$\times 10^{-4}$	CL=90%	829
$2K^- 2e^+ + \text{c.c.}$	L	< 1.52	$\times 10^{-4}$	CL=90%	791
$2K^- 2\mu^+ + \text{c.c.}$	L	< 9.4	$\times 10^{-5}$	CL=90%	710
$\pi^- \pi^- e^+ \mu^+ + \text{c.c.}$	L	< 7.9	$\times 10^{-5}$	CL=90%	911
$K^- \pi^- e^+ \mu^+ + \text{c.c.}$	L	< 2.18	$\times 10^{-4}$	CL=90%	848
$2K^- e^+ \mu^+ + \text{c.c.}$	L	< 5.7	$\times 10^{-5}$	CL=90%	754
$p e^-$	L, B	$[s] < 1.0$	$\times 10^{-5}$	CL=90%	696
$\bar{p} e^+$	L, B	$[t] < 1.1$	$\times 10^{-5}$	CL=90%	696

$D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

$$\text{Mass } m = 2006.85 \pm 0.05 \text{ MeV} \quad (S = 1.1)$$

$$m_{D^{*0}} - m_{D^0} = 142.014 \pm 0.030 \text{ MeV} \quad (S = 1.5)$$

$$\text{Full width } \Gamma < 2.1 \text{ MeV, CL} = 90\%$$

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_j/Γ)	p (MeV/c)
$D^0 \pi^0$	$(64.7 \pm 0.9) \%$	43
$D^0 \gamma$	$(35.3 \pm 0.9) \%$	137

$D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

 I, J, P need confirmation.

 Mass $m = 2010.26 \pm 0.05$ MeV

$$m_{D^*(2010)^+} - m_{D^+} = 140.603 \pm 0.015$$
 MeV

$$m_{D^*(2010)^+} - m_{D^0} = 145.4257 \pm 0.0017$$
 MeV

 Full width $\Gamma = 83.4 \pm 1.8$ keV

 $D^*(2010)^-$ modes are charge conjugates of the modes below.

 $D^*(2010)^\pm$ DECAY MODES

 Fraction (Γ_i/Γ)

 p (MeV/c)

$D^0 \pi^+$	$(67.7 \pm 0.5) \%$	39
$D^+ \pi^0$	$(30.7 \pm 0.5) \%$	38
$D^+ \gamma$	$(1.6 \pm 0.4) \%$	136

 $D_0^*(2300)^0$

$$I(J^P) = \frac{1}{2}(0^+)$$

 Mass $m = 2300 \pm 19$ MeV

 Full width $\Gamma = 274 \pm 40$ MeV

 $D_0^*(2300)^0$ DECAY MODES

 Fraction (Γ_i/Γ)

 p (MeV/c)

$D^+ \pi^-$	seen	369
-------------	------	-----

 $D_1(2420)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

 Mass $m = 2420.8 \pm 0.5$ MeV ($S = 1.3$)

$$m_{D_1^0} - m_{D^{*+}} = 410.6 \pm 0.5$$
 ($S = 1.3$)

 Full width $\Gamma = 31.7 \pm 2.5$ MeV ($S = 3.5$)

 $\bar{D}_1(2420)^0$ modes are charge conjugates of modes below.

 $D_1(2420)^0$ DECAY MODES

 Fraction (Γ_i/Γ)

 p (MeV/c)

$D^*(2010)^+ \pi^-$	seen	353
$D^0 \pi^+ \pi^-$	seen	425
$D^+ \pi^-$	not seen	472
$D^{*0} \pi^+ \pi^-$	not seen	279

$D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $J^P = 2^+$ assignment strongly favored.

$$\text{Mass } m = 2460.7 \pm 0.4 \text{ MeV} \quad (S = 3.1)$$

$$m_{D_2^{*0}} - m_{D^+} = 591.0 \pm 0.4 \text{ MeV} \quad (S = 2.9)$$

$$m_{D_2^{*0}} - m_{D^{*+}} = 450.4 \pm 0.4 \text{ MeV} \quad (S = 2.9)$$

$$\text{Full width } \Gamma = 47.5 \pm 1.1 \text{ MeV} \quad (S = 1.8)$$

 $\bar{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

 $D_2^*(2460)^0$ DECAY MODES

 Fraction (Γ_i/Γ)

 p (MeV/c)

$D^+ \pi^-$	seen	505
$D^*(2010)^+ \pi^-$	seen	389
$D^0 \pi^+ \pi^-$	not seen	462
$D^{*0} \pi^+ \pi^-$	not seen	324

 $D_2^*(2460)^\pm$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $J^P = 2^+$ assignment strongly favored.

$$\text{Mass } m = 2465.4 \pm 1.3 \text{ MeV} \quad (S = 3.1)$$

$$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7 \text{ MeV}$$

$$\text{Full width } \Gamma = 46.7 \pm 1.2 \text{ MeV}$$

 $D_2^*(2460)^\pm$ modes are charge conjugates of modes below.

 $D_2^*(2460)^\pm$ DECAY MODES

 Fraction (Γ_i/Γ)

 p (MeV/c)

$D^0 \pi^+$	seen	513
$D^{*0} \pi^+$	seen	396
$D^+ \pi^+ \pi^-$	not seen	462
$D^{*+} \pi^+ \pi^-$	not seen	326

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^- 2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] Submodes of the $D^+ \rightarrow K^- 2\pi^+ \pi^0$ and $K_S^0 2\pi^+ \pi^-$ modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 *Review*, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $\bar{K}^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
- [l] This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.
- [n] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.17 ± 0.17 %.
- [o] This is a doubly Cabibbo-suppressed mode.
- [p] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [q] This branching fraction includes all the decay modes of the resonance in the final state.
- [r] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [s] This limit is for either D^0 or \bar{D}^0 to $p e^-$.
- [t] This limit is for either D^0 or \bar{D}^0 to $\bar{p} e^+$.