

24. Big Bang Nucleosynthesis

Revised October 2019 by B.D. Fields (Astronomy, Illinois U.; Physics, Illinois U.), P. Molaro (INAF-OATS Trieste) and S. Sarkar (Rudolf Peierls, Oxford U.).

24.1 Abstract

Big-Bang nucleosynthesis (BBN) offers the deepest reliable probe of the early Universe, being based on well-understood Standard Model physics [1]. Predictions of the abundances of the light elements, D, ^3He , ^4He , and ^7Li , synthesized at the end of the *first three minutes*, are in good overall agreement with the primordial abundances inferred from observational data, thus validating the standard hot Big-Bang cosmology (see [2–5] for reviews). This is particularly impressive given that these abundances span nine orders of magnitude – from $^4\text{He}/\text{H} \sim 0.08$ down to $^7\text{Li}/\text{H} \sim 10^{-10}$ (ratios by number). Thus BBN provides powerful constraints on possible deviations from the standard cosmology, and on new physics beyond the Standard Model [6–9].

24.2 Theory

The synthesis of the light elements is sensitive to physical conditions in the early radiation-dominated era at a temperature $T \sim 1$ MeV, corresponding to an age $t \sim 1$ s. At higher temperatures, weak interactions were in thermal equilibrium, thus fixing the ratio of the neutron and proton number densities to be $n/p = e^{-Q/T}$, where $Q = 1.293$ MeV is the neutron-proton mass difference. As the temperature dropped, the neutron-proton inter-conversion rate per nucleon, $\Gamma_{n \leftrightarrow p} \sim G_{\text{F}}^2 T^5$, fell faster than the Hubble expansion rate, $H \sim \sqrt{g_* G_{\text{N}}} T^2$, where g_* counts the number of relativistic particle species determining the energy density in radiation (see The Cosmological Parameters—Sec. 22 of this *Review*). This resulted in departure from chemical equilibrium (*freeze-out*) at $T_{\text{fr}} \sim (g_* G_{\text{N}}/G_{\text{F}}^4)^{1/6} \simeq 1$ MeV. The neutron fraction at this time, $n/p = e^{-Q/T_{\text{fr}}} \simeq 1/6$, is thus sensitive to every known physical interaction, since Q is determined by both strong and electromagnetic interactions while T_{fr} depends on the weak as well as gravitational interactions. Moreover, the sensitivity to the Hubble expansion rate affords a probe of, *e.g.*, the number of relativistic neutrino species [10]. After freeze-out, the neutrons were free to β -decay, so the neutron fraction dropped to $n/p \simeq 1/7$ by the time nuclear reactions began. A simplified analytic model of freeze-out yields the n/p ratio to an accuracy of $\sim 1\%$ [11, 12].

The rates of these reactions depend on the density of baryons (strictly speaking, nucleons), which is usually expressed normalized to the relic blackbody photon density as $\eta \equiv n_b/n_\gamma$. As we shall see, all the light-element abundances can be explained with $\eta_{10} \equiv \eta \times 10^{10}$ in the range 5.8–6.5 (95% CL). With n_γ fixed by the present CMB temperature 2.7255 K (see The Cosmological Parameters—Sec. 29 of this *Review*), this can be stated as the allowed range for the baryon mass density today, $\rho_b = (3.9\text{--}4.6) \times 10^{-31}$ g cm $^{-3}$, or as the baryonic fraction of the critical density, $\Omega_b = \rho_b/\rho_{\text{crit}} \simeq \eta_{10} h^{-2}/274 = (0.021\text{--}0.024) h^{-2}$, where $h \equiv H_0/100$ km s $^{-1}$ Mpc $^{-1}$ is the present Hubble parameter (see The Cosmological Parameters—Sec. 25.1 of this *Review*).

The nucleosynthesis chain begins with the formation of deuterium in the process $p(n, \gamma)\text{D}$. However, photo-dissociation by the high number density of photons delays production of deuterium (and other complex nuclei) until well after T drops below the binding energy of deuterium, $\Delta_{\text{D}} = 2.23$ MeV. The quantity $\eta^{-1} e^{-\Delta_{\text{D}}/T}$, *i.e.*, the number of photons per baryon above the deuterium photo-dissociation threshold, falls below unity at $T \simeq 0.1$ MeV; nuclei can then begin to form without being immediately photo-dissociated again. Only 2-body reactions, such as $\text{D}(p, \gamma)^3\text{He}$ and $^3\text{He}(\text{D}, p)^4\text{He}$ are important because the density by this time has become rather low – comparable to that of air!

Nearly all neutrons end up bound in the most stable light element ^4He . Heavier nuclei do not

form in any significant quantity both because of the absence of stable nuclei with mass number 5 or 8 (which impedes nucleosynthesis via $n^4\text{He}$, $p^4\text{He}$ or $^4\text{He}^4\text{He}$ reactions), and the large Coulomb barriers for reactions such as $^3\text{He}(^4\text{He}, \gamma)^7\text{Li}$ and $^3\text{He}(^4\text{He}, \gamma)^7\text{Be}$. Hence the primordial mass fraction of ^4He , $Y_p \equiv \rho(^4\text{He})/\rho_b$, can be estimated by the simple counting argument

$$Y_p = \frac{2(n/p)}{1 + n/p} \simeq 0.25 \quad (24.1)$$

where strictly speaking this gives the baryon fraction in ^4He , which is what we will quote throughout. This differs slightly from the mass fraction due to small binding energy corrections.

There is little sensitivity here to the actual nuclear reaction rates, which are, however, important in determining the other ‘left-over’ abundances: D and ^3He at the level of a few times 10^{-5} by number relative to H, and $^7\text{Li}/\text{H}$ at the level of about 10^{-10} (when η_{10} is in the range 1–10). These values can be understood in terms of approximate analytic arguments [12, 13]. The experimental parameter most important in determining Y_p is the neutron lifetime, τ_n , which normalizes (the inverse of) $\Gamma_{n\rightarrow p}$. Its value has recently been significantly revised downwards to $\tau_n = 879.4 \pm 0.6$ s (see *N Baryons Listing*).

The elemental abundances shown in Fig. 24.1 as a function of η_{10} were calculated [14] using an updated version [15] of the Wagoner code [1]; other versions [16–19] too are publicly available. The ^4He curve includes small corrections due to radiative processes at zero and finite temperatures [20], non-equilibrium neutrino heating during e^\pm annihilation [21], and finite nucleon mass effects [22]; the range primarily reflects the 2σ uncertainty in the neutron lifetime. The spread in the curves for D, ^3He , and ^7Li corresponds to the 2σ uncertainties in nuclear cross sections, as estimated by Monte Carlo methods [15, 23–25]. The input nuclear data have been carefully reassessed [2, 14, 15, 23–25, 25–29, 29], leading to improved precision for the abundance predictions. In particular, the uncertainty in $^7\text{Li}/\text{H}$ at interesting values of η has been reduced recently by a factor ~ 2 , a consequence of a similar reduction in the error budget [30] for the dominant mass-7 production channel $^3\text{He}(^4\text{He}, \gamma)^7\text{Be}$. Polynomial fits to the predicted abundances and the error correlation matrix have been given in refs. [24, 31]. The boxes in Fig 24.1 show the observationally inferred primordial abundances with their associated uncertainties, as discussed below.

24.3 Light Element Abundances

BBN theory predicts the universal abundances of D, ^3He , ^4He , and ^7Li which are essentially fixed by $t \sim 180$ s. However, abundances are observed at much later epochs, after stellar nucleosynthesis commenced. Stars produce heavy elements such as C, N, O, and Fe (“metals”), while the ejected remains of stellar processing alters the light element abundances from their primordial values. Thus, one seeks astrophysical sites with low metal abundances to measure light element abundances that are closer to primordial.

BBN is the only significant source of deuterium which is entirely destroyed when it is cycled into stars [32]. Thus, any detection provides a lower limit to primordial D/H, and an upper limit on η_{10} . The best proxy to the primordial value of D is its measure in distant and chemically unprocessed matter, where stellar processing (astration) is minimal [32]. This has become possible with the advent of large telescopes, but after two decades of observational efforts we have only about a dozen determinations listed in Table 24.1 [33–46].

High-resolution spectra reveal the presence of D in high-redshift, low-metallicity quasar absorption systems via its isotope-shifted Lyman- α absorption features, though, unfortunately, these are often obscured or contaminated by the hydrogen features of the Lyman- α forest.

The nuclear reaction cross sections important for BBN have all been measured at the relevant energies. Recently however there have been substantial advances in the precision of light element




Figure 24.1: The primordial abundances of ${}^4\text{He}$, D, ${}^3\text{He}$, and ${}^7\text{Li}$ as predicted by the standard model of Big-Bang nucleosynthesis — the bands show the 95% CL range [47]. Boxes indicate the observed light element abundances. The narrow vertical band indicates the CMB measure of the cosmic baryon density, while the wider band indicates the BBN D+ ${}^4\text{He}$ concordance range (both at 95% CL).

observations (e.g., D/H) and in the determination of cosmological parameters (e.g., from *Planck*). This motivates corresponding improvement in BBN predictions and thus in the key reaction cross sections. For example, it has been suggested [48, 49] that $d(p, \gamma){}^3\text{He}$ measurements may suffer from

systematic errors and be inferior to *ab initio* theory; if so, this could alter D/H abundances at a level that is now significant. Ongoing low-background cross section measurements should resolve this issue [50].

A few DLA systems show D lines resolved up to the higher members of the Lyman series. Recent determinations [37, 39] and re-analyses [38, 46, 51] provide strikingly improved precision over earlier work. A weighted mean of the 11 most precise measurements in Table 24.1 provides a value with a one percent precision:

$$D/H|_p \times 10^6 = (25.47 \pm 0.25). \quad (24.2)$$

Considering all the 16 extant determinations, the weighted mean is $D/H|_p \times 10^6 = (25.36 \pm 0.26)$, while different selections provide $D/H|_p \times 10^6 = (25.27 \pm 0.30)$ [51] or $D/H|_p \times 10^6 = (25.45 \pm 0.25)$ [52], all consistent with each other within 1σ . Metallicities of the absorbers are $(0.001 - 0.03) \times$ Solar, i.e. at a level where no significant astration is expected [34, 53]. D/H shows no correlation with metallicity, redshift, or the hydrogen column density $N(H)$ ($= \int_{\text{los}} n_H ds$) integrated over the line-of-sight through the absorber. In the Galaxy D/H measurements are anti-correlated with metal abundances, which suggests that interstellar D partly resides in dust particles [54]. However, in the absorbers where deuterium is measured, the dust content is quite small as implied by Solar proportions of the abundances of refractory and non refractory elements. This is consistent with the measured D/H being truly representative of the primordial value.

Table 24.1: D/H measurements. For systems with multiple measurements we used the most recent one which is generally more precise.

QSO	z_{em}	z_{em}	$\log(NHI)$	[X/H]	(D/H) $\times 10^6$	Ref
QSO 2206–199	2.56	2.076	20.436 \pm 0.008	-2.04	16.5 \pm 3.5	[55]
QSO 0347–3819	3.22	3.025	20.63 \pm 0.09	-1.25	22.4 \pm 6.7	[56]
SDSS J1134+5742	3.52	3.411	17.95 \pm 0.05	<-4.2	20.4 \pm 6.1	[41]
QSO CTQ 247	3.02	2.621	20.45 \pm 0.10	-1.99	28.0 \pm 8.0	[43]
SDSS 1337+3152	3.17	3.168	20.41 \pm 0.15	-2.68	12.0 \pm 5.0	[57]
SDSS J1419+0829	3.03	3.049	20.392 \pm 0.003	-1.92	25.1 \pm 0.5	[36]
HS 0105+1619	2.65	2.536	19.40 \pm 0.01	-1.77	25.8 \pm 1.5	[36]
QSO B0913+0715	2.78	2.618	20.312 \pm 0.008	-2.40	25.3 \pm 1.0	[36]
SDSS J1358+0349	2.89	2.853	20.524 \pm 0.006	-2.33	26.2 \pm 0.7	[37]
SDSS J1358+6522	3.17	3.067	20.50 \pm 0.01	-2.33	25.8 \pm 1.0	[36]
SDSS J1558–0031	2.82	2.702	20.75 \pm 0.03	-1.55	24.0 \pm 1.4	[36]
PKS 1937–1009	3.78	3.256	18.09 \pm 0.03	-1.87	24.5 \pm 2.8	[58]
QSO J1444+2919	2.66	2.437	19.983 \pm 0.010	-2.04	19.7 \pm 3.3	[39]
PKS 1937–1009	3.78	3.572	17.925 \pm 0.006	-2.26	26.2 \pm 0.5	[46]
QSO 1009+2956	2.63	2.504	17.362 \pm 0.005	-2.50	24.8 \pm 4.1	[52]
QSO 1243+307	2.55	2.525	19.761 \pm 0.026	-2.77	23.9 \pm 1.0	[51]
Weighted mean of all 16					25.38 \pm 0.25	
Weighted mean of the most recent 11					25.47 \pm 0.25	

The primordial ${}^4\text{He}$ abundance is best determined through recombination emission lines of He and H in the most metal-poor extragalactic HII (ionized) regions, *viz.* blue compact galaxies, generally found at low redshift. There is now a large body of data on ${}^4\text{He}$ and CNO in these galaxies,

with over 1000 such systems in the Sloan Digital Sky Survey alone [59]. These data confirm that the stellar contribution to the helium abundance is positively correlated with metal production, so extrapolation to zero metallicity gives the primordial ${}^4\text{He}$ abundance Y_p . However, HII regions are complex systems and several physical parameters enter in the He/H determination, notably the electron density and temperature, as well as reddening. Thus, systematic effects dominate the uncertainties in the abundance determination [60,61]. A major step forward has been the inclusion of the He $\lambda 10830$ infrared emission line which shows a strong dependence on the electron density and is thus useful to break the degeneracy with the temperature, allowing for a more robust helium abundance determination. In recent works the underlying ${}^4\text{He}$ stellar absorption, and/or the newly derived values of the HeI-recombination and H-excitation-collisional coefficients are addressed and the ${}^4\text{He}$ abundances have increased significantly. Some recent results are $Y_p = 0.2451 \pm 0.0026$ [62], $Y_p = 0.2449 \pm 0.0040$ [63] and $Y_p = 0.2551 \pm 0.0022$ [59] — see Ref. [64] and references therein for previous determinations. Taking all this into account our recommended ${}^4\text{He}$ abundance is

$$Y_p = 0.245 \pm 0.003, \quad (24.3)$$

but the matter is far from settled given that the measurements are only marginally consistent.

The best suited objects for the primordial ${}^7\text{Li}$ determination are metal-poor stars in the Galactic halo, which have metallicities going down to 10^{-6} of the Solar value [65]. Observations have long shown [66–69] that ${}^7\text{Li}$ does not vary significantly in halo dwarfs with metallicities $\lesssim 1/30$ of Solar — the *Spite plateau* [66, 70]. Recent observations show a puzzling drop in the Li/H abundance in metal-poor stars with $[\text{Fe}/\text{H}] < -3.0$ [71–73]. This becomes particularly acute at the very low metallicity end where only one star out of the seven dwarfs with metallicities $[\text{Fe}/\text{H}] \lesssim -4.5$ shows a ${}^7\text{Li}$ abundance close to the Spite Plateau, while in the others where it ought to be present it is either lower or totally absent [65, 74]. The reason for the increase in scatter at low metallicity is unknown and prevents derivation of the primordial ${}^7\text{Li}$ value by extrapolating to zero metallicity [72, 73].

The primordial ${}^7\text{Li}$ in different samples of stars or globular clusters has been reported as $\text{Li}/\text{H}|_p = (1.7 \pm 0.3) \times 10^{-10}$ [69], $\text{Li}/\text{H}|_p = (2.19 \pm 0.28) \times 10^{-10}$ [75], and $\text{Li}/\text{H}|_p = (1.86 \pm 0.23) \times 10^{-10}$ [76]. We note that different parameters (*e.g.*, the temperature) of the stellar atmosphere in which the ${}^7\text{Li}$ absorption line is formed lead to slightly different values. To estimate the primordial ${}^7\text{Li}$ value we consider only stars with metallicity in the range $-2.8 < [\text{Fe}/\text{H}] < -1.5$ [73], where no scatter in excess of the observational errors is observed. This yields:

$$\text{Li}/\text{H}|_p = (1.6 \pm 0.3) \times 10^{-10}. \quad (24.4)$$

Strictly speaking the suggested primordial ${}^7\text{Li}$ abundance should be considered a *lower bound* rather than a measure. In fact, ${}^7\text{Li}$ in Pop II stars may have been partially destroyed due to mixing of the outer layers with the hotter interior [77]. Such processes can be constrained by the absence of significant scatter in ${}^7\text{Li}$ versus T_{eff} [68], but ${}^7\text{Li}$ depletion by a factor as large as ~ 1.8 may have occurred [68, 78]. A recent model predicts that an initial ${}^7\text{Li}$ abundance of $\text{Li}/\text{H}|_p = 5.3 \times 10^{-10}$, corresponding to the baryon density indicated by CMB and D/H abundance, is significantly destroyed in the pre-main-sequence phase and then partially restored by late accretion of fresh ${}^7\text{Li}$ normal material ending onto the Spite Plateau [79, 80].

A ${}^6\text{Li}$ plateau (analogous to the ${}^7\text{Li}$ plateau) has also been claimed [81]. This has however been challenged by new observations and analyses which show that stellar convective motions can generate asymmetries in the line shape that mimic the presence of ${}^6\text{Li}$ [82–84]. Recent high-precision measurements are sensitive to the tiny isotopic shift in absorption indicate ${}^6\text{Li}/{}^7\text{Li} \leq 0.05$, thus confirming that ${}^7\text{Li}$ is dominant [81, 82].

The primordial abundance of ${}^3\text{He}$ has the poorest observational determination of all of the light nuclides. The only data available come from the Solar system and from solar-metallicity HII regions in the Galaxy [85]. Therefore, inferring the primordial ${}^3\text{He}$ abundance is problematic, compounded by the fact that stellar nucleosynthesis models for ${}^3\text{He}$ are in conflict with observations. Consequently, we consider it inappropriate to use ${}^3\text{He}$ (and also $\text{D}+{}^3\text{He}$) as a cosmological probe.

24.4 Concordance, Dark Matter, and the CMB

We now use the observed light element abundances to test the theory. We first consider standard BBN, which is based on Standard Model physics alone, so $N_\nu = 3$ and the only free parameter is the baryon-to-photon ratio η . (The implications of BBN for physics beyond the Standard Model will be considered below). Thus, any abundance measurement determines η , and additional measurements overconstrain the theory and thereby provide a consistency check.

While the η ranges spanned by the boxes in Fig 24.1 do not all overlap, they are all within a factor ~ 2 of each other. In particular, the lithium abundance corresponds to η values that are inconsistent with that of the (now very precise) D/H abundance as well as the less-constraining ${}^4\text{He}$ abundance. This discrepancy marks the *lithium problem*. The problem could simply reflect difficulty in determining the primordial lithium abundance, or could hint at a more fundamental omission in the theory. The possibility that lithium reveals new physics is addressed in detail in the next section. If however we exclude the lithium constraint because its inferred abundance may suffer from systematic uncertainties, then D/H and ${}^4\text{He}$ are in agreement. The concordant η range is essentially that implied by D/H , namely

$$5.8 \leq \eta_{10} \leq 6.5 \text{ (95\% CL)}. \quad (24.5)$$

Despite the lithium problem, the overall concordance remains remarkable: using only well-established microphysics we can extrapolate back to $t \sim 1$ s to predict light element abundances spanning nine orders of magnitude, in approximate agreement with observation. This is a major success for the standard cosmology, and inspires confidence in extrapolation back to such early times.

This concordance provides a measure of the baryon content:

$$0.021 \leq \Omega_b h^2 \leq 0.024 \text{ (95\% CL)}, \quad (24.6)$$

a result that plays a key role in our understanding of the matter budget of the Universe. First of all $\Omega_b \ll 1$, *i.e.*, baryons cannot close the Universe [86]. Furthermore, the cosmic density of (optically) luminous matter is $\Omega_{\text{lum}} \simeq 0.0024 h^{-1}$ [87], so that $\Omega_b \gg \Omega_{\text{lum}}$: most baryons are optically dark, probably in the form of a diffuse intergalactic medium [88]. Finally, given that $\Omega_m \sim 0.3$ (see Dark Matter and Cosmological Parameters reviews), we infer that most matter in the Universe is not only dark, but also takes some non-baryonic (more precisely, non-nucleonic) form.

The BBN prediction for the cosmic baryon density can be tested through precision measurements of CMB temperature fluctuations (see Cosmic Microwave Background review). One can determine η from the amplitudes of the acoustic peaks in the CMB angular power spectrum [89], making it possible to compare two measures of η using very different physics, at two widely separated epochs. In the standard cosmology, there is no change in η between BBN and CMB decoupling, thus, a comparison of η_{BBN} and η_{CMB} is a key test. Agreement would endorse the standard picture, while disagreement could point to new physics during/between the BBN and CMB epochs.

The analysis described in the Cosmic Microwave Background review, based on *Planck* TT, TE, EE + lowE data and lensing, yields $\Omega_b h^2 = 0.02237 \pm 0.00015$ [90], which corresponds to $\eta_{10} = 6.12 \pm 0.04$ [91]. This result depends weakly on the primordial helium abundance, and the fiducial *Planck* analysis uses BBN theory to fix $Y_p(\eta)$. Without BBN theory, the *Planck* TT, TE, EE + lowE

data plus lensing give $\Omega_b h^2 = 0.02230 \pm 0.00020$, corresponding to $\eta_{10} = 6.105 \pm 0.055$. As shown in Fig. 24.1, this CMB estimate of the baryon density (narrow vertical band) is consistent with the BBN range, *i.e.*, in good agreement with the value inferred from high-redshift D/H measurements and local ${}^4\text{He}$ determinations; together these observations span diverse environments from redshifts $z \sim 1000$ to the present.

The ${}^4\text{He}$ abundance is proportional to the n/p ratio when the weak-interaction rate falls behind the Hubble expansion rate at $T_{\text{fr}} \sim 1$ MeV. The presence of additional neutrino flavors (or of any other relativistic species) at this time increases g_* , hence the expansion rate, leading to a larger value of T_{fr} , n/p , and therefore Y_p [10, 92]. In the Standard Model at $T = 1$ MeV, $g_* = 5.5 + \frac{7}{4}N_\nu$, where N_ν is the *effective* number of (nearly) massless neutrino flavors. The helium curves in 24.1 were computed taking $N_\nu = 3$; small corrections for non-equilibrium neutrino heating [21] are included in the thermal evolution and lead to an effective $N_\nu = 3.045$ compared to assuming instantaneous neutrino freezeout (see The Cosmological Parameters—Sec. 22 of this *Review*). The computed ${}^4\text{He}$ abundance scales as $\Delta Y_p \simeq 0.013 \Delta N_\nu$ [11]. Clearly the central value for N_ν from BBN will depend on η , which is independently determined (with weaker sensitivity to N_ν) by the adopted D or ${}^7\text{Li}$ abundance. For example, if the best value for the observed primordial ${}^4\text{He}$ abundance is 0.249, then, for $\eta_{10} \sim 6$, the central value for N_ν is very close to 3. A maximum likelihood analysis on η and N_ν based on ${}^4\text{He}$ and D abundances nearly identical to those above finds the (correlated) 95% CL ranges to be $5.6 < \eta_{10} < 6.6$ and $2.3 < N_\nu < 3.4$ [5]. Identical results are obtained using a simpler method to extract such bounds based on χ^2 statistics, given a set of input abundances [93].

The CMB damping tail is sensitive to the primordial ${}^4\text{He}$ abundance independently of both BBN and local ${}^4\text{He}$ measurements [94]. The *Planck* analysis using TT, TE, EE+lowE and lensing but not the BBN $Y_p(\eta)$ relation gives a ${}^4\text{He}$ mass fraction $0.239_{-0.025}^{+0.024}$, and nucleon fraction $Y_p = 0.240_{-0.25}^{+0.24}$, both at 95% CL [90]. This is consistent with the HII region helium abundance determination. Moreover, this value is consistent with the Standard ($N_\nu = 3$) BBN prediction for Y_p with the *Planck*-determined baryon density.

This concordance represents a successful CMB-only test of BBN.

The precision determination of the baryon density using the CMB motivates using this as an input to BBN calculations. Within the context of the Standard Model, BBN then becomes a zero-parameter theory, and the light element abundances are completely determined to within the uncertainties in η_{CMB} and the BBN theoretical errors. Comparison with the observed abundances then can be used to test the astrophysics of post-BBN light element evolution [95]. Alternatively, one can consider possible physics beyond the Standard Model (*e.g.*, which might change the expansion rate during BBN) and then use all of the abundances to test such models; this is discussed in 23.6 below.

24.5 The Lithium Problem

As Fig. 24.1 shows, stellar Li/H measurements are inconsistent with the D/H (and CMB), given the error budgets we have quoted. Recent updates in nuclear cross sections and stellar abundance systematics *increase* the discrepancy to over 5σ , depending on the stellar abundance analysis adopted [14]. For instance it is found $\text{Li}/\text{H}|_p = (5.623 \pm 0.247) \times 10^{-10}$ in [96], *i.e.* a factor 3.5 higher than equation 24.4 and at a 10σ CL.

The question then becomes pressing as to whether this mismatch comes from systematic errors in the observed abundances, and/or uncertainties in stellar astrophysics or nuclear inputs, or whether there might be new physics at work [9]. Nuclear inputs (cross sections) for BBN reactions are constrained by extensive laboratory measurements; to increase ${}^7\text{Be}$ destruction requires enhancement of otherwise subdominant processes that can be attained by missed resonances in a few reactions such as ${}^7\text{Be}(d, p)2\alpha$ if the compound nuclear state properties are particularly favor-

able [29, 97–99]. However, experimental searches have now closed off these possibilities [100–102], making a *nuclear fix* increasingly unlikely.

Another conventional means to solve the lithium problem is by *in situ* destruction over the long lifetimes of the host halo stars. Stellar depletion mechanisms include diffusion, rotationally induced mixing, or pre-main-sequence depletion. These effects certainly occur, but to reduce lithium to the required levels generally requires some *ad hoc* mechanism and fine tuning of the initial stellar parameters [79, 80, 103]. A putative signature of diffusion has been reported for the globular clusters NGC 6397 and NGC 6752, where the “turnoff” stars exhibit slightly lower (by a factor ~ 1.3) abundances of Fe II TiII, ScII, CaI and MgI, than in more evolved stars [78, 104]. General features of diffusive models are a dispersion in the Li abundances and a pronounced downturn in the Li abundances at the hot end of the Li plateau. Some extra turbulence needs to be invoked to limit diffusion in the hotter stars and to restore uniform Li abundance along the Spite plateau [103]. Li destruction in the pre-Main sequence phase and partially restored by the accretion has been also proposed [80]

As nuclear and astrophysical solutions to the lithium problem become increasingly constrained (even if difficult to rule out definitively), the possibility of new physics arises. Nucleosynthesis models in which the baryon-to-photon ratio is inhomogeneous can alter abundances for a given η_{BBN} , but will overproduce ${}^7\text{Li}$ [105]. Entropy generation by some non-standard process could have decreased η between the BBN era and CMB decoupling, however the lack of spectral distortions in the CMB rules out any significant energy injection up to a redshift $z \sim 10^7$ [106]. The most intriguing resolution of the lithium problem thus involves new physics during BBN [7–9]

We summarize the general features of such solutions here, and later consider examples in the context of specific particle physics models. Many proposed solutions introduce perturbations to light-element formation during BBN; while all element abundances may suffer perturbations, the interplay of ${}^7\text{Li}$ and D is often the most important *i.e.* observations of D often provide the strongest constraints on the allowed perturbations to ${}^7\text{Li}$. In this connection it is important to note that the new, very precise determination of D/H will significantly constrain the ability of such models to ameliorate or solve the lithium problem.

A well studied class of models invokes the injection of suprathreshold hadronic or electromagnetic particles due to decays of dark matter particles. The effects are complex and depend on the nature of the decaying particles and their branchings and spectra. However, the models that most successfully solve the lithium problem generally feature non-thermal nucleons, which dissociate all light elements. Dissociation of even a small fraction of ${}^4\text{He}$ introduces a large abundance of free neutrons, which quickly thermalize. The thermal neutrons drive the ${}^7\text{Be}(n, p){}^7\text{Li}$ conversion of ${}^7\text{Be}$. The resulting ${}^7\text{Li}$ has a lower Coulomb barrier relative to ${}^7\text{Be}$ and is readily destroyed via ${}^7\text{Li}(p, \alpha){}^4\text{He}$ [107, 108]. But ${}^4\text{He}$ dissociation also produces D directly as well as via nonthermal neutron $n(p, \gamma)d$ reactions. This introduces a tension between Li/H reduction and D/H enhancement that becomes increasingly restrictive with the increasing precision of deuterium observations. Indeed, this now forces particle injection scenarios to make very small ${}^7\text{Li}$ perturbations — far short of the level needed. An exception is a recent model wherein MeV-scale decays by construction avoid ${}^4\text{He}$ dissociation and associated D/H overproduction, instead *borrowing* neutrons by dissociating only deuterons [109].

Another important class of models retains the standard cosmic particle content, but changes their interactions via time variations in the fundamental constants [110–115]. Here too, the details are model-dependent, but scenarios that solve or alleviate the lithium problem often feature perturbations to the deuteron binding energy. A weaker D binding leads to the D bottleneck being overcome later, so that element formation commences at a lower temperature and lower density. This leads in turn to slower nuclear rates that freeze out earlier. The net result is a *higher* final

D/H, due to less efficient processing into ${}^4\text{He}$, but also *lower* Li, due to suppressed production via ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$.

The *cosmological lithium problem* remains an unresolved issue in BBN. Nevertheless, the remarkable concordance between the CMB and the D (as well as ${}^4\text{He}$) abundance, is a non-trivial success, and provides important constraints on the early Universe.

24.6 Beyond the Standard Model

Given the simple physics underlying BBN, it is remarkable that it still provides the most effective test for the cosmological viability of ideas concerning physics beyond the Standard Model. Although baryogenesis and inflation must have occurred at higher temperatures in the early Universe, we do not as yet have ‘standard models’ for these, so BBN still marks the boundary between the established and the speculative in Big Bang cosmology. It might appear possible to push the boundary back to the quark-hadron transition at $T \sim \Lambda_{\text{QCD}}$, or electroweak symmetry breaking at $T \sim 1/\sqrt{G_{\text{F}}}$; however, so far no observable relics of these epochs have been identified, either theoretically or observationally. Thus, although the Standard Model provides a precise description of physics up to the Fermi scale, cosmology cannot be traced in detail before the BBN era.

The CMB power spectrum in the damping tail is independently sensitive to N_ν (*e.g.* [116]). The CMB value N_ν^{CMB} probes the cosmic radiation content at (re)combination, so a discrepancy would imply new physics or astrophysics. Indeed, observations by the South Pole Telescope implied $N_\nu^{\text{CMB}} = 3.85 \pm 0.62$ [117], prompting discussion of *dark radiation* such as sterile neutrinos [118]. However, *Planck* 2018 results give $N_\nu^{\text{CMB}} = 2.92_{+0.37}^{+0.36}$, 95% CL, when using Planck TT, TE, EE+lowE, a result quite consistent with the 3.045 of the Standard Model neutrinos [90].

Just as one can use the measured helium abundance to place limits on g_* [92, 111, 119–121], any changes in the strong, weak, electromagnetic, or gravitational coupling constants, arising *e.g.*, from the dynamics of new dimensions, can be similarly constrained [122], as can any speed-up of the expansion rate in, *e.g.*, scalar-tensor theories of gravity [123].

The limits on N_ν can be translated into limits on other types of particles or particle masses that would affect the expansion rate of the Universe during nucleosynthesis. For example, consider *sterile* neutrinos with only right-handed interactions of strength $G_{\text{R}} < G_{\text{F}}$. Such particles would decouple at higher temperature than (left-handed) neutrinos, so their number density ($\propto T^3$) relative to neutrinos would be reduced by any subsequent entropy release, *e.g.*, due to annihilations of massive particles that become non-relativistic between the two decoupling temperatures. Thus (relativistic) particles with less than full strength weak interactions contribute less to the energy density than particles that remain in equilibrium up to the time of nucleosynthesis [124]. If we impose $N_\nu < 4$ as an illustrative constraint, then the three right-handed neutrinos must have a temperature $3(T_{\nu_{\text{R}}}/T_{\nu_{\text{L}}})^4 < 1$. Since the temperature of the decoupled ν_{R} is determined by entropy conservation (see The Cosmological Parameters—Sec. 22 of this *Review*), $T_{\nu_{\text{R}}}/T_{\nu_{\text{L}}} = [(43/4)/g_*(T_{\text{d}})]^{1/3} < 0.76$, where T_{d} is the decoupling temperature of the ν_{R} . This requires $g_*(T_{\text{d}}) > 24$, so decoupling must have occurred at $T_{\text{d}} > 140$ MeV. The decoupling temperature is related to G_{R} through $(G_{\text{R}}/G_{\text{F}})^2 \sim (T_{\text{d}}/3 \text{ MeV})^{-3}$, where 3 MeV is the decoupling temperature for ν_{L} s. This yields a limit $G_{\text{R}} \lesssim 10^{-2}G_{\text{F}}$. The above argument sets lower limits on the masses of new Z' gauge bosons to which right-handed neutrinos would be coupled in models of superstrings [125], or extended technicolour [126]. Similarly a Dirac magnetic moment for neutrinos, which would allow the right-handed states to be produced through scattering and thus increase g_* , can be significantly constrained [127], as can any new interactions for neutrinos that have a similar effect [128–130]. Right-handed states can be populated directly by helicity-flip scattering if the neutrino mass is large enough, and this property has been used to infer a bound of $m_{\nu_\tau} \lesssim 1$ MeV (taking $N_\nu < 4$) [131]. If there is mixing between active and sterile neutrinos then the effect on BBN is more complicated [132, 133].

BBN limits on the cosmic expansion rate constrain supersymmetric scenarios in which the neutralino or gravitino are very light, so that they contribute to g_* [134]. A gravitino in the mass range $\sim 10^{-4} - 10$ eV will affect the expansion rate of the Universe similarly to a light neutralino (which is however now probably ruled out by collider data, especially the decays of the Higgs-like boson). The net contribution to N_ν then ranges between 0.74 and 1.69, depending on the gravitino and slepton masses [135].

The limit on the expansion rate during BBN can also be translated into bounds on the mass/lifetime of non-relativistic particles that decay during BBN. This results in an even faster speed-up rate, and typically also changes the entropy [136–138]. If the decays include Standard Model particles, the resulting electromagnetic [139] [95, 125, 140] and/or hadronic [141, 142] cascades can strongly perturb the light elements, which leads to even stronger constraints. Such arguments have been applied to rule out an MeV mass for ν_τ , which decays during nucleosynthesis [143].

Decaying-particle arguments have proved very effective in probing supersymmetry. Light-element abundances generally are complementary to accelerator data in constraining SUSY parameter space, with BBN reaching to values kinematically inaccessible to the LHC. Much recent interest has focused on the case in which the next-to-lightest supersymmetric particle is metastable and decays during or after BBN. The constraints on unstable particles discussed above imply stringent bounds on the allowed abundance of such particles [107]; if the metastable particle is charged (*e.g.*, the stau), then it is possible for it to form atom-like electromagnetic bound states with nuclei, and the resulting impact on light elements can be quite complex [8, 97, 144]. Moreover, SUSY decays can destroy ${}^7\text{Li}$ and/or produce ${}^6\text{Li}$, leading to a possible supersymmetric solution to the lithium problems noted above [145] (see [7] for a review).

These arguments impose powerful constraints on supersymmetric inflationary cosmology [95, 125, 140–142], particularly thermal leptogenesis [146]. These limits can be evaded only if the gravitino is massive enough to decay before BBN, *i.e.*, $m_{3/2} \gtrsim 50$ TeV [147] (which would be unnatural), or if it is in fact the lightest supersymmetric particle and thus stable [125, 140, 148, 149]. Similar constraints apply to moduli – very weakly coupled fields in string theory that obtain an electroweak-scale mass from supersymmetry breaking [150].

Finally, we mention that BBN places powerful constraints on the possibility that there are new large dimensions in nature, perhaps enabling the scale of quantum gravity to be as low as the electroweak scale [151]. Thus, Standard Model fields may be localized on a *brane*, while gravity alone propagates in the *bulk*. It has been further noted that the new dimensions may be non-compact, even infinite [152], and the cosmology of such models has attracted considerable attention. The expansion rate in the early Universe can be significantly modified, so BBN is able to set interesting constraints on such possibilities [153, 154].

References

- [1] R. V. Wagoner, W. A. Fowler and F. Hoyle, *Astrophys. J.* **148**, 3 (1967).
- [2] D. N. Schramm and M. S. Turner, *Rev. Mod. Phys.* **70**, 303 (1998), [[arXiv:astro-ph/9706069](#)].
- [3] G. Steigman, *Ann. Rev. Nucl. Part. Sci.* **57**, 463 (2007), [[arXiv:0712.1100](#)].
- [4] F. Iocco *et al.*, *Phys. Rept.* **472**, 1 (2009), [[arXiv:0809.0631](#)].
- [5] R. H. Cyburt *et al.*, *Rev. Mod. Phys.* **88**, 015004 (2016), [[arXiv:1505.01076](#)].
- [6] S. Sarkar, *Rept. on Prog. in Phys.* **59**, 1493 (1996).
- [7] K. Jedamzik and M. Pospelov, *New J. Phys.* **11**, 105028 (2009), [[arXiv:0906.2087](#)].
- [8] M. Pospelov and J. Pradler, *Ann. Rev. Nucl. Part. Sci.* **60**, 539 (2010), [[arXiv:1011.1054](#)].
- [9] B. D. Fields, *Ann. Rev. Nucl. Part. Sci.* **61**, 47 (2011), [[arXiv:1203.3551](#)].

- [10] P.J.E. Peebles, *Phys. Rev. Lett.* **16**, 411 (1966).
- [11] J. Bernstein, L. S. Brown and G. Feinberg, *Rev. Mod. Phys.* **61**, 25 (1989).
- [12] S. Mukhanov, *Int. J. Theor. Phys.* **143**, 669 (2004).
- [13] R. Esmailzadeh, G. D. Starkman and S. Dimopoulos, *Astrophys. J.* **378**, 504 (1991).
- [14] R. H. Cyburt, B. D. Fields and K. A. Olive, *JCAP* **0811**, 012 (2008), [[arXiv:0808.2818](#)].
- [15] R. H. Cyburt, B. D. Fields and K. A. Olive, *New Astron.* **6**, 215 (2001), [[arXiv:astro-ph/0102179](#)].
- [16] L. Kawano, Technical Report FERMILAB-PUB-92-004-A (1992), URL <https://ui.adsabs.harvard.edu/abs/1992STIN...9225163K>.
- [17] O. Pisanti *et al.*, *Comput. Phys. Commun.* **178**, 956 (2008), [[arXiv:0705.0290](#)].
- [18] A. Arbey, *Comput. Phys. Commun.* **183**, 1822 (2012), [[arXiv:1106.1363](#)].
- [19] R. Consiglio *et al.*, *Computer Physics Communications* **233**, 237 (2018), [[arXiv:1712.04378](#)].
- [20] S. Esposito *et al.*, *Nucl. Phys.* **B568**, 421 (2000), [[arXiv:astro-ph/9906232](#)].
- [21] S. Dodelson and M. S. Turner, *Phys. Rev.* **D46**, 3372 (1992).
- [22] D. Seckel (1993), [[hep-ph/9305311](#)].
- [23] M. S. Smith, L. H. Kawano and R. A. Malaney, *Astrophys. J. Suppl.* **85**, 219 (1993).
- [24] G. Fiorentini *et al.*, *Phys. Rev.* **D58**, 063506 (1998), [[arXiv:astro-ph/9803177](#)].
- [25] A. Coc *et al.*, *Astrophys. J.* **744**, 158 (2012), [[arXiv:1107.1117](#)].
- [26] K. M. Nollett and S. Burles, *Phys. Rev.* **D61**, 123505 (2000), [[arXiv:astro-ph/0001440](#)].
- [27] R. H. Cyburt, *Phys. Rev.* **D70**, 023505 (2004), [[arXiv:astro-ph/0401091](#)].
- [28] P. D. Serpico *et al.*, *JCAP* **0412**, 010 (2004), [[arXiv:astro-ph/0408076](#)].
- [29] R. N. Boyd *et al.*, *Phys. Rev.* **D82**, 105005 (2010), [[arXiv:1008.0848](#)].
- [30] R.H. Cyburt and B. Davids, *Phys. Rev.* **C78**, 012 (2008).
- [31] S. Burles, K. M. Nollett and M. S. Turner, *Astrophys. J.* **552**, L1 (2001), [[arXiv:astro-ph/0010171](#)].
- [32] R. I. Epstein, J. M. Lattimer and D. N. Schramm, *Nature* **263**, 198 (1976).
- [33] S. D’Odorico *et al.*, *Astron. & Astrophys.* **368**, L21 (2001).
- [34] D. Romano *et al.*, *Mon. Not. Roy. Astron. Soc.* **369**, 295 (2006), [[arXiv:astro-ph/0603190](#)].
- [35] M. Pettini and D. V. Bowen, *Astrophys. J.* **560**, 41 (2001), [[arXiv:astro-ph/0104474](#)].
- [36] R. Cooke *et al.*, *Astrophys. J.* **781**, 1, 31 (2014), [[arXiv:1308.3240](#)].
- [37] R. J. Cooke *et al.*, *Astrophys. J.* **830**, 2, 148 (2016), [[arXiv:1607.03900](#)].
- [38] E.O. Zavarygin *et al.*, [arXiv:1706.09512](#)(2017).
- [39] S. A. Balashev *et al.*, *Mon. Not. Roy. Astron. Soc.* **458**, 2, 2188 (2016), [[arXiv:1511.01797](#)].
- [40] S. A. Levshakov *et al.*, *Astrophys. J.* **565**, 696 (2002), [[arXiv:astro-ph/0105529](#)].
- [41] M. Fumagalli, J. M. O’Meara and J. X. Prochaska, *Science* **334**, 1245 (2011), [[arXiv:1111.2334](#)].
- [42] R. Srianand *et al.*, *Mon. Not. Roy. Astron. Soc.* **405**, 1888 (2010), [[arXiv:1002.4620](#)].
- [43] P. Noterdaeme *et al.*, *Astron. & Astrophys.* **542**, L33 (2012), [[arXiv:1205.3777](#)].
- [44] M. Pettini and R. Cooke, *Mon. Not. Roy. Astron. Soc.* **425**, 2477 (2012), [[arXiv:1205.3785](#)].
- [45] S. Riemer-Sørensen *et al.*, *Mon. Not. Roy. Astron. Soc.* **447**, 2925 (2015), [[arXiv:1412.4043](#)].

- [46] S. Riemer-Sørensen *et al.*, *Mon. Not. Roy. Astron. Soc.* **468**, 3, 3239 (2017), [[arXiv:1703.06656](#)].
- [47] B. D. Fields *et al.*, *JCAP* (2019).
- [48] K. M. Nollett and G. P. Holder (2011), [[arXiv:1112.2683](#)].
- [49] L. E. Marcucci *et al.*, *Phys. Rev. Lett.* **116**, 10, 102501 (2016), [Erratum: *Phys. Rev. Lett.*117,no.4,049901(2016)], [[arXiv:1510.07877](#)].
- [50] C. Gustavino, *Euro. Phys. J. Web of Conferences*, 136, 01009 (2017).
- [51] R. J. Cooke, M. Pettini and C. C. Steidel, *Astrophys. J.* **855**, 2, 102 (2018), [[arXiv:1710.11129](#)].
- [52] E. O. Zavarygin *et al.*, *Mon. Not. R. Astron. Soc* **477**, 4, 5536 (2018), [[arXiv:1706.09512](#)].
- [53] F. van de Voort *et al.*, *Mon. Not. R. Astron. Soc* **477**, 1, 80 (2018), [[arXiv:1704.08254](#)].
- [54] J. L. Linsky *et al.*, *Astrophys. J.* **647**, 1106 (2006), [[arXiv:astro-ph/0608308](#)].
- [55] M. Pettini and D. V. Bowen, *Astrophys. J.* **560**, 1, 41 (2001), [[arXiv:astro-ph/0104474](#)].
- [56] S. D’Odorico, M. Dessauges-Zavadsky and P. Molaro, *Astron. & Astrophys.* **368**, L21 (2001), [[arXiv:astro-ph/0102162](#)].
- [57] R. Srianand *et al.*, *Mon. Not. R. Astron. Soc* **405**, 3, 1888 (2010), [[arXiv:1002.4620](#)].
- [58] S. Riemer-Sørensen *et al.*, *Mon. Not. R. Astron. Soc* **447**, 3, 2925 (2015), [[arXiv:1412.4043](#)].
- [59] Y. I. Izotov, T. X. Thuan and N. G. Guseva, *Mon. Not. Roy. Astron. Soc.* **445**, 1, 778 (2014), [[arXiv:1408.6953](#)].
- [60] Y. I. Izotov *et al.*, *Astrophys. J.* **527**, 757 (1999), [[arXiv:astro-ph/9907228](#)].
- [61] K. A. Olive and E. D. Skillman, *Astrophys. J.* **617**, 29 (2004), [[arXiv:astro-ph/0405588](#)].
- [62] M. Valerdi *et al.*, *Astrophys. J.* **876**, 2, 98 (2019), [[arXiv:1904.01594](#)].
- [63] E. Aver, K. A. Olive and E. D. Skillman, *JCAP* **1507**, 07, 011 (2015), [[arXiv:1503.08146](#)].
- [64] Y.I. Izotov *et al.*, *Astron. & Astrophys.* **558**, A57 (2013).
- [65] D. S. Aguado *et al.*, *Astrophys. J. Lett.* **874**, 2, L21 (2019), [[arXiv:1904.04892](#)].
- [66] M. Spite and F. Spite, *Nature* **297**, 483 (1982).
- [67] E. Vangioni-Flam *et al.*, *New Astron.* **4**, 245 (1999), [[arXiv:astro-ph/9811327](#)].
- [68] S. G. Ryan *et al.*, *Astrophys. J.* **530**, L57 (2000), [[arXiv:astro-ph/9905211](#)].
- [69] P. Bonifacio and P. Molaro, *Mon. Not. Roy. Astron. Soc.* **285**, 847 (1997), [[arXiv:astro-ph/9611043](#)].
- [70] R. Rebolo, P. Molaro and J. E. Beckman, *Astron. & Astrophys.* **192**, 192 (1988).
- [71] P. Bonifacio *et al.*, *Astron. & Astrophys.* **462**, 851 (2007).
- [72] W. Aoki *et al.*, *Astrophys. J.* **698**, 1803 (2009), [[arXiv:0904.1448](#)].
- [73] L. Sbordone *et al.*, *Astron. & Astrophys.* **522**, A26 (2010).
- [74] P. Bonifacio *et al.*, *Astron. & Astrophys.* **612**, A65 (2018), [[arXiv:1801.03935](#)].
- [75] P. Bonifacio *et al.*, *Astron. & Astrophys.* **390**, 91 (2002), [[arXiv:astro-ph/0204332](#)].
- [76] J. Melendez *et al.*, *Astron. & Astrophys.* **515**, L3 (2010).
- [77] M.H. Pinsonneault *et al.*, *Astrophys. J.* **574**, 389 (2002).
- [78] A. J. Korn *et al.*, *Nature* **442**, 657 (2006), [[arXiv:astro-ph/0608201](#)].
- [79] P. Molaro *et al.*, *Memorie della Soc. Astronomica Italiana Supp.* **22**, 233 (2012).

- [80] X. Fu *et al.*, Mon. Not. R. Astron. Soc **452**, 325 (2015).
- [81] M. Asplund *et al.*, Astrophys. J. **644**, 229 (2006), [[arXiv:astro-ph/0510636](#)].
- [82] K. Lind *et al.*, Astron. & Astrophys. **554**, 96 (2013).
- [83] R. Cayrel *et al.*, Astron. & Astrophys. **473**, L37 (2007).
- [84] M. Steffen *et al.*, Memorie della Soc. Astronomica Italiana Supp. **22**, 152 (2012).
- [85] D. S. Balsler and T. M. Bania, Astrophys. J. **156**, 6, 280 (2018), [[arXiv:1810.09422](#)].
- [86] H. Reeves *et al.*, Astrophys. J. **179**, 909 (1973).
- [87] M. Fukugita and P. J. E. Peebles, Astrophys. J. **616**, 643 (2004), [[arXiv:astro-ph/0406095](#)].
- [88] R. Cen and J. P. Ostriker, Astrophys. J. **514**, 1 (1999), [[arXiv:astro-ph/9806281](#)].
- [89] G. Jungman *et al.*, Phys. Rev. **D54**, 1332 (1996), [[arXiv:astro-ph/9512139](#)].
- [90] Planck Collaboration *et al.*, arXiv e-prints arXiv:1807.06209 (2018), [[arXiv:1807.06209](#)].
- [91] P.A.R. Ade *et al.*, Astron. & Astrophys. **594**, A13 (2016).
- [92] G. Steigman, D. N. Schramm and J. E. Gunn, Phys. Lett. **B66**, 202 (1977), [,159(1977)].
- [93] E. Lisi, S. Sarkar and F. L. Villante, Phys. Rev. **D59**, 123520 (1999), [[hep-ph/9901404](#)].
- [94] R. Trotta and S. H. Hansen, Phys. Rev. **D69**, 023509 (2004), [[arXiv:astro-ph/0306588](#)].
- [95] R. H. Cyburt, B. D. Fields and K. A. Olive, Phys. Lett. **B567**, 227 (2003), [[arXiv:astro-ph/0302431](#)].
- [96] C. Pitrou *et al.*, Phys. Rept. **754**, 1 (2018), [[arXiv:1801.08023](#)].
- [97] R. H. Cyburt *et al.*, JCAP **1305**, 014 (2013), [[arXiv:1303.0574](#)].
- [98] N. Chakraborty, B. D. Fields and K. A. Olive, Phys. Rev. **D83**, 063006 (2011), [[arXiv:1011.0722](#)].
- [99] C. Brogini *et al.*, JCAP **1206**, 030 (2012), [[arXiv:1202.5232](#)].
- [100] P. D. O'Malley *et al.*, Phys. Rev. **C84**, 042801 (2011).
- [101] F. Hammache *et al.*, Phys. Rev. **C88**, 6, 062802 (2013), [[arXiv:1312.0894](#)].
- [102] M. W. Paris *et al.*, Nucl. Data Sheets **120**, 184 (2014), [[arXiv:1304.3153](#)].
- [103] O. Richard, G. Michaud and J. Richer, Astrophys. J. **619**, 538 (2005), [[arXiv:astro-ph/0409672](#)].
- [104] P. Gruyters *et al.*, Astron. Astrophys. **555**, 31 (2013).
- [105] K. Jedamzik and J. B. Rehm, Phys. Rev. **D64**, 023510 (2001), [[arXiv:astro-ph/0101292](#)].
- [106] D. J. Fixsen *et al.*, Astrophys. J. **473**, 576 (1996), [[arXiv:astro-ph/9605054](#)].
- [107] M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. **D71**, 083502 (2005), [[arXiv:astro-ph/0408426](#)].
- [108] K. Jedamzik, Phys. Rev. **D70**, 063524 (2004), [[arXiv:astro-ph/0402344](#)].
- [109] A. Goudelis, M. Pospelov and J. Pradler, Phys. Rev. Lett. **116**, 21, 211303 (2016), [[arXiv:1510.08858](#)].
- [110] J. D. Barrow, Phys. Rev. **D35**, 1805 (1987).
- [111] B. A. Campbell and K. A. Olive, Phys. Lett. **B345**, 429 (1995), [[hep-ph/9411272](#)].
- [112] L. Bergstrom, S. Iguri and H. Rubinstein, Phys. Rev. **D60**, 045005 (1999), [[arXiv:astro-ph/9902157](#)].
- [113] V. V. Flambaum and E. V. Shuryak, Phys. Rev. **D65**, 103503 (2002), [[hep-ph/0201303](#)].

- [114] A. Coc *et al.*, Phys. Rev. **D76**, 023511 (2007), [[arXiv:astro-ph/0610733](#)].
- [115] J. C. Berengut *et al.*, Phys. Rev. **D87**, 8, 085018 (2013), [[arXiv:1301.1738](#)].
- [116] Z. Hou *et al.*, Phys. Rev. **D87**, 083008 (2013), [[arXiv:1104.2333](#)].
- [117] R. Keisler *et al.*, Astrophys. J. **743**, 28 (2011), [[arXiv:1105.3182](#)].
- [118] J. Hamann *et al.*, Phys. Rev. Lett. **105**, 181301 (2010), [[arXiv:1006.5276](#)].
- [119] F. S. Accetta, L. M. Krauss and P. Romanelli, Phys. Lett. **B248**, 146 (1990).
- [120] K. M. Nollett and R. E. Lopez, Phys. Rev. **D66**, 063507 (2002), [[arXiv:astro-ph/0204325](#)].
- [121] C. Bambi, M. Giannotti and F. L. Villante, Phys. Rev. **D71**, 123524 (2005), [[arXiv:astro-ph/0503502](#)].
- [122] E. W. Kolb, M. J. Perry and T. P. Walker, Phys. Rev. **D33**, 869 (1986).
- [123] A. Coc *et al.*, Phys. Rev. **D73**, 083525 (2006), [[arXiv:astro-ph/0601299](#)].
- [124] K. A. Olive, D. N. Schramm and G. Steigman, Nucl. Phys. **B180**, 497 (1981).
- [125] J. R. Ellis *et al.*, Phys. Lett. **167B**, 457 (1986).
- [126] L. M. Krauss, J. Terning and T. Appelquist, Phys. Rev. Lett. **71**, 823 (1993), [[hep-ph/9305265](#)].
- [127] J. A. Morgan, Phys. Lett. **102B**, 247 (1981).
- [128] E. W. Kolb, M. S. Turner and T. P. Walker, Phys. Rev. **D34**, 2197 (1986).
- [129] J. A. Grifols and E. Masso, Mod. Phys. Lett. **A2**, 205 (1987).
- [130] K. S. Babu, R. N. Mohapatra and I. Z. Rothstein, Phys. Rev. Lett. **67**, 545 (1991).
- [131] A. D. Dolgov, S. H. Hansen and D. V. Semikoz, Nucl. Phys. **B524**, 621 (1998), [[hep-ph/9712284](#)].
- [132] K. Enqvist, K. Kainulainen and M. J. Thomson, Nucl. Phys. **B373**, 498 (1992).
- [133] A. D. Dolgov, Phys. Rept. **370**, 333 (2002), [[hep-ph/0202122](#)].
- [134] J. A. Grifols, R. N. Mohapatra and A. Riotto, Phys. Lett. **B400**, 124 (1997), [[hep-ph/9612253](#)].
- [135] H. K. Dreiner *et al.*, Phys. Rev. **D85**, 065027 (2012), [[arXiv:1111.5715](#)].
- [136] K. Sato and M. Kobayashi, Prog. Theor. Phys. **58**, 1775 (1977).
- [137] D. A. Dicus *et al.*, Phys. Rev. **D17**, 1529 (1978).
- [138] R. J. Scherrer and M. S. Turner, Astrophys. J. **331**, 19 (1988), [[Astrophys. J.331,33\(1988\)](#)].
- [139] D. Lindley, Mon. Not. R. Astron. Soc **188**, 15 (1979).
- [140] J. R. Ellis *et al.*, Nucl. Phys. **B373**, 399 (1992).
- [141] M. H. Reno and D. Seckel, Phys. Rev. **D37**, 3441 (1988).
- [142] S. Dimopoulos *et al.*, Nucl. Phys. **B311**, 699 (1989).
- [143] S. Sarkar and A. M. Cooper-Sarkar, Phys. Lett. **148B**, 347 (1984), [[I.362\(1984\)](#)].
- [144] M. Kawasaki, K. Kohri and T. Moroi, Phys. Lett. **B649**, 436 (2007), [[hep-ph/0703122](#)].
- [145] K. Jedamzik *et al.*, JCAP **0607**, 007 (2006), [[hep-ph/0512044](#)].
- [146] S. Davidson *et al.*, Phys. Rev. **466**, 105 (2008).
- [147] S. Weinberg, Phys. Rev. Lett. **48**, 1303 (1982).
- [148] R. H. Cyburt *et al.*, Phys. Rev. **D67**, 103521 (2003), [[arXiv:astro-ph/0211258](#)].

- [149] M. Bolz, A. Brandenburg and W. Buchmuller, Nucl. Phys. **B606**, 518 (2001), [Erratum: Nucl. Phys.B790,336(2008)], [[hep-ph/0012052](#)].
- [150] G. D. Coughlan *et al.*, Physics Letters B **131**, 1-3, 59 (1983).
- [151] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Rev. **D59**, 086004 (1999), [[hep-ph/9807344](#)].
- [152] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999), [[hep-ph/9905221](#)].
- [153] J. M. Cline, C. Grojean and G. Servant, Phys. Rev. Lett. **83**, 4245 (1999), [[hep-ph/9906523](#)].
- [154] P. Binetruy *et al.*, Phys. Lett. **B477**, 285 (2000), [[hep-th/9910219](#)].