

CHARMED MESONS ($C = \pm 1$)

$D^+ = c\bar{d}$, $D^0 = c\bar{u}$, $\bar{D}^0 = \bar{c}u$, $D^- = \bar{c}d$, similarly for D^* 's

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1869.65 \pm 0.05$ MeV

Mean life $\tau = (1040 \pm 7) \times 10^{-15}$ s

$$c\tau = 311.8 \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{anything})/\Gamma(c \rightarrow \text{anything}) = 0.096 \pm 0.004 \quad [a]$$

$$\Gamma(c \rightarrow D^*(2010)^+ \text{anything})/\Gamma(c \rightarrow \text{anything}) = 0.255 \pm 0.017$$

CP -violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_L^0 e^\pm \nu) = (-0.6 \pm 1.6)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.41 \pm 0.09)\%$$

$$A_{CP}(K_L^0 K^\pm) \text{ in } D^\pm \rightarrow K_L^0 K^\pm = (-4.2 \pm 3.4) \times 10^{-2}$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.18 \pm 0.16)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (-0.3 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (-0.1 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.0 \pm 1.2)\%$$

$$A_{CP}(\pi^\pm \pi^0) = (2.4 \pm 1.2)\%$$

$$A_{CP}(\pi^\pm \eta) = (1.0 \pm 1.5)\% \quad (S = 1.4)$$

$$A_{CP}(\pi^\pm \eta'(958)) = (-0.6 \pm 0.7)\%$$

$$A_{CP}(\bar{K}^0 / K^0 K^\pm) = (0.11 \pm 0.17)\%$$

$$A_{CP}(K_S^0 K^\pm) = (-0.01 \pm 0.07)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_S^0 K^\pm \pi^0 = (1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K_L^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_L^0 K^\pm \pi^0 = (-1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.37 \pm 0.29)\%$$

$$A_{CP}(K^\pm K^{*0}) = (-0.3 \pm 0.4)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.01 \pm 0.09)\% \quad (S = 1.8)$$

$$A_{CP}(K^\pm K_0^*(1430)^0) = (8^{+7}_{-6})\%$$

$$A_{CP}(K^\pm K_2^*(1430)^0) = (43^{+20}_{-26})\%$$

$$A_{CP}(K^\pm K_0^*(700)) = (-12^{+18}_{-13})\%$$

$$A_{CP}(a_0(1450)^0 \pi^\pm) = (-19^{+14}_{-16})\%$$

$$A_{CP}(\phi(1680) \pi^\pm) = (-9 \pm 26)\%$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = (-2 \pm 4)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) = (-4 \pm 7)\%$$

$$A_{CP}(K^\pm \pi^0) = (-4 \pm 11)\%$$

χ^2 tests of CP -violation (CPV)

Local CPV in $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm = 78.1\%$

Local CPV in $D^\pm \rightarrow K^+ K^- \pi^\pm = 31\%$

CP violating asymmetries of P -odd (T -odd) moments

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-12 \pm 11) \times 10^{-3} [b]$$

D^+ form factors

$$f_+(0)|V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell = 0.719 \pm 0.011 \quad (S = 1.6)$$

$$r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell = -2.13 \pm 0.14$$

$$r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell = -3 \pm 12 \quad (S = 1.5)$$

$$f_+(0)|V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell = 0.1407 \pm 0.0025$$

$$r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell = -2.00 \pm 0.13$$

$$r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell = -4 \pm 5$$

$$f_+(0)|V_{cd}| \text{ in } D^+ \rightarrow \eta e^+ \nu_e = (8.3 \pm 0.5) \times 10^{-2}$$

$$r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e = -5.3 \pm 2.7 \quad (S = 1.9)$$

$$r_v \equiv V(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e = 1.24 \pm 0.11$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e = 1.06 \pm 0.16$$

$$r_v \equiv V(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e = 1.64 \pm 0.10 \quad (S = 1.2)$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e = 0.84 \pm 0.06$$

$$r_v \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 1.49 \pm 0.05 \quad (S = 2.1)$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.802 \pm 0.021$$

$$r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.0 \pm 0.4$$

$$\Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 1.13 \pm 0.08$$

$$\Gamma_+/ \Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.22 \pm 0.06 \quad (S = 1.6)$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07 \pm 0.30) %		—
μ^+ anything	(17.6 \pm 3.2) %		—
K^- anything	(25.7 \pm 1.4) %		—
\bar{K}^0 anything + K^0 anything	(61 \pm 5) %		—
K^+ anything	(5.9 \pm 0.8) %		—

$K^*(892)^-$ anything	(6 \pm 5) %	-
$K^*(892)^0$ anything	(23 \pm 5) %	-
$K^*(892)^0$ anything	< 6.6 %	CL=90%
η anything	(6.3 \pm 0.7) %	-
η' anything	(1.04 \pm 0.18) %	-
ϕ anything	(1.12 \pm 0.04) %	-

Leptonic and semileptonic modes

$e^+ \nu_e$	< 8.8 $\times 10^{-6}$ CL=90%	935
$\gamma e^+ \nu_e$	< 3.0 $\times 10^{-5}$ CL=90%	935
$\mu^+ \nu_\mu$	(3.74 \pm 0.17) $\times 10^{-4}$	932
$\tau^+ \nu_\tau$	(1.20 \pm 0.27) $\times 10^{-3}$	90
$\overline{K}^0 e^+ \nu_e$	(8.73 \pm 0.10) %	869
$\overline{K}^0 \mu^+ \nu_\mu$	(8.76 \pm 0.19) %	865
$K^- \pi^+ e^+ \nu_e$	(4.02 \pm 0.18) %	S=3.2
$\overline{K}^*(892)^0 e^+ \nu_e, \overline{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.77 \pm 0.17) %	722
$(K^- \pi^+) [0.8-1.0]\text{GeV} e^+ \nu_e$	(3.39 \pm 0.09) %	864
$(K^- \pi^+)_{S-wave} e^+ \nu_e$	(2.28 \pm 0.11) $\times 10^{-3}$	-
$\overline{K}^*(1410)^0 e^+ \nu_e, \overline{K}^*(1410)^0 \rightarrow K^- \pi^+$	< 6 $\times 10^{-3}$ CL=90%	-
$\overline{K}_2^*(1430)^0 e^+ \nu_e, \overline{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	< 5 $\times 10^{-4}$ CL=90%	-
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 $\times 10^{-3}$ CL=90%	864
$\overline{K}^*(892)^0 e^+ \nu_e$	(5.40 \pm 0.10) %	S=1.1
$K^- \pi^+ \mu^+ \nu_\mu$	(3.65 \pm 0.34) %	851
$\overline{K}^*(892)^0 \mu^+ \nu_\mu, \overline{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.52 \pm 0.10) %	717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(1.9 \pm 0.5) $\times 10^{-3}$	851
$\overline{K}^*(892)^0 \mu^+ \nu_\mu$	(5.27 \pm 0.15) %	717
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.5 $\times 10^{-3}$ CL=90%	825
$\overline{K}_1(1270)^0 e^+ \nu_e, \overline{K}_1^0 \rightarrow K^- \pi^+ \pi^0$	(1.06 \pm 0.15) $\times 10^{-3}$	-
$\overline{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.3 $\times 10^{-4}$ CL=90%	380
$\overline{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.5 $\times 10^{-3}$ CL=90%	105
$\pi^0 e^+ \nu_e$	(3.72 \pm 0.17) $\times 10^{-3}$	S=2.0
$\pi^0 \mu^+ \nu_\mu$	(3.50 \pm 0.15) $\times 10^{-3}$	927
$\eta e^+ \nu_e$	(1.11 \pm 0.07) $\times 10^{-3}$	855
$\pi^- \pi^+ e^+ \nu_e$	(2.45 \pm 0.10) $\times 10^{-3}$	924
$f_0(500)^0 e^+ \nu_e, f_0(500)^0 \rightarrow \pi^+ \pi^-$	(6.3 \pm 0.5) $\times 10^{-4}$	-
$\rho^0 e^+ \nu_e$	(2.18 \pm 0.17) $\times 10^{-3}$	774
$\rho^0 \mu^+ \nu_\mu$	(2.4 \pm 0.4) $\times 10^{-3}$	770

$\omega e^+ \nu_e$	(1.69 \pm 0.11) $\times 10^{-3}$	771
$\eta'(958) e^+ \nu_e$	(2.0 \pm 0.4) $\times 10^{-4}$	690
$a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0$	(1.7 \pm 0.8) $\times 10^{-4}$	-
$\phi e^+ \nu_e$	< 1.3 $\times 10^{-5}$ CL=90%	657
$D^0 e^+ \nu_e$	< 1.0 $\times 10^{-4}$ CL=90%	5
Hadronic modes with a \bar{K} or $\bar{K} K \bar{K}$		
$K_S^0 \pi^+$	(1.562 \pm 0.031) %	S=1.7
$K_L^0 \pi^+$	(1.46 \pm 0.05) %	863
$K^- 2\pi^+$	[c] (9.38 \pm 0.16) %	S=1.6
$(K^- \pi^+)_{S-\text{wave}} \pi^+$	(7.52 \pm 0.17) %	846
$\bar{K}_0^*(1430)^0 \pi^+,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[d] (1.25 \pm 0.06) %	382
$\bar{K}^*(892)^0 \pi^+,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.04 \pm 0.12) %	714
$\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	not seen	381
$\bar{K}_2^*(1430)^0 \pi^+,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[d] (2.3 \pm 0.7) $\times 10^{-4}$	371
$\bar{K}^*(1680)^0 \pi^+,$ $\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[d] (2.2 \pm 1.1) $\times 10^{-4}$	58
$K^-(2\pi^+)_{I=2}$	(1.45 \pm 0.26) %	-
$K_S^0 \pi^+ \pi^0$	[c] (7.36 \pm 0.21) %	845
$K_S^0 \rho^+$	(6.14 \pm 0.60) %	677
$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	(1.5 \pm 1.2) $\times 10^{-3}$	-
$\bar{K}^*(892)^0 \pi^+,$ $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	(2.64 \pm 0.32) $\times 10^{-3}$	714
$\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^{*0} \rightarrow$ $K_S^0 \pi^0$	(2.7 \pm 0.9) $\times 10^{-3}$	-
$\bar{K}_0^*(1680)^0 \pi^+, \bar{K}_0^{*0} \rightarrow$ $K_S^0 \pi^0$	(10 \pm 7) $\times 10^{-4}$	-
$\bar{\kappa}^0 \pi^+, \bar{\kappa}^0 \rightarrow K_S^0 \pi^0$	(6 \pm 5) $\times 10^{-3}$	-
$K_S^0 \pi^+ \pi^0$ nonresonant	(3 \pm 4) $\times 10^{-3}$	845
$K_S^0 \pi^+ \pi^0$ nonresonant and $\bar{\kappa}^0 \pi^+$	(1.37 \pm 0.21) %	-
$(K_S^0 \pi^0)_{S-\text{wave}} \pi^+$	(1.27 \pm 0.27) %	845
$K_S^0 \pi^+ \eta'(958)$	(1.90 \pm 0.21) $\times 10^{-3}$	481
$K^- 2\pi^+ \pi^0$	[e] (6.25 \pm 0.18) %	816
$K_S^0 2\pi^+ \pi^-$	[e] (3.10 \pm 0.09) %	814

$K^- 3\pi^+ \pi^-$	[c]	(5.7 \pm 0.5) $\times 10^{-3}$	S=1.1	772
$\overline{K}^*(892)^0 2\pi^+ \pi^-$,		(1.2 \pm 0.4) $\times 10^{-3}$		645
$\overline{K}^*(892)^0 \rightarrow K^- \pi^+$				
$\overline{K}^*(892)^0 \rho^0 \pi^+$,		(2.3 \pm 0.4) $\times 10^{-3}$		239
$\overline{K}^*(892)^0 \rightarrow K^- \pi^+$				
$\overline{K}^*(892)^0 a_1(1260)^+$	[f]	(9.3 \pm 1.9) $\times 10^{-3}$		†
$K^- \rho^0 2\pi^+$		(1.72 \pm 0.28) $\times 10^{-3}$		524
$K^- 3\pi^+ \pi^-$ nonresonant		(4.0 \pm 2.9) $\times 10^{-4}$		772
$K^+ 2K_S^0$		(2.54 \pm 0.13) $\times 10^{-3}$		545
$K^+ K^- K_S^0 \pi^+$		(2.4 \pm 0.5) $\times 10^{-4}$		436
Pionic modes				
$\pi^+ \pi^0$		(1.247 \pm 0.033) $\times 10^{-3}$		925
$2\pi^+ \pi^-$		(3.27 \pm 0.18) $\times 10^{-3}$		909
$\rho^0 \pi^+$		(8.3 \pm 1.5) $\times 10^{-4}$		767
$\pi^+ (\pi^+ \pi^-)_{S-\text{wave}}$		(1.83 \pm 0.16) $\times 10^{-3}$		909
$\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-$		(1.38 \pm 0.12) $\times 10^{-3}$		—
$f_0(980) \pi^+$,		(1.56 \pm 0.33) $\times 10^{-4}$		669
$f_0(980) \rightarrow \pi^+ \pi^-$				
$f_0(1370) \pi^+$,		(8 \pm 4) $\times 10^{-5}$		—
$f_0(1370) \rightarrow \pi^+ \pi^-$				
$f_2(1270) \pi^+$,		(5.0 \pm 0.9) $\times 10^{-4}$		485
$f_2(1270) \rightarrow \pi^+ \pi^-$				
$\rho(1450)^0 \pi^+$,		< 8 $\times 10^{-5}$ CL=95%		338
$\rho(1450)^0 \rightarrow \pi^+ \pi^-$				
$f_0(1500) \pi^+$,		(1.1 \pm 0.4) $\times 10^{-4}$		—
$f_0(1500) \rightarrow \pi^+ \pi^-$				
$f_0(1710) \pi^+$,		< 5 $\times 10^{-5}$ CL=95%		—
$f_0(1710) \rightarrow \pi^+ \pi^-$				
$f_0(1790) \pi^+$,		< 7 $\times 10^{-5}$ CL=95%		—
$f_0(1790) \rightarrow \pi^+ \pi^-$				
$(\pi^+ \pi^+)_{S-\text{wave}} \pi^-$		< 1.2 $\times 10^{-4}$ CL=95%		909
$2\pi^+ \pi^-$ nonresonant		< 1.1 $\times 10^{-4}$ CL=95%		909
$\pi^+ 2\pi^0$		(4.7 \pm 0.4) $\times 10^{-3}$		910
$2\pi^+ \pi^- \pi^0$		(1.16 \pm 0.08) %		883
$3\pi^+ 2\pi^-$		(1.66 \pm 0.16) $\times 10^{-3}$	S=1.1	845
$\eta \pi^+$				
$\eta \pi^+ \pi^0$		(3.77 \pm 0.09) $\times 10^{-3}$		848
$\omega \pi^+$		(1.38 \pm 0.35) $\times 10^{-3}$		831
$\eta'(958) \pi^+$		(2.8 \pm 0.6) $\times 10^{-4}$		764
$\eta'(958) \pi^+ \pi^0$		(4.97 \pm 0.19) $\times 10^{-3}$		681
		(1.6 \pm 0.5) $\times 10^{-3}$		654

Hadronic modes with a $K\bar{K}$ pair

$K^+ K_S^0$	$(3.04 \pm 0.09) \times 10^{-3}$	S=2.2	793
$K_L^0 K^+$	$(3.21 \pm 0.16) \times 10^{-3}$		793
$K_S^0 K^+ \pi^0$	$(5.07 \pm 0.30) \times 10^{-3}$		744
$K_L^0 K^+ \pi^0$	$(5.24 \pm 0.31) \times 10^{-3}$		744
$K^+ K^- \pi^+$	[c] $(9.68 \pm 0.18) \times 10^{-3}$		744
$\phi \pi^+$	$(5.70 \pm 0.14) \times 10^{-3}$		647
$\phi \pi^+, \phi \rightarrow K^+ K^-$	$(2.69 \pm 0.07) \times 10^{-3}$		647
$K^+ \bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(2.49 \pm 0.08) \times 10^{-3}$		613
$K^+ \bar{K}_0^*(1430)^0,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	$(1.82 \pm 0.35) \times 10^{-3}$		-
$K^+ \bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow$ $K^- \pi^+$	$(1.6 \pm 1.2) \times 10^{-4}$		-
$K^+ \bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^- \pi^+$	$(6.8 \pm 3.5) \times 10^{-4}$		-
$a_0(1450)^0 \pi^+, a_0^0 \rightarrow$ $K^+ K^-$	$(4.5 \pm 7.0) \times 10^{-4}$		-
$\phi(1680) \pi^+, \phi \rightarrow K^+ K^-$	$(4.9 \pm 4.0) \times 10^{-5}$		-
$K_S^0 K_S^0 \pi^+$	$(2.70 \pm 0.13) \times 10^{-3}$		741
$K^+ K_S^0 \pi^+ \pi^-$	$(1.74 \pm 0.18) \times 10^{-3}$		678
$K_S^0 K^- 2\pi^+$	$(2.38 \pm 0.17) \times 10^{-3}$		678
$K^+ K^- 2\pi^+ \pi^-$	$(2.3 \pm 1.2) \times 10^{-4}$		601

A few poorly measured branching fractions:

$\phi \pi^+ \pi^0$	$(2.3 \pm 1.0) \%$		619
$\phi \rho^+$	$< 1.5 \quad \% \quad \text{CL}=90\%$		260
$K^+ K^- \pi^+ \pi^0 \text{non-}\phi$	$(1.5 \pm 0.7) \%$		682
$K^*(892)^+ K_S^0$	$(1.7 \pm 0.8) \%$		612

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(2.08 \pm 0.21) \times 10^{-4}$	S=1.4	864
$K^+ \eta$	$(1.25 \pm 0.16) \times 10^{-4}$	S=1.1	776
$K^+ \eta'(958)$	$(1.85 \pm 0.20) \times 10^{-4}$		571
$K^+ \pi^+ \pi^-$	$(4.91 \pm 0.09) \times 10^{-4}$		846
$K^+ \rho^0$	$(1.9 \pm 0.5) \times 10^{-4}$		679
$K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow$ $K^+ \pi^-$	$(2.3 \pm 0.4) \times 10^{-4}$		714
$K^+ f_0(980), f_0(980) \rightarrow$ $\pi^+ \pi^-$	$(4.4 \pm 2.6) \times 10^{-5}$		-

$K_2^*(1430)^0 \pi^+$, $K_2^*(1430)^0 \rightarrow K^+ \pi^-$		(3.9 \pm 2.7) $\times 10^{-5}$	-
$K^+ \pi^-$ nonresonant		not seen	846
$2K^+ K^-$		(6.14 \pm 0.11) $\times 10^{-5}$	550
$\phi(1020)^0 K^+$		< 2.1 $\times 10^{-5}$ CL=90%	-
$K^+ \phi(1020)$, $\phi \rightarrow K^+ K^-$		(4.4 \pm 0.6) $\times 10^{-6}$	-
$K^+(K^+ K^-)$ <i>S-wave</i>		(5.77 \pm 0.12) $\times 10^{-5}$	550
$\Delta C = 1$ weak neutral current (<i>C1</i>) modes, or Lepton Family number (<i>LF</i>) or Lepton number (<i>L</i>) violating modes			
$\pi^+ e^+ e^-$	<i>C1</i>	< 1.1 $\times 10^{-6}$ CL=90%	930
$\pi^+ \pi^0 e^+ e^-$		< 1.4 $\times 10^{-5}$ CL=90%	925
$\pi^+ \phi$, $\phi \rightarrow e^+ e^-$	[<i>g</i>]	(1.7 \pm 1.4) $\times 10^{-6}$	-
$\pi^+ \mu^+ \mu^-$	<i>C1</i>	< 7.3 $\times 10^{-8}$ CL=90%	918
$\pi^+ \phi$, $\phi \rightarrow \mu^+ \mu^-$	[<i>g</i>]	(1.8 \pm 0.8) $\times 10^{-6}$	-
$\rho^+ \mu^+ \mu^-$	<i>C1</i>	< 5.6 $\times 10^{-4}$ CL=90%	757
$K^+ e^+ e^-$	[<i>h</i>]	< 1.0 $\times 10^{-6}$ CL=90%	870
$K^+ \pi^0 e^+ e^-$		< 1.5 $\times 10^{-5}$ CL=90%	864
$K_S^0 \pi^+ e^+ e^-$		< 2.6 $\times 10^{-5}$ CL=90%	-
$K_S^0 K^+ e^+ e^-$		< 1.1 $\times 10^{-5}$ CL=90%	-
$K^+ \mu^+ \mu^-$	[<i>h</i>]	< 4.3 $\times 10^{-6}$ CL=90%	856
$\pi^+ e^+ \mu^-$	<i>LF</i>	< 2.9 $\times 10^{-6}$ CL=90%	927
$\pi^+ e^- \mu^+$	<i>LF</i>	< 3.6 $\times 10^{-6}$ CL=90%	927
$K^+ e^+ \mu^-$	<i>LF</i>	< 1.2 $\times 10^{-6}$ CL=90%	866
$K^+ e^- \mu^+$	<i>LF</i>	< 2.8 $\times 10^{-6}$ CL=90%	866
$\pi^- 2e^+$	<i>L</i>	< 1.1 $\times 10^{-6}$ CL=90%	930
$\pi^- 2\mu^+$	<i>L</i>	< 2.2 $\times 10^{-8}$ CL=90%	918
$\pi^- e^+ \mu^+$	<i>L</i>	< 2.0 $\times 10^{-6}$ CL=90%	927
$\rho^- 2\mu^+$	<i>L</i>	< 5.6 $\times 10^{-4}$ CL=90%	757
$K^- 2e^+$	<i>L</i>	< 9 $\times 10^{-7}$ CL=90%	870
$K_S^0 \pi^- 2e^+$		< 3.3 $\times 10^{-6}$ CL=90%	863
$K^- \pi^0 2e^+$		< 8.5 $\times 10^{-6}$ CL=90%	864
$K^- 2\mu^+$	<i>L</i>	< 1.0 $\times 10^{-5}$ CL=90%	856
$K^- e^+ \mu^+$	<i>L</i>	< 1.9 $\times 10^{-6}$ CL=90%	866
$K^*(892)^- 2\mu^+$	<i>L</i>	< 8.5 $\times 10^{-4}$ CL=90%	703

D⁰

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1864.83 \pm 0.05$ MeV

$$m_{D^\pm} - m_{D^0} = 4.822 \pm 0.015 \text{ MeV}$$

$$\text{Mean life } \tau = (410.1 \pm 1.5) \times 10^{-15} \text{ s}$$

$$c\tau = 122.9 \text{ }\mu\text{m}$$

Mixing and related parameters

$$|m_{D_1^0} - m_{D_2^0}| = (0.95^{+0.41}_{-0.44}) \times 10^{10} \text{ } \hbar \text{ s}^{-1}$$

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.29^{+0.14}_{-0.18}) \times 10^{-2}$$

$$|q/p| = 0.92^{+0.12}_{-0.09}$$

$$A_\Gamma = (-0.125 \pm 0.526) \times 10^{-3}$$

$$\phi^{K_S^0 \pi\pi} = -0.09^{+0.10}_{-0.13}$$

$$K^+ \pi^- \text{ relative strong phase: } \cos \delta = 0.97 \pm 0.11$$

$$K^- \pi^+ \pi^0 \text{ coherence factor } R_{K\pi\pi^0} = 0.82 \pm 0.06$$

$$K^- \pi^+ \pi^0 \text{ average relative strong phase } \delta^{K\pi\pi^0} = (199 \pm 14)^\circ$$

$$K^- \pi^- 2\pi^+ \text{ coherence factor } R_{K3\pi} = 0.53^{+0.18}_{-0.21}$$

$$K^- \pi^- 2\pi^+ \text{ average relative strong phase } \delta^{K3\pi} = (125^{+22}_{-14})^\circ$$

$$D^0 \rightarrow K^- \pi^- 2\pi^+, R_{K3\pi} (y \cos \delta^{K3\pi} - x \sin \delta^{K3\pi}) = (-3.0 \pm 0.7) \times 10^{-3} \text{ TeV}^{-1}$$

$$K_S^0 K^+ \pi^- \text{ coherence factor } R_{K_S^0 K\pi} = 0.70 \pm 0.08$$

$$K_S^0 K^+ \pi^- \text{ average relative strong phase } \delta^{K_S^0 K\pi} = (0 \pm 16)^\circ$$

$$K^* K \text{ coherence factor } R_{K^* K} = 0.94 \pm 0.12$$

$$K^* K \text{ average relative strong phase } \delta^{K^* K} = (-17 \pm 18)^\circ$$

CP-violation decay-rate asymmetries (labeled by the D^0 decay)

$$A_{CP}(K^+ K^-) = (-0.07 \pm 0.11)\%$$

$$A_{CP}(2K_S^0) = (0.4 \pm 1.4)\%$$

$$A_{CP}(\pi^+ \pi^-) = (0.13 \pm 0.14)\%$$

$$A_{CP}(\pi^0 \pi^0) = (0.0 \pm 0.6)\%$$

$$A_{CP}(\rho \gamma) = (6 \pm 15) \times 10^{-2}$$

$$A_{CP}(\phi \gamma) = (-9 \pm 7) \times 10^{-2}$$

$$A_{CP}(\overline{K}^*(892)^0 \gamma) = (-0.3 \pm 2.0) \times 10^{-2}$$

$$A_{CP}(\pi^+ \pi^- \pi^0) = (0.3 \pm 0.4)\%$$

$$A_{CP}(\rho(770)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (1.2 \pm 0.9)\% [i]$$

$$A_{CP}(\rho(770)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-3.1 \pm 3.0)\% [i]$$

$$A_{CP}(\rho(770)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (-1.0 \pm 1.7)\% [i]$$

$$A_{CP}(\rho(1450)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 70)\% [i]$$

$$A_{CP}(\rho(1450)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-20 \pm 40)\% [i]$$

$$A_{CP}(\rho(1450)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (6 \pm 9)\% [i]$$

$$A_{CP}(\rho(1700)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (-5 \pm 14)\% [i]$$

$$A_{CP}(\rho(1700)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (13 \pm 9)\% [i]$$

$$A_{CP}(\rho(1700)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (8 \pm 11)\% [i]$$

$$A_{CP}(f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 35)\% [i]$$

$$A_{CP}(f_0(1370) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (25 \pm 18)\% [i]$$

$$\begin{aligned}
A_{CP}(f_0(1500)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (0 \pm 18)\% [i] \\
A_{CP}(f_0(1710)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (0 \pm 24)\% [i] \\
A_{CP}(f_2(1270)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (-4 \pm 6)\% [i] \\
A_{CP}(\sigma(400)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (6 \pm 8)\% [i] \\
A_{CP}(\text{nonresonant } \pi^+\pi^-\pi^0) &= (-13 \pm 23)\% [i] \\
A_{CP}(a_1(1260)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (5 \pm 6)\% \\
A_{CP}(a_1(1260)^-\pi^+ \rightarrow 2\pi^+2\pi^-) &= (14 \pm 18)\% \\
A_{CP}(\pi(1300)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (-2 \pm 15)\% \\
A_{CP}(\pi(1300)^-\pi^+ \rightarrow 2\pi^+2\pi^-) &= (-6 \pm 30)\% \\
A_{CP}(a_1(1640)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (9 \pm 26)\% \\
A_{CP}(\pi_2(1670)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (7 \pm 18)\% \\
A_{CP}(\sigma f_0(1370) \rightarrow 2\pi^+2\pi^-) &= (-15 \pm 19)\% \\
A_{CP}(\sigma\rho(770)^0 \rightarrow 2\pi^+2\pi^-) &= (3 \pm 27)\% \\
A_{CP}(2\rho(770)^0 \rightarrow 2\pi^+2\pi^-) &= (-6 \pm 6)\% \\
A_{CP}(2f_2(1270) \rightarrow 2\pi^+2\pi^-) &= (-28 \pm 24)\% \\
A_{CP}(K^+K^-\pi^0) &= (-1.0 \pm 1.7)\% \\
A_{CP}(K^*(892)^+K^- \rightarrow K^+K^-\pi^0) &= (-0.9 \pm 1.3)\% [i] \\
A_{CP}(K^*(1410)^+K^- \rightarrow K^+K^-\pi^0) &= (-21 \pm 24)\% [i] \\
A_{CP}((K^+\pi^0)_{S-\text{wave}}K^- \rightarrow K^+K^-\pi^0) &= (7 \pm 15)\% [i] \\
A_{CP}(\phi(1020)\pi^0 \rightarrow K^+K^-\pi^0) &= (1.1 \pm 2.2)\% [i] \\
A_{CP}(f_0(980)\pi^0 \rightarrow K^+K^-\pi^0) &= (-3 \pm 19)\% [i] \\
A_{CP}(a_0(980)^0\pi^0 \rightarrow K^+K^-\pi^0) &= (-5 \pm 16)\% [i] \\
A_{CP}(f'_2(1525)\pi^0 \rightarrow K^+K^-\pi^0) &= (0 \pm 160)\% [i] \\
A_{CP}(K^*(892)^-K^+ \rightarrow K^+K^-\pi^0) &= (-5 \pm 4)\% [i] \\
A_{CP}(K^*(1410)^-K^+ \rightarrow K^+K^-\pi^0) &= (-17 \pm 29)\% [i] \\
A_{CP}((K^-\pi^0)_{S-\text{wave}}K^+ \rightarrow K^+K^-\pi^0) &= (-10 \pm 40)\% [i] \\
A_{CP}(K_S^0\pi^0) &= (-0.20 \pm 0.17)\% \\
A_{CP}(K_S^0\eta) &= (0.5 \pm 0.5)\% \\
A_{CP}(K_S^0\eta') &= (1.0 \pm 0.7)\% \\
A_{CP}(K_S^0\phi) &= (-3 \pm 9)\% \\
A_{CP}(K^-\pi^+) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+\pi^-) &= (-0.9 \pm 1.4)\% \\
A_{CP}(D_{CP(\pm 1)} \rightarrow K^\mp\pi^\pm) &= (12.7 \pm 1.5)\% \\
A_{CP}(K^-\pi^+\pi^0) &= (0.1 \pm 0.5)\% \\
A_{CP}(K^+\pi^-\pi^0) &= (0 \pm 5)\% \\
A_{CP}(K_S^0\pi^+\pi^-) &= (-0.1 \pm 0.8)\% \\
A_{CP}(K^*(892)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (0.4 \pm 0.5)\% \\
A_{CP}(K^*(892)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (1 \pm 6)\% \\
A_{CP}(\overline{K}^0\rho^0 \rightarrow K_S^0\pi^+\pi^-) &= (-0.1 \pm 0.5)\% \\
A_{CP}(\overline{K}^0\omega \rightarrow K_S^0\pi^+\pi^-) &= (-13 \pm 7)\% \\
A_{CP}(\overline{K}^0f_0(980) \rightarrow K_S^0\pi^+\pi^-) &= (-0.4 \pm 2.7)\% \\
A_{CP}(\overline{K}^0f_2(1270) \rightarrow K_S^0\pi^+\pi^-) &= (-4 \pm 5)\%
\end{aligned}$$

$$\begin{aligned}
A_{CP}(\overline{K^0} f_0(1370) \rightarrow K_S^0 \pi^+ \pi^-) &= (-1 \pm 9)\% \\
A_{CP}(\overline{K^0} \rho^0(1450) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 10)\% \\
A_{CP}(\overline{K^0} f_0(600) \rightarrow K_S^0 \pi^+ \pi^-) &= (-3 \pm 5)\% \\
A_{CP}(K^*(1410)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (-2 \pm 9)\% \\
A_{CP}(K_0^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (4 \pm 4)\% \\
A_{CP}(K_0^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (12 \pm 15)\% \\
A_{CP}(K_2^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (3 \pm 6)\% \\
A_{CP}(K_2^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (-10 \pm 32)\% \\
A_{CP}(K^- \pi^+ \pi^+ \pi^-) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+ \pi^- \pi^+ \pi^-) &= (-2 \pm 4)\% \\
A_{CP}(K^+ K^- \pi^+ \pi^-) &= (1.3 \pm 1.7)\% \\
A_{CP}(K_1^*(1270)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-2.3 \pm 1.7)\% \\
A_{CP}(K_1^*(1270)^+ K^- \rightarrow K^{*0} \pi^+ K^-) &= (-1 \pm 10)\% \\
A_{CP}(K_1^*(1270)^- K^+ \rightarrow \overline{K}^{*0} \pi^- K^+) &= (-10 \pm 32)\% \\
A_{CP}(K_1^*(1270)^- K^+ \rightarrow K^+ K^- \pi^+ \pi^-) &= (1.7 \pm 3.5)\% \\
A_{CP}(K_1^*(1270)^+ K^- \rightarrow \rho^0 K^+ K^-) &= (-7 \pm 17)\% \\
A_{CP}(K_1^*(1270)^- K^+ \rightarrow \rho^0 K^- K^+) &= (10 \pm 13)\% \\
A_{CP}(K_1(1400)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-4.4 \pm 2.1)\% \\
A_{CP}(K^*(1410)^+ K^- \rightarrow K^{*0} \pi^+ K^-) &= (-20 \pm 17)\% \\
A_{CP}(K^*(1410)^- K^+ \rightarrow \overline{K}^{*0} \pi^- K^+) &= (-1 \pm 14)\% \\
A_{CP}(K^*(1680)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-17 \pm 29)\% \\
A_{CP}(K^{*0} \overline{K}^{*0}) \text{ in } D^0, \overline{D}^0 \rightarrow K^{*0} \overline{K}^{*0} &= (-5 \pm 14)\% \\
A_{CP}(K^{*0} \overline{K}^{*0} S\text{-wave}) &= (-3.9 \pm 2.2)\% \\
A_{CP}(\phi \rho^0) \text{ in } D^0, \overline{D}^0 \rightarrow \phi \rho^0 &= (1 \pm 9)\% \\
A_{CP}(\phi \rho^0 S\text{-wave}) &= (-3 \pm 5)\% \\
A_{CP}(\phi \rho^0 D\text{-wave}) &= (-37 \pm 19)\% \\
A_{CP}(\phi (\pi^+ \pi^-)_{S\text{-wave}}) &= (6 \pm 6)\% \\
A_{CP}(K^*(892)^0 (K^- \pi^+)_{S\text{-wave}}) &= (-10 \pm 40)\% \\
A_{CP}(K^+ K^- \pi^+ \pi^- \text{ non-resonant}) &= (8 \pm 20)\% \\
A_{CP}((K^- \pi^+)_{P\text{-wave}} (K^+ \pi^-)_{S\text{-wave}}) &= (3 \pm 11)\% \\
A_{CP}(K^+ K^- \mu^+ \mu^-) \text{ in } D^0, \overline{D}^0 \rightarrow K^+ K^- \mu^+ \mu^- &= (0 \pm 11)\% \\
A_{CP}(\pi^+ \pi^- \mu^+ \mu^-) \text{ in } D^0, \overline{D}^0 \rightarrow \pi^+ \pi^- \mu^+ \mu^- &= (5 \pm 4)\%
\end{aligned}$$

***CP*-even fractions (labeled by the D^0 decay)**

$$\begin{aligned}
&CP\text{-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^0 \text{ decays} = (97.3 \pm 1.7)\% \\
&CP\text{-even fraction in } D^0 \rightarrow K^+ K^- \pi^0 \text{ decays} = (73 \pm 6)\% \\
&CP\text{-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- \text{ decays} = (76.9 \pm 2.3)\% \\
&CP\text{-even fraction in } D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0 \text{ decays} = (23.8 \pm 1.7)\% \\
&CP\text{-even fraction in } D^0 \rightarrow K^+ K^- \pi^+ \pi^- \text{ decays} = (75 \pm 4)\%
\end{aligned}$$

***CP*-violation asymmetry difference**

$$\Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-0.154 \pm 0.029)\%$$

χ^2 tests of CP -violation (CPV) p-values

Local CPV in D^0 , $\bar{D}^0 \rightarrow \pi^+ \pi^- \pi^0 = 4.9\%$

Local CPV in D^0 , $\bar{D}^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- = (0.6 \pm 0.2)\%$

Local CPV in D^0 , $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^- = 96\%$

Local CPV in D^0 , $\bar{D}^0 \rightarrow K^+ K^- \pi^0 = 16.6\%$

Local CPV in D^0 , $\bar{D}^0 \rightarrow K^+ K^- \pi^+ \pi^- = 9.1\%$

T -violation decay-rate asymmetry

$$A_T(K^+ K^- \pi^+ \pi^-) = (2.9 \pm 2.2) \times 10^{-3} [b]$$

$$A_{T\text{viol}}(K_S \pi^+ \pi^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow K_S \pi^+ \pi^- \pi^0 = (-0.3^{+1.4}_{-1.6}) \times 10^{-3}$$

CPT -violation decay-rate asymmetry

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$r_V \equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 1.46 \pm 0.07$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 0.68 \pm 0.06$$

$$f_+(0) \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.736 \pm 0.004$$

$$f_+(0)|V_{cs}| \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.7166 \pm 0.0030$$

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = -2.40 \pm 0.16$$

$$r_2 \equiv a_2/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 5 \pm 4$$

$$f_+(0) \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 0.637 \pm 0.009$$

$$f_+(0)|V_{cd}| \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 0.1436 \pm 0.0026 \quad (S = 1.5)$$

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = -1.97 \pm 0.28 \quad (S = 1.4)$$

$$r_2 \equiv a_2/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = -0.2 \pm 2.2 \quad (S = 1.7)$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ p Confidence leve(MeV/c)
Topological modes		
0-prongs	[j] (15 ± 6) %	—
2-prongs	(71 ± 6) %	—
4-prongs	[k] (14.6 ± 0.5) %	—
6-prongs	[l] (6.5 ± 1.3) × 10 ⁻⁴	—

Inclusive modes

e^+ anything	[η]	(6.49 \pm 0.11) %		
μ^+ anything		(6.8 \pm 0.6) %		
K^- anything		(54.7 \pm 2.8) %	S=1.3	
\bar{K}^0 anything + K^0 anything		(47 \pm 4) %		
K^+ anything		(3.4 \pm 0.4) %		
$K^*(892)^-$ anything		(15 \pm 9) %		
$\bar{K}^*(892)^0$ anything		(9 \pm 4) %		
$K^*(892)^+$ anything		< 3.6 %	CL=90%	
$K^*(892)^0$ anything		(2.8 \pm 1.3) %		
η anything		(9.5 \pm 0.9) %		
η' anything		(2.48 \pm 0.27) %		
ϕ anything		(1.08 \pm 0.04) %		
invisibles		< 9.4 $\times 10^{-5}$	CL=90%	

Semileptonic modes

$K^- e^+ \nu_e$		(3.542 \pm 0.035) %	S=1.3	867
$K^- \mu^+ \nu_\mu$		(3.41 \pm 0.04) %		864
$K^*(892)^- e^+ \nu_e$		(2.15 \pm 0.16) %		719
$K^*(892)^- \mu^+ \nu_\mu$		(1.89 \pm 0.24) %		714
$K^- \pi^0 e^+ \nu_e$		(1.6 \pm 1.3) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$		(1.44 \pm 0.04) %		860
$(\bar{K}^0 \pi^-)_{S-wave} e^+ \nu_e$		(7.9 \pm 1.7) $\times 10^{-4}$		860
$K^- \pi^+ \pi^- e^+ \nu_e$		(2.8 \pm 1.4) $\times 10^{-4}$		843
$K_1(1270)^- e^+ \nu_e$		(7.6 \pm 4.0) $\times 10^{-4}$		511
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$		< 1.3 $\times 10^{-3}$	CL=90%	821
$(\bar{K}^*(892) \pi)^- \mu^+ \nu_\mu$		< 1.5 $\times 10^{-3}$	CL=90%	692
$\pi^- e^+ \nu_e$		(2.91 \pm 0.04) $\times 10^{-3}$		927
$\pi^- \mu^+ \nu_\mu$		(2.67 \pm 0.12) $\times 10^{-3}$	S=1.3	924
$\pi^- \pi^0 e^+ \nu_e$		(1.45 \pm 0.07) $\times 10^{-3}$		922
$\rho^- e^+ \nu_e$		(1.50 \pm 0.12) $\times 10^{-3}$	S=1.9	771
$a(980)^- e^+ \nu_e, \quad a^- \rightarrow \eta \pi^-$		(1.33 \pm 0.34) $\times 10^{-4}$		—

Hadronic modes with one \bar{K}

$K^- \pi^+$		(3.950 \pm 0.031) %	S=1.2	861
$K_S^0 \pi^0$		(1.240 \pm 0.022) %		860
$K_L^0 \pi^0$		(10.0 \pm 0.7) $\times 10^{-3}$		860
$K_S^0 \pi^+ \pi^-$	[c]	(2.80 \pm 0.18) %	S=1.1	842
$K_S^0 \rho^0$		(6.3 \pm 0.6) $\times 10^{-3}$		674
$K_S^0 \omega, \quad \omega \rightarrow \pi^+ \pi^-$		(2.0 \pm 0.6) $\times 10^{-4}$		670
$K_S^0 (\pi^+ \pi^-)_{S-wave}$		(3.3 \pm 0.8) $\times 10^{-3}$		842

$K_S^0 f_0(980)$, $f_0 \rightarrow \pi^+ \pi^-$	$(1.20 \pm 0.40) \times 10^{-3}$	549
$K_S^0 f_0(1370)$, $f_0 \rightarrow \pi^+ \pi^-$	$(2.8 \pm 0.9) \times 10^{-3}$	†
$K_S^0 f_2(1270)$, $f_2 \rightarrow \pi^+ \pi^-$	$(9 \pm 10) \times 10^{-5}$	262
$K^*(892)^- \pi^+$, $K^{*-} \rightarrow K_S^0 \pi^-$	$(1.64 \pm 0.14) \%$	711
$K_0^*(1430)^- \pi^+$, $K_0^{*-} \rightarrow K_S^0 \pi^-$	$(2.67 \pm 0.40) \times 10^{-3}$	378
$K_2^*(1430)^- \pi^+$, $K_2^{*-} \rightarrow K_S^0 \pi^-$	$(3.4 \pm 1.9) \times 10^{-4}$	367
$K^*(1680)^- \pi^+$, $K^{*-} \rightarrow K_S^0 \pi^-$	$(4.4 \pm 3.5) \times 10^{-4}$	46
$K^*(892)^+ \pi^-$, $K^{*+} \rightarrow K_S^0 \pi^+$	[o] $(1.13 \pm 0.60) \times 10^{-4}$	711
$K_0^*(1430)^+ \pi^-$, $K_0^{*+} \rightarrow K_S^0 \pi^+$	[o] $< 1.4 \times 10^{-5}$ CL=95% —	
$K_2^*(1430)^+ \pi^-$, $K_2^{*+} \rightarrow K_S^0 \pi^+$	[o] $< 3.4 \times 10^{-5}$ CL=95% —	
$K_S^0 \pi^+ \pi^-$ nonresonant	$(2.5 \pm 6.0) \times 10^{-4}$	842
$K^- \pi^+ \pi^0$	[c] $(14.4 \pm 0.5) \%$ S=2.0 844	
$K^- \rho^+$	$(11.3 \pm 0.7) \%$	675
$K^- \rho(1700)^+$, $\rho^+ \rightarrow \pi^+ \pi^0$	$(8.2 \pm 1.8) \times 10^{-3}$	†
$K^*(892)^- \pi^+$, $K^*(892)^- \rightarrow K^- \pi^0$	$(2.31 \pm 0.40) \%$	711
$\bar{K}^*(892)^0 \pi^0$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.95 \pm 0.24) \%$	711
$K_0^*(1430)^- \pi^+$, $K_0^{*-} \rightarrow K^- \pi^0$	$(4.8 \pm 2.2) \times 10^{-3}$	378
$\bar{K}_0^*(1430)^0 \pi^0$, $\bar{K}_0^{*0} \rightarrow K^- \pi^+$	$(5.9 \pm 5.0) \times 10^{-3}$	379
$K^*(1680)^- \pi^+$, $K^{*-} \rightarrow K^- \pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$	46
$K^- \pi^+ \pi^0$ nonresonant	$(1.15 \pm 0.60) \%$	844
$K_S^0 2\pi^0$	$(9.1 \pm 1.1) \times 10^{-3}$	S=2.2 843
$K_S^0 (2\pi^0)_{S-wave}$	$(2.6 \pm 0.7) \times 10^{-3}$	—
$\bar{K}^*(892)^0 \pi^0$, $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(8.1 \pm 0.7) \times 10^{-3}$	711
$\bar{K}^*(1430)^0 \pi^0$, $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(4 \pm 23) \times 10^{-5}$	—
$\bar{K}^*(1680)^0 \pi^0$, $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(1.0 \pm 0.4) \times 10^{-3}$	—

$K_S^0 f_2(1270)$, $f_2 \rightarrow 2\pi^0$	(2.3 \pm 1.1) $\times 10^{-4}$	-	-
$2K_S^0$, one $K_S^0 \rightarrow 2\pi^0$	(3.2 \pm 1.1) $\times 10^{-4}$	-	-
$K^- 2\pi^+ \pi^-$	[c] (8.23 \pm 0.14) %	S=1.1	813
$K^- \pi^+ \rho^0$ total	(6.87 \pm 0.31) %		609
$K^- \pi^+ \rho^0$ 3-body	(6.1 \pm 1.6) $\times 10^{-3}$		609
$\bar{K}^*(892)^0 \rho^0$, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(1.01 \pm 0.05) %		416
$\bar{K}^*(892)^0 \rho^0$ transverse,	(1.2 \pm 0.4) %		417
$\bar{K}^{*0} \rightarrow K^- \pi^+$			
$K^- a_1(1260)^+$, $a_1^+ \rightarrow \rho^0 \pi^+$	(4.33 \pm 0.32) %		327
$K_1(1270)^- \pi^+$, $K_1^- \rightarrow K^- \pi^+ \pi^-$ total	(3.9 \pm 0.4) $\times 10^{-3}$		-
$K_1(1270)^- \pi^+$, $K_1^- \rightarrow \bar{K}^*(892)^0 \pi^-$, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(6.6 \pm 2.3) $\times 10^{-4}$		484
$K^- 2\pi^+ \pi^-$ nonresonant	(1.81 \pm 0.07) %		813
$K_S^0 \pi^+ \pi^- \pi^0$	[p] (5.2 \pm 0.6) %		813
$K_S^0 \eta$, $\eta \rightarrow \pi^+ \pi^- \pi^0$	(1.17 \pm 0.03) $\times 10^{-3}$		772
$K_S^0 \omega$, $\omega \rightarrow \pi^+ \pi^- \pi^0$	(9.9 \pm 0.6) $\times 10^{-3}$		670
$K^- \pi^+ 2\pi^0$	(8.86 \pm 0.23) %		815
$K^- 2\pi^+ \pi^- \pi^0$	(4.3 \pm 0.4) %		771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(1.3 \pm 0.6) %		643
$K^- \pi^+ \omega$, $\omega \rightarrow \pi^+ \pi^- \pi^0$	(2.8 \pm 0.5) %		605
$\bar{K}^*(892)^0 \omega$, $\bar{K}^{*0} \rightarrow K^- \pi^+$, $\omega \rightarrow \pi^+ \pi^- \pi^0$	(6.5 \pm 3.0) $\times 10^{-3}$		410
$K_S^0 \eta \pi^0$	(5.7 \pm 1.1) $\times 10^{-3}$		721
$K_S^0 a_0(980)$, $a_0 \rightarrow \eta \pi^0$	(6.8 \pm 2.1) $\times 10^{-3}$		-
$\bar{K}^*(892)^0 \eta$, $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$	(1.7 \pm 0.5) $\times 10^{-3}$		-
$K_S^0 2\pi^+ 2\pi^-$	(2.66 \pm 0.30) $\times 10^{-3}$		768
$K_S^0 \rho^0 \pi^+ \pi^-$, no $K^*(892)^-$	(1.1 \pm 0.7) $\times 10^{-3}$		-
$K^*(892)^- 2\pi^+ \pi^-$, $K^*(892)^- \rightarrow K_S^0 \pi^-$, no ρ^0	(5 \pm 7) $\times 10^{-4}$		642
$K^*(892)^- \rho^0 \pi^+$, $K^*(892)^- \rightarrow K_S^0 \pi^-$	(1.6 \pm 0.6) $\times 10^{-3}$		230
$K_S^0 2\pi^+ 2\pi^-$ nonresonant	< 1.2 $\times 10^{-3}$	CL=90%	768
$K^- 3\pi^+ 2\pi^-$	(2.2 \pm 0.6) $\times 10^{-4}$		713

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. These nine modes below are all corrected for unseen decays of the resonances.

$K_S^0 \eta$	$(5.09 \pm 0.13) \times 10^{-3}$	772
$K_S^0 \omega$	$(1.11 \pm 0.06) \%$	670
$K_S^0 \eta'(958)$	$(9.49 \pm 0.32) \times 10^{-3}$	565
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	$(1.9 \pm 0.9) \%$	643
$K^- \pi^+ \omega$	$(3.1 \pm 0.6) \%$	605
$\bar{K}^*(892)^0 \omega$	$(1.1 \pm 0.5) \%$	410
$K^- \pi^+ \eta'(958)$	$(6.43 \pm 0.34) \times 10^{-3}$	479
$K_S^0 \eta'(958) \pi^0$	$(2.52 \pm 0.27) \times 10^{-3}$	479
$\bar{K}^*(892)^0 \eta'(958)$	$< 1.0 \times 10^{-3}$	CL=90% 119

Hadronic modes with three K 's

$K_S^0 K^+ K^-$	$(4.42 \pm 0.32) \times 10^{-3}$	544
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	$(2.9 \pm 0.4) \times 10^{-3}$	—
$K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0$	$(5.9 \pm 1.8) \times 10^{-4}$	—
$K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0$	$< 1.1 \times 10^{-4}$	CL=95% —
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	$< 9 \times 10^{-5}$	CL=95% —
$K_S^0 \phi, \phi \rightarrow K^+ K^-$	$(2.03 \pm 0.15) \times 10^{-3}$	520
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	$(1.7 \pm 1.1) \times 10^{-4}$	—
$3K_S^0$	$(7.5 \pm 0.7) \times 10^{-4}$	S=1.4 539
$K^+ 2K^- \pi^+$	$(2.25 \pm 0.32) \times 10^{-4}$	434
$K^+ K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(4.5 \pm 1.8) \times 10^{-5}$	†
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	$(4.1 \pm 1.7) \times 10^{-5}$	422
$\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(1.08 \pm 0.21) \times 10^{-4}$	†
$K^+ 2K^- \pi^+ \text{nonresonant}$	$(3.4 \pm 1.5) \times 10^{-5}$	434
$2K_S^0 K^\pm \pi^\mp$	$(5.9 \pm 1.3) \times 10^{-4}$	427

Pionic modes

$\pi^+ \pi^-$	$(1.455 \pm 0.024) \times 10^{-3}$	S=1.3 922
$2\pi^0$	$(8.26 \pm 0.25) \times 10^{-4}$	923
$\pi^+ \pi^- \pi^0$	$(1.49 \pm 0.06) \%$	S=2.1 907
$\rho^+ \pi^-$	$(1.01 \pm 0.04) \%$	764
$\rho^0 \pi^0$	$(3.86 \pm 0.23) \times 10^{-3}$	764
$\rho^- \pi^+$	$(5.15 \pm 0.25) \times 10^{-3}$	764
$\rho(1450)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.6 \pm 2.1) \times 10^{-5}$	—
$\rho(1450)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	$(4.5 \pm 2.0) \times 10^{-5}$	—
$\rho(1450)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	$(2.7 \pm 0.4) \times 10^{-4}$	—
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	$(6.1 \pm 1.5) \times 10^{-4}$	—

$\rho(1700)^0 \pi^0$, $\rho^0 \rightarrow \pi^+ \pi^-$	(7.4 \pm 1.8) $\times 10^{-4}$	-
$\rho(1700)^- \pi^+$, $\rho^- \rightarrow \pi^- \pi^0$	(4.8 \pm 1.1) $\times 10^{-4}$	-
$f_0(980) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	(3.7 \pm 0.9) $\times 10^{-5}$	-
$f_0(500) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	(1.22 \pm 0.22) $\times 10^{-4}$	-
$f_0(1370) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	(5.5 \pm 2.1) $\times 10^{-5}$	-
$f_0(1500) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	(5.8 \pm 1.6) $\times 10^{-5}$	-
$f_0(1710) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	(4.6 \pm 1.6) $\times 10^{-5}$	-
$f_2(1270) \pi^0$, $f_2 \rightarrow \pi^+ \pi^-$	(1.97 \pm 0.21) $\times 10^{-4}$	-
$\pi^+ \pi^- \pi^0$ nonresonant	(1.3 \pm 0.4) $\times 10^{-4}$	907
$3\pi^0$	(2.0 \pm 0.5) $\times 10^{-4}$	908
$2\pi^+ 2\pi^-$	(7.56 \pm 0.20) $\times 10^{-3}$	880
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow 2\pi^+ \pi^-$ total	(4.54 \pm 0.31) $\times 10^{-3}$	-
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+$ S-wave	(3.14 \pm 0.21) $\times 10^{-3}$	-
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+$ D-wave	(1.9 \pm 0.5) $\times 10^{-4}$	-
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	(6.4 \pm 0.7) $\times 10^{-4}$	-
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \rho^0 \pi^-$ S-wave	(2.3 \pm 0.9) $\times 10^{-4}$	-
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \sigma \pi^-$	(6.1 \pm 3.4) $\times 10^{-5}$	-
$\pi(1300)^+ \pi^-$, $\pi(1300)^+ \rightarrow \pi(1300)^- \pi^+$	(5.1 \pm 2.7) $\times 10^{-4}$	-
$\pi(1300)^- \pi^+$, $\pi(1300)^- \rightarrow \sigma \pi^-$	(2.3 \pm 2.2) $\times 10^{-4}$	-
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+$ D-wave	(3.2 \pm 1.6) $\times 10^{-4}$	-
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	(1.8 \pm 1.4) $\times 10^{-4}$	-
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow f_2(1270)^0 \pi^+$	(2.0 \pm 0.9) $\times 10^{-4}$	-
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow \pi^+ \pi^-$	(2.6 \pm 1.0) $\times 10^{-4}$	-
$2\rho^0$ total	(1.85 \pm 0.13) $\times 10^{-3}$	518
$2\rho^0$, parallel helicities	(8.3 \pm 3.2) $\times 10^{-5}$	-
$2\rho^0$, perpendicular helicities	(4.8 \pm 0.6) $\times 10^{-4}$	-
$2\rho^0$, longitudinal helicities	(1.27 \pm 0.10) $\times 10^{-3}$	-
$2\rho(770)^0$, S-wave	(1.8 \pm 1.3) $\times 10^{-4}$	-
$2\rho(770)^0$, P-wave	(5.3 \pm 1.3) $\times 10^{-4}$	-
$2\rho(770)^0$, D-wave	(6.2 \pm 3.0) $\times 10^{-4}$	-
Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$	(1.51 \pm 0.12) $\times 10^{-3}$	-
3-body total	(6.2 \pm 0.9) $\times 10^{-4}$	-
$\sigma \pi^+ \pi^-$	(6.2 \pm 0.9) $\times 10^{-4}$	-

$\sigma \rho(770)^0$	(5.0 \pm 2.5) $\times 10^{-4}$	-
$f_0(980)\pi^+\pi^-$, $f_0 \rightarrow \pi^+\pi^-$	(1.8 \pm 0.5) $\times 10^{-4}$	-
$f_2(1270)\pi^+\pi^-$, $f_2 \rightarrow \pi^+\pi^-$	(3.7 \pm 0.6) $\times 10^{-4}$	-
$2f_2(1270)$, $f_2 \rightarrow \pi^+\pi^-$	(1.6 \pm 1.8) $\times 10^{-4}$	-
$f_0(1370)\sigma$, $f_0 \rightarrow \pi^+\pi^-$	(1.6 \pm 0.5) $\times 10^{-3}$	-
$\pi^+\pi^-2\pi^0$	(1.02 \pm 0.09) %	882
$\eta\pi^0$	[q] (6.3 \pm 0.6) $\times 10^{-4}$	S=1.1 846
$\omega\pi^0$	[q] (1.17 \pm 0.35) $\times 10^{-4}$	761
$\omega\eta$	(1.98 \pm 0.18) $\times 10^{-3}$	S=1.1 648
$2\pi^+2\pi^-\pi^0$	(4.2 \pm 0.5) $\times 10^{-3}$	844
$\eta\pi^+\pi^-$	[q] (1.09 \pm 0.16) $\times 10^{-3}$	827
$\omega\pi^+\pi^-$	[q] (1.6 \pm 0.5) $\times 10^{-3}$	738
$\eta 2\pi^0$	(3.8 \pm 1.3) $\times 10^{-4}$	829
$3\pi^+3\pi^-$	(4.3 \pm 1.2) $\times 10^{-4}$	795
$\eta'(958)\pi^0$	(9.2 \pm 1.0) $\times 10^{-4}$	678
$\eta'(958)\pi^+\pi^-$	(4.5 \pm 1.7) $\times 10^{-4}$	650
2η	(2.11 \pm 0.19) $\times 10^{-3}$	S=2.2 754
$2\eta\pi^0$	(7.3 \pm 2.2) $\times 10^{-4}$	699
3η	< 1.3 $\times 10^{-4}$	CL=90% 421
$\eta\eta'(958)$	(1.01 \pm 0.19) $\times 10^{-3}$	537

Hadronic modes with a $K\bar{K}$ pair

K^+K^-	(4.08 \pm 0.06) $\times 10^{-3}$	S=1.6 791
$2K_S^0$	(1.41 \pm 0.05) $\times 10^{-4}$	S=1.1 789
$K_S^0 K^- \pi^+$	(3.3 \pm 0.5) $\times 10^{-3}$	S=1.1 739
$K^*(892)^0 K_S^0$, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(8.2 \pm 1.6) $\times 10^{-5}$	608
$K^*(892)^+ K^-$, $K^{*+} \rightarrow K_S^0 \pi^+$	(1.89 \pm 0.30) $\times 10^{-3}$	-
$\bar{K}^*(1410)^0 K_S^0$, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(1.3 \pm 1.9) $\times 10^{-4}$	-
$K^*(1410)^+ K^-$, $K^{*+} \rightarrow K_S^0 \pi^+$	(3.2 \pm 1.9) $\times 10^{-4}$	-
$(K^-\pi^+)_{S-wave} K_S^0$	(6.0 \pm 2.9) $\times 10^{-4}$	739
$(K_S^0\pi^+)_{S-wave} K^-$	(3.9 \pm 1.0) $\times 10^{-4}$	739
$a_0(980)^-\pi^+$, $a_0^- \rightarrow K_S^0 K^-$	(1.3 \pm 1.4) $\times 10^{-4}$	-
$a_0(1450)^-\pi^+$, $a_0^- \rightarrow K_S^0 K^-$	(2.5 \pm 2.0) $\times 10^{-5}$	-
$a_2(1320)^-\pi^+$, $a_2^- \rightarrow K_S^0 K^-$	(5 \pm 5) $\times 10^{-6}$	-
$\rho(1450)^-\pi^+$, $\rho^- \rightarrow K_S^0 K^-$	(4.6 \pm 2.5) $\times 10^{-5}$	-

$K_S^0 K^+ \pi^-$	$(2.17 \pm 0.34) \times 10^{-3}$	S=1.1	739
$K^*(892)^0 K_S^0, \ K^{*0} \rightarrow K^+ \pi^-$	$(1.12 \pm 0.21) \times 10^{-4}$		608
$K^*(892)^- K^+, \ K^{*-} \rightarrow K_S^0 \pi^-$	$(6.2 \pm 1.0) \times 10^{-4}$		-
$K^*(1410)^0 K_S^0, \ K^{*0} \rightarrow K^+ \pi^+$	$(5 \pm 8) \times 10^{-5}$		-
$K^*(1410)^- K^+, \ K^{*-} \rightarrow K_S^0 \pi^-$	$(2.6 \pm 2.0) \times 10^{-4}$		-
$(K^+ \pi^-)_{S-wave} K_S^0$	$(3.7 \pm 1.9) \times 10^{-4}$		739
$(K_S^0 \pi^-)_{S-wave} K^+$	$(1.4 \pm 0.6) \times 10^{-4}$		739
$a_0(980)^+ \pi^-, \ a_0^+ \rightarrow K_S^0 K^+$	$(6 \pm 4) \times 10^{-4}$		-
$a_0(1450)^+ \pi^-, \ a_0^+ \rightarrow K_S^0 K^+$	$(3.2 \pm 2.5) \times 10^{-5}$		-
$\rho(1700)^+ \pi^-, \ \rho^+ \rightarrow K_S^0 K^+$	$(1.1 \pm 0.6) \times 10^{-5}$		-
$K^+ K^- \pi^0$	$(3.42 \pm 0.14) \times 10^{-3}$		743
$K^*(892)^+ K^-, \ K^*(892)^+ \rightarrow K^+ \pi^0$	$(1.52 \pm 0.07) \times 10^{-3}$		-
$K^*(892)^- K^+, \ K^*(892)^- \rightarrow K^- \pi^0$	$(5.4 \pm 0.4) \times 10^{-4}$		-
$(K^+ \pi^0)_{S-wave} K^-$	$(2.43 \pm 0.18) \times 10^{-3}$		743
$(K^- \pi^0)_{S-wave} K^+$	$(1.3 \pm 0.5) \times 10^{-4}$		743
$f_0(980) \pi^0, \ f_0 \rightarrow K^+ K^-$	$(3.6 \pm 0.6) \times 10^{-4}$		-
$\phi \pi^0, \ \phi \rightarrow K^+ K^-$	$(6.6 \pm 0.4) \times 10^{-4}$		-
$2K_S^0 \pi^0$	$< 5.9 \times 10^{-4}$		740
$K^+ K^- \pi^+ \pi^-$	$(2.47 \pm 0.11) \times 10^{-3}$		677
$\phi(\pi^+ \pi^-)_{S-wave}, \ \phi \rightarrow K^+ K^-$	$(10 \pm 5) \times 10^{-5}$		614
$(\phi \rho^0)_{S-wave}, \ \phi \rightarrow K^+ K^-$	$(6.9 \pm 0.6) \times 10^{-4}$		250
$(\phi \rho^0)_{P-wave}, \ \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.9) \times 10^{-5}$		-
$(\phi \rho^0)_{D-wave}, \ \phi \rightarrow K^+ K^-$	$(4.2 \pm 1.4) \times 10^{-5}$		-
$(K^*(892)^0 \bar{K}^*(892)^0)_{S-wave}, \ K^{*0} \rightarrow K^\pm \pi^\mp$	$(2.24 \pm 0.13) \times 10^{-4}$		-
$(K^*(892)^0 \bar{K}^*(892)^0)_{P-wave}, \ K^* \rightarrow K^\pm \pi^\mp$	$(1.20 \pm 0.08) \times 10^{-4}$		-
$(K^*(892)^0 \bar{K}^*(892)^0)_{D-wave}, \ K^* \rightarrow K^\pm \pi^\mp$	$(4.7 \pm 0.4) \times 10^{-5}$		-
$K^*(892)^0 (K^- \pi^+)_{S-wave}$	$(1.4 \pm 0.6) \times 10^{-4}$		-
3-body, $K^{*0} \rightarrow K^+ \pi^-$			-
$K_1(1270)^+ K^-, \ K_1^+ \rightarrow K^{*0} \pi^+$	$(1.4 \pm 0.9) \times 10^{-4}$		-
$K_1(1270)^+ K^-, \ K_1^+ \rightarrow K^*(1430)^0 \pi^+$	$(1.5 \pm 0.5) \times 10^{-4}$		-
$K^+ \pi^-$			-

$K_1(1270)^+ K^-$, $K_1^+ \rightarrow \rho^0 K^+$	$(2.2 \pm 0.6) \times 10^{-4}$	-
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow \omega(782) K^+$, $\omega \rightarrow \pi^+ \pi^-$	$(1.5 \pm 1.2) \times 10^{-5}$	-
$K_1(1270)^- K^+$, $K_1^- \rightarrow \rho^0 K^-$	$(1.3 \pm 0.4) \times 10^{-4}$	-
$K_1(1400)^+ K^-$, $K_1^+ \rightarrow K^*(892)^0 \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(4.6 \pm 0.4) \times 10^{-4}$	-
$K^*(1410)^- K^+$, $K^{*-} \rightarrow \bar{K}^{*0} \pi^-$	$(7.0 \pm 1.1) \times 10^{-5}$	-
$K_1(1680)^+ K^-$, $K_1^+ \rightarrow K^{*0} \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(8.9 \pm 3.2) \times 10^{-5}$	-
$K^+ K^- \pi^+ \pi^-$ non-resonant	$(2.7 \pm 0.6) \times 10^{-4}$	-
$2K_S^0 \pi^+ \pi^-$	$(1.22 \pm 0.23) \times 10^{-3}$	673
$K_S^0 K^- 2\pi^+ \pi^-$	$< 1.4 \times 10^{-4}$	CL=90% 595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$	600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi \pi^0$	$(1.17 \pm 0.04) \times 10^{-3}$	645
$\phi \eta$	$(1.8 \pm 0.5) \times 10^{-4}$	489
$\phi \omega$	$< 2.1 \times 10^{-3}$	CL=90% 238

Radiative modes

$\rho^0 \gamma$	$(1.82 \pm 0.32) \times 10^{-5}$	771
$\omega \gamma$	$< 2.4 \times 10^{-4}$	CL=90% 768
$\phi \gamma$	$(2.81 \pm 0.19) \times 10^{-5}$	654
$\bar{K}^*(892)^0 \gamma$	$(4.2 \pm 0.7) \times 10^{-4}$	719

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

$K^+ \ell^- \bar{\nu}_\ell$ via \bar{D}^0	$< 2.2 \times 10^{-5}$	CL=90%	-
K^+ or $K^*(892)^+ e^- \bar{\nu}_e$ via \bar{D}^0	$< 6 \times 10^{-5}$	CL=90%	-
$K^+ \pi^-$ via DCS	$(1.50 \pm 0.07) \times 10^{-4}$	S=3.0	861
$K^+ \pi^-$ via \bar{D}^0	$(1.364 \pm 0.026) \times 10^{-4}$	-	-
$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \bar{D}^0$	$< 1.6 \times 10^{-5}$	CL=95%	861
$K^*(892)^+ \pi^-$, $K^{*+} \rightarrow K_S^0 \pi^+$	$< 1.8 \times 10^{-4}$	CL=95%	-
$K_0^*(1430)^+ \pi^-$, $K_0^{*+} \rightarrow K_S^0 \pi^+$	$(1.13 \pm 0.60) \times 10^{-4}$	711	-
$K_2^*(1430)^+ \pi^-$, $K_2^{*+} \rightarrow K_S^0 \pi^+$	$< 3.4 \times 10^{-5}$	-	-

$K^+ \pi^- \pi^0$	DC	$(3.06 \pm 0.15) \times 10^{-4}$	844
$K^+ \pi^- \pi^0$ via \bar{D}^0		$(7.6 \pm 0.5) \times 10^{-4}$	-
$K^+ \pi^+ 2\pi^-$ via DCS		$(2.49 \pm 0.07) \times 10^{-4}$	-
$K^+ \pi^+ 2\pi^-$	DC	$(2.65 \pm 0.06) \times 10^{-4}$	813
$K^+ \pi^+ 2\pi^-$ via \bar{D}^0		$(7.9 \pm 3.0) \times 10^{-6}$	812
μ^- anything via \bar{D}^0		$< 4 \times 10^{-4}$	CL=90%

**$\Delta C = 1$ weak neutral current ($C1$) modes,
Lepton Family number (LF) violating modes,
Lepton (L) or Baryon (B) number violating modes**

$\gamma\gamma$	$C1$	$< 8.5 \times 10^{-7}$	CL=90%	932
$e^+ e^-$	$C1$	$< 7.9 \times 10^{-8}$	CL=90%	932
$\mu^+ \mu^-$	$C1$	$< 6.2 \times 10^{-9}$	CL=90%	926
$\pi^0 e^+ e^-$	$C1$	$< 4 \times 10^{-6}$	CL=90%	928
$\pi^0 \mu^+ \mu^-$	$C1$	$< 1.8 \times 10^{-4}$	CL=90%	915
$\eta e^+ e^-$	$C1$	$< 3 \times 10^{-6}$	CL=90%	852
$\eta \mu^+ \mu^-$	$C1$	$< 5.3 \times 10^{-4}$	CL=90%	838
$\pi^+ \pi^- e^+ e^-$	$C1$	$< 7 \times 10^{-6}$	CL=90%	922
$\rho^0 e^+ e^-$	$C1$	$< 1.0 \times 10^{-4}$	CL=90%	771
$\pi^+ \pi^- \mu^+ \mu^-$	$C1$	$(9.6 \pm 1.2) \times 10^{-7}$		894
$\pi^+ \pi^- \mu^+ \mu^-$ (non-res)		$< 5.5 \times 10^{-7}$	CL=90%	-
$\rho^0 \mu^+ \mu^-$	$C1$	$< 2.2 \times 10^{-5}$	CL=90%	754
$\omega e^+ e^-$	$C1$	$< 6 \times 10^{-6}$	CL=90%	768
$\omega \mu^+ \mu^-$	$C1$	$< 8.3 \times 10^{-4}$	CL=90%	751
$K^- K^+ e^+ e^-$	$C1$	$< 1.1 \times 10^{-5}$	CL=90%	791
$\phi e^+ e^-$	$C1$	$< 5.2 \times 10^{-5}$	CL=90%	654
$K^- K^+ \mu^+ \mu^-$	$C1$	$(1.54 \pm 0.32) \times 10^{-7}$		710
$K^- K^+ \mu^+ \mu^-$ (non-res)		$< 3.3 \times 10^{-5}$	CL=90%	-
$\phi \mu^+ \mu^-$	$C1$	$< 3.1 \times 10^{-5}$	CL=90%	631
$\bar{K}^0 e^+ e^-$	[h]	$< 2.4 \times 10^{-5}$	CL=90%	866
$\bar{K}^0 \mu^+ \mu^-$	[h]	$< 2.6 \times 10^{-4}$	CL=90%	852
$K^- \pi^+ e^+ e^-$, $675 < m_{ee} < 875$ MeV		$(4.0 \pm 0.5) \times 10^{-6}$		-
$K^- \pi^+ e^+ e^-$, $1.005 < m_{ee} < 1.035$ GeV		$< 5 \times 10^{-7}$	CL=90%	-
$\bar{K}^*(892)^0 e^+ e^-$	[h]	$< 4.7 \times 10^{-5}$	CL=90%	719
$K^- \pi^+ \mu^+ \mu^-$	$C1$	$< 3.59 \times 10^{-4}$	CL=90%	829
$K^- \pi^+ \mu^+ \mu^-$, $675 < m_{\mu\mu} < 875$ MeV		$(4.2 \pm 0.4) \times 10^{-6}$		-
$\bar{K}^*(892)^0 \mu^+ \mu^-$	[h]	$< 2.4 \times 10^{-5}$	CL=90%	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	$C1$	$< 8.1 \times 10^{-4}$	CL=90%	863
$\mu^\pm e^\mp$	LF	$[r] < 1.3 \times 10^{-8}$	CL=90%	929
$\pi^0 e^\pm \mu^\mp$	LF	$[r] < 8.6 \times 10^{-5}$	CL=90%	924
$\eta e^\pm \mu^\mp$	LF	$[r] < 1.0 \times 10^{-4}$	CL=90%	848

$\pi^+ \pi^- e^\pm \mu^\mp$	<i>LF</i>	$[r] < 1.5$	$\times 10^{-5}$	CL=90%	911
$\rho^0 e^\pm \mu^\mp$	<i>LF</i>	$[r] < 4.9$	$\times 10^{-5}$	CL=90%	767
$\omega e^\pm \mu^\mp$	<i>LF</i>	$[r] < 1.2$	$\times 10^{-4}$	CL=90%	764
$K^- K^+ e^\pm \mu^\mp$	<i>LF</i>	$[r] < 1.8$	$\times 10^{-4}$	CL=90%	754
$\phi e^\pm \mu^\mp$	<i>LF</i>	$[r] < 3.4$	$\times 10^{-5}$	CL=90%	648
$\bar{K}^0 e^\pm \mu^\mp$	<i>LF</i>	$[r] < 1.0$	$\times 10^{-4}$	CL=90%	863
$K^- \pi^+ e^\pm \mu^\mp$	<i>LF</i>	$[r] < 5.53$	$\times 10^{-4}$	CL=90%	848
$\bar{K}^*(892)^0 e^\pm \mu^\mp$	<i>LF</i>	$[r] < 8.3$	$\times 10^{-5}$	CL=90%	714
$2\pi^- 2e^+ + \text{c.c.}$	<i>L</i>	< 1.12	$\times 10^{-4}$	CL=90%	922
$2\pi^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 2.9	$\times 10^{-5}$	CL=90%	894
$K^- \pi^- 2e^+$		< 2.8	$\times 10^{-6}$	CL=90%	861
$K^- \pi^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 3.9	$\times 10^{-4}$	CL=90%	829
$2K^- 2e^+ + \text{c.c.}$	<i>L</i>	< 1.52	$\times 10^{-4}$	CL=90%	791
$2K^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 9.4	$\times 10^{-5}$	CL=90%	710
$\pi^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 7.9	$\times 10^{-5}$	CL=90%	911
$K^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 2.18	$\times 10^{-4}$	CL=90%	848
$2K^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 5.7	$\times 10^{-5}$	CL=90%	754
$p e^-$	<i>L,B</i>	$[s] < 1.0$	$\times 10^{-5}$	CL=90%	696
$\bar{p} e^+$	<i>L,B</i>	$[t] < 1.1$	$\times 10^{-5}$	CL=90%	696

 $D^*(2007)^0$

$I(J^P) = \frac{1}{2}(1^-)$

*I, J, P need confirmation.*Mass $m = 2006.85 \pm 0.05$ MeV ($S = 1.1$) $m_{D^{*0}} - m_{D^0} = 142.014 \pm 0.030$ MeV ($S = 1.5$)Full width $\Gamma < 2.1$ MeV, CL = 90% $\bar{D}^*(2007)^0$ modes are charge conjugates of modes below. **$D^*(2007)^0$ DECAY MODES**Fraction (Γ_i/Γ) p (MeV/c)

$D^0 \pi^0$	$(64.7 \pm 0.9) \%$	43
$D^0 \gamma$	$(35.3 \pm 0.9) \%$	137

 $D^*(2010)^{\pm}$

$I(J^P) = \frac{1}{2}(1^-)$

*I, J, P need confirmation.*Mass $m = 2010.26 \pm 0.05$ MeV $m_{D^*(2010)^+} - m_{D^+} = 140.603 \pm 0.015$ MeV $m_{D^*(2010)^+} - m_{D^0} = 145.4257 \pm 0.0017$ MeVFull width $\Gamma = 83.4 \pm 1.8$ keV $D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^{\pm}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0\pi^+$	(67.7 ± 0.5) %	39
$D^+\pi^0$	(30.7 ± 0.5) %	38
$D^+\gamma$	(1.6 ± 0.4) %	136

$D_0^*(2300)^0$

$$I(J^P) = \frac{1}{2}(0^+)$$

was $D_0^*(2400)^0$

Mass $m = 2300 \pm 19$ MeV
 Full width $\Gamma = 274 \pm 40$ MeV

$D_0^*(2300)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^+\pi^-$	seen	369

$D_1(2420)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 2420.8 \pm 0.5$ MeV ($S = 1.3$)
 $m_{D_1^0} - m_{D^{*+}} = 410.6 \pm 0.5$ MeV ($S = 1.3$)
 Full width $\Gamma = 31.7 \pm 2.5$ MeV ($S = 3.5$)

$\overline{D}_1(2420)^0$ modes are charge conjugates of modes below.

$D_1(2420)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^*(2010)^+\pi^-$	seen	353
$D^0\pi^+\pi^-$	seen	425
$D^+\pi^-$	not seen	472
$D^{*0}\pi^+\pi^-$	not seen	279

$D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored.

Mass $m = 2460.7 \pm 0.4$ MeV ($S = 3.1$)
 $m_{D_2^{*0}} - m_{D^+} = 591.0 \pm 0.4$ MeV ($S = 2.9$)
 $m_{D_2^{*0}} - m_{D^{*+}} = 450.4 \pm 0.4$ MeV ($S = 2.9$)
 Full width $\Gamma = 47.5 \pm 1.1$ MeV ($S = 1.8$)

$\overline{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

$D_2^*(2460)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^+ \pi^-$	seen	505
$D^*(2010)^+ \pi^-$	seen	389
$D^0 \pi^+ \pi^-$	not seen	462
$D^{*0} \pi^+ \pi^-$	not seen	324

$D_2^*(2460)^\pm$	$I(J^P) = \frac{1}{2}(2^+)$
-------------------------------------	-----------------------------

$J^P = 2^+$ assignment strongly favored.

Mass $m = 2465.4 \pm 1.3$ MeV (S = 3.1)

$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7$ MeV

Full width $\Gamma = 46.7 \pm 1.2$ MeV

$D_2^*(2460)^-$ modes are charge conjugates of modes below.

$D_2^*(2460)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	seen	513
$D^{*0} \pi^+$	seen	396
$D^+ \pi^+ \pi^-$	not seen	462
$D^{*+} \pi^+ \pi^-$	not seen	326

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^- 2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] Submodes of the $D^+ \rightarrow K^- 2\pi^+ \pi^0$ and $K_S^0 2\pi^+ \pi^-$ modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 Review, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $\overline{K}^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
- [l] This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.
- [n] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to $6.17 \pm 0.17\%$.
- [o] This is a doubly Cabibbo-suppressed mode.
- [p] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [q] This branching fraction includes all the decay modes of the resonance in the final state.
- [r] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [s] This limit is for either D^0 or \overline{D}^0 to $p e^-$.
- [t] This limit is for either D^0 or \overline{D}^0 to $\overline{p} e^+$.