

$$I(J^P) = 0(0^-)$$


The angular distributions of the decays of the  $\phi$  and  $\bar{K}^*(892)^0$  in the  $\phi\pi^+$  and  $K^+\bar{K}^*(892)^0$  modes strongly indicate that the spin is zero. The parity given is that expected of a  $c\bar{s}$  ground state.

## $D_s^\pm$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements. Measurements of the  $D_s^\pm$  mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements have been omitted altogether.

| VALUE (MeV)   | EVTS | DOCUMENT ID   | TECN | COMMENT                                    |
|---|------|---|------|--|
| <b>1968.35 ± 0.07 OUR FIT</b>   |      |   |      |  |
| <b>1969.0 ± 1.4 OUR AVERAGE</b>   |      | Error includes scale factor of 1.5. See the ideogram below. |      |  |
| 1967.0 ± 1.0 ± 1.0  | 54   | BARLAG  | 90C  | ACCM $\pi^-$ Cu 230 GeV                    |
| 1969.3 ± 1.4 ± 1.4  |      | ALBRECHT  | 88   | ARG $e^+e^-$ 9.4–10.6 GeV                  |
| 1972.7 ± 1.5 ± 1.0  | 21   | BECKER  | 87B  | SILI 200 GeV $\pi, K, p$                   |
| 1972.4 ± 3.7 ± 3.7  | 27   | BLAYLOCK  | 87   | MRK3 $e^+e^-$ 4.14 GeV                     |
| 1963 ± 3 ± 3  | 30   | DERRICK   | 85B  | HRS $e^+e^-$ 29 GeV                        |
| 1970 ± 5 ± 5  | 104  | CHEN  | 83C  | CLEO $e^+e^-$ 10.5 GeV                     |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |      |   |      |  |
| 1968.3 ± 0.7 ± 0.7  | 290  | <sup>1</sup> ANJOS  | 88   | E691 Photoproduction                       |
| 1980 ± 15   | 6    | USHIDA  | 86   | EMUL $\nu$ wideband                        |
| 1973.6 ± 2.6 ± 3.0  | 163  | ALBRECHT  | 85D  | ARG $e^+e^-$ 10 GeV                        |
| 1948 ± 28 ± 10  | 65   | AIHARA  | 84D  | TPC $e^+e^-$ 29 GeV                        |
| 1975 ± 9 ± 10   | 49   | ALTHOFF   | 84   | TASS $e^+e^-$ 14–25 GeV                    |
| 1975 ± 4  | 3    | BAILEY  | 84   | ACCM hadron <sup>+</sup> Be → $\phi\pi^+X$ |

<sup>1</sup> ANJOS 88 enters the fit via  $m_{D_s^\pm} - m_{D^\pm}$  (see below).



### $m_{D_s^\pm} - m_{D^\pm}$

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

| VALUE (MeV)                   | EVTs | DOCUMENT ID | TECN | COMMENT                                |
|-------------------------------|------|-------------|------|--|
| <b>98.69±0.05 OUR FIT</b>     |      |             |      |  |
| <b>98.69±0.05 OUR AVERAGE</b> |      |             |      |  |
| 98.68±0.03±0.04               |      | AAIJ        | 13V  | LHCB $D_s^+ \rightarrow K^+ K^- \pi^+$ |
| 99.41±0.38±0.21               |      | ACOSTA      | 03D  | CDF2 $\bar{p}p$ , $\sqrt{s}=1.96$ TeV  |
| 98.4 ±0.1 ±0.3                | 48k  | AUBERT      | 02G  | BABR $e^+e^- \approx \Upsilon(4S)$     |
| 99.5 ±0.6 ±0.3                |      | BROWN       | 94   | CLE2 $e^+e^- \approx \Upsilon(4S)$     |
| 98.5 ±1.5                     | 555  | CHEN        | 89   | CLEO $e^+e^-$ 10.5 GeV                 |
| 99.0 ±0.8                     | 290  | ANJOS       | 88   | E691 Photoproduction                   |

### $D_s^\pm$ MEAN LIFE

Measurements with an error greater than  $100 \times 10^{-15}$  s or with fewer than 100 events have been omitted from the Listings.

| VALUE ( $10^{-15}$ s)      | EVTs  | DOCUMENT ID       | TECN | COMMENT                                  |
|----------------------------|-------|-------------------|------|--|
| <b>504 ± 4 OUR AVERAGE</b> |       |                   |      | Error includes scale factor of 1.2.      |
| 506.4± 3.0± 1.7±1.7        |       | <sup>1</sup> AAIJ | 17AN | LHCB $pp$ at 7, 8 TeV                    |
| 507.4± 5.5± 5.1            | 13.6k | LINK              | 05J  | FOCS $\phi \pi^+$ and $\bar{K}^{*0} K^+$ |
| 472.5±17.2± 6.6            | 760   | IORI              | 01   | SELX 600 GeV $\Sigma^-, \pi^-, p$        |

|  |      |                        |     |      |                                  |
|--|------|------------------------|-----|------|----------------------------------|
| 518 ±14 ± 7                            | 1662 | AITALA                 | 99  | E791 | $\pi^-$ nucleus, 500 GeV         |
| 486.3±15.0 <sup>+</sup> <sub>5.1</sub> | 2167 | <sup>2</sup> BONVICINI | 99  | CLE2 | $e^+e^- \approx \mathcal{R}(4S)$ |
| 475 ±20 ± 7                            | 900  | FRABETTI               | 93F | E687 | $\gamma\text{Be}, \phi\pi^+$     |
| 500 ±60 ±30                            | 104  | FRABETTI               | 90  | E687 | $\gamma\text{Be}, \phi\pi^+$     |
| 470 ±40 ±20                            | 228  | RAAB                   | 88  | E691 | Photoproduction                  |

<sup>1</sup>This AAIJ 17AN value is derived from the difference between the  $D_S^-$  and  $D^-$  widths.

The 3rd uncertainty,  $\pm 1.7 \times 10^{-15}$  s, arises from the uncertainty of the  $D^-$  width.

<sup>2</sup>BONVICINI 99 obtains  $1.19 \pm 0.04$  for the ratio of  $D_S^+$  to  $D^0$  lifetimes.

## $D_S^+$ DECAY MODES

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance.  $D_S^-$  modes are charge conjugates of the modes below.

| Mode  | Fraction ( $\Gamma_j/\Gamma$ )   | Scale factor/<br>Confidence level |
|---|----------------------------------|-----------------------------------|
| <b>Inclusive modes</b>  |                                  |                                   |
| $\Gamma_1$ $e^+$ semileptonic                                   | [a] ( 6.5 ±0.4 ) %               |                                   |
| $\Gamma_2$ $\pi^+$ anything                                     | (119.3 ±1.4 ) %                  |                                   |
| $\Gamma_3$ $\pi^-$ anything                                     | ( 43.2 ±0.9 ) %                  |                                   |
| $\Gamma_4$ $\pi^0$ anything                                     | (123 ±7 ) %                      |                                   |
| $\Gamma_5$ $K^-$ anything                                       | ( 18.7 ±0.5 ) %                  |                                   |
| $\Gamma_6$ $K^+$ anything                                       | ( 28.9 ±0.7 ) %                  |                                   |
| $\Gamma_7$ $K_S^0$ anything                                     | ( 19.0 ±1.1 ) %                  |                                   |
| $\Gamma_8$ $\eta$ anything                                      | [b] ( 29.9 ±2.8 ) %              |                                   |
| $\Gamma_9$ $\omega$ anything                                    | ( 6.1 ±1.4 ) %                   |                                   |
| $\Gamma_{10}$ $\eta'$ anything                                  | [c] ( 10.3 ±1.4 ) %              | S=1.1                             |
| $\Gamma_{11}$ $f_0(980)$ anything, $f_0 \rightarrow \pi^+\pi^-$ | < 1.3 %                          | CL=90%                            |
| $\Gamma_{12}$ $\phi$ anything                                   | ( 15.7 ±1.0 ) %                  |                                   |
| $\Gamma_{13}$ $K^+K^-$ anything                                 | ( 15.8 ±0.7 ) %                  |                                   |
| $\Gamma_{14}$ $K_S^0K^+$ anything                               | ( 5.8 ±0.5 ) %                   |                                   |
| $\Gamma_{15}$ $K_S^0K^-$ anything                               | ( 1.9 ±0.4 ) %                   |                                   |
| $\Gamma_{16}$ $2K_S^0$ anything                                 | ( 1.70±0.32 ) %                  |                                   |
| $\Gamma_{17}$ $2K^+$ anything                                   | < 2.6 × 10 <sup>-3</sup>         | CL=90%                            |
| $\Gamma_{18}$ $2K^-$ anything                                   | < 6 × 10 <sup>-4</sup>           | CL=90%                            |
| <b>Leptonic and semileptonic modes</b>                          |                                  |                                   |
| $\Gamma_{19}$ $e^+\nu_e$  | < 8.3 × 10 <sup>-5</sup>         | CL=90%                            |
| $\Gamma_{20}$ $\mu^+\nu_\mu$                                    | ( 5.49±0.16 ) × 10 <sup>-3</sup> |                                   |
| $\Gamma_{21}$ $\tau^+\nu_\tau$                                  | ( 5.48±0.23 ) %                  |                                   |
| $\Gamma_{22}$ $\gamma e^+\nu_e$                                 | < 1.3 × 10 <sup>-4</sup>         | CL=90%                            |
| $\Gamma_{23}$ $K^+K^-e^+\nu_e$                                  | —                                |                                   |
| $\Gamma_{24}$ $\phi e^+\nu_e$                                   | [d] ( 2.39±0.16 ) %              | S=1.3                             |
| $\Gamma_{25}$ $\phi\mu^+\nu_\mu$                                | ( 1.9 ±0.5 ) %                   |                                   |

|               |   |     |                                  |        |
|---------------|---|-----|----------------------------------|--------|
| $\Gamma_{26}$ | $\eta e^+ \nu_e + \eta'(958) e^+ \nu_e$           | [d] | ( 3.03±0.24 ) %                  |        |
| $\Gamma_{27}$ | $\eta e^+ \nu_e$                                  | [d] | ( 2.32±0.08 ) %                  |        |
| $\Gamma_{28}$ | $\eta'(958) e^+ \nu_e$                            | [d] | ( 8.0 ±0.7 ) × 10 <sup>-3</sup>  |        |
| $\Gamma_{29}$ | $\eta \mu^+ \nu_\mu$                              |     | ( 2.4 ±0.5 ) %                   |        |
| $\Gamma_{30}$ | $\eta'(958) \mu^+ \nu_\mu$                        |     | ( 1.1 ±0.5 ) %                   |        |
| $\Gamma_{31}$ | $\omega e^+ \nu_e$                                | [e] | < 2.0 × 10 <sup>-3</sup>         | CL=90% |
| $\Gamma_{32}$ | $K^0 e^+ \nu_e$                                   |     | ( 3.4 ±0.4 ) × 10 <sup>-3</sup>  |        |
| $\Gamma_{33}$ | $K^*(892)^0 e^+ \nu_e$                            | [d] | ( 2.15±0.28 ) × 10 <sup>-3</sup> | S=1.1  |
| $\Gamma_{34}$ | $f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-$ |     |                                  |        |

### Hadronic modes with a $K\bar{K}$ pair

|               |  |       |   |        |
|---------------|--|-------|---|--------|
| $\Gamma_{35}$ | $K^+ K_S^0$  |       | ( 1.46±0.04 ) %                           | S=1.1  |
| $\Gamma_{36}$ | $K^+ K_L^0$  |       | ( 1.49±0.06 ) %                           |        |
| $\Gamma_{37}$ | $K^+ \bar{K}^0$  |       | ( 2.95±0.14 ) %                           |        |
| $\Gamma_{38}$ | $K^+ K^- \pi^+$  | [f]   | ( 5.39±0.15 ) %                           | S=1.2  |
| $\Gamma_{39}$ | $\phi \pi^+$   | [d,g] | ( 4.5 ±0.4 ) %                            |        |
| $\Gamma_{40}$ | $\phi \pi^+, \phi \rightarrow K^+ K^-$                                       | [g]   | ( 2.24±0.08 ) %                           |        |
| $\Gamma_{41}$ | $K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$                   |       | ( 2.58±0.08 ) %                           |        |
| $\Gamma_{42}$ | $f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$                                    |       | ( 1.14±0.31 ) %                           |        |
| $\Gamma_{43}$ | $f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-$                                   |       | ( 7 ±5 ) × 10 <sup>-4</sup>               |        |
| $\Gamma_{44}$ | $f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$                                   |       | ( 6.6 ±2.8 ) × 10 <sup>-4</sup>           |        |
| $\Gamma_{45}$ | $K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^{*0} \rightarrow K^- \pi^+$              |       | ( 1.8 ±0.4 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{46}$ | $K^+ K_S^0 \pi^0$  |       | ( 1.52±0.22 ) %                           |        |
| $\Gamma_{47}$ | $2K_S^0 \pi^+$   |       | ( 7.7 ±0.6 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{48}$ | $K^0 \bar{K}^0 \pi^+$  |       | —   |        |
| $\Gamma_{49}$ | $K^*(892)^+ \bar{K}^0$   | [d]   | ( 5.4 ±1.2 ) %                            |        |
| $\Gamma_{50}$ | $K^+ K^- \pi^+ \pi^0$  |       | ( 6.2 ±0.6 ) %                            | S=1.1  |
| $\Gamma_{51}$ | $\phi \rho^+$  | [d]   | ( 8.4 <sup>+1.9</sup> <sub>-2.3</sub> ) % |        |
| $\Gamma_{52}$ | $K_S^0 K^- 2\pi^+$   |       | ( 1.65±0.10 ) %                           |        |
| $\Gamma_{53}$ | $K^*(892)^+ \bar{K}^*(892)^0$  | [d]   | ( 7.2 ±2.6 ) %                            |        |
| $\Gamma_{54}$ | $K^+ K_S^0 \pi^+ \pi^-$  |       | ( 9.9 ±0.8 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{55}$ | $K^+ K^- 2\pi^+ \pi^-$   |       | ( 8.6 ±1.5 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{56}$ | $\phi 2\pi^+ \pi^-$  | [d]   | ( 1.21±0.16 ) %                           |        |
| $\Gamma_{57}$ | $\phi \rho^0 \pi^+, \phi \rightarrow K^+ K^-$                                |       | ( 6.5 ±1.3 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{58}$ | $\phi a_1(1260)^+, \phi \rightarrow K^+ K^-, a_1^+ \rightarrow \rho^0 \pi^+$ |       | ( 7.4 ±1.2 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{59}$ | $\phi 2\pi^+ \pi^- \text{ non-}\rho, \phi \rightarrow K^+ K^-$               |       | ( 1.8 ±0.7 ) × 10 <sup>-3</sup>           |        |
| $\Gamma_{60}$ | $K^+ K^- \rho^0 \pi^+ \text{ non-}\phi$                                      | <     | 2.6 × 10 <sup>-4</sup>                    | CL=90% |
| $\Gamma_{61}$ | $K^+ K^- 2\pi^+ \pi^- \text{ nonresonant}$                                   |       | ( 9 ±7 ) × 10 <sup>-4</sup>               |        |
| $\Gamma_{62}$ | $2K_S^0 2\pi^+ \pi^-$  |       | ( 8.4 ±3.5 ) × 10 <sup>-4</sup>           |        |

### Hadronic modes without $K$ 's

|               |  |                       |                  |        |
|---------------|--|-----------------------|------------------|--------|
| $\Gamma_{63}$ | $\pi^+ \pi^0$  | $< 3.4$               | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{64}$ | $2\pi^+ \pi^-$   | ( 1.08±0.04 ) %       |                  | S=1.1  |
| $\Gamma_{65}$ | $\rho^0 \pi^+$   | ( 1.9 ±1.2 )          | $\times 10^{-4}$ |        |
| $\Gamma_{66}$ | $\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$                                  | [h] ( 9.0 ±0.4 )      | $\times 10^{-3}$ |        |
| $\Gamma_{67}$ | $f_0(980) \pi^+, f_0 \rightarrow \pi^+ \pi^-$                          |                       |                  |        |
| $\Gamma_{68}$ | $f_0(1370) \pi^+, f_0 \rightarrow \pi^+ \pi^-$                         |                       |                  |        |
| $\Gamma_{69}$ | $f_0(1500) \pi^+, f_0 \rightarrow \pi^+ \pi^-$                         |                       |                  |        |
| $\Gamma_{70}$ | $f_2(1270) \pi^+, f_2 \rightarrow \pi^+ \pi^-$                         | ( 1.09±0.20 )         | $\times 10^{-3}$ |        |
| $\Gamma_{71}$ | $\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$                   | ( 3.0 ±1.9 )          | $\times 10^{-4}$ |        |
| $\Gamma_{72}$ | $\pi^+ 2\pi^0$   | ( 6.5 ±1.3 )          | $\times 10^{-3}$ |        |
| $\Gamma_{73}$ | $2\pi^+ \pi^- \pi^0$   | —                     |                  |        |
| $\Gamma_{74}$ | $\eta \pi^+$   | [d] ( 1.68±0.10 ) %   |                  | S=1.2  |
| $\Gamma_{75}$ | $\omega \pi^+$   | [d] ( 1.92±0.30 )     | $\times 10^{-3}$ |        |
| $\Gamma_{76}$ | $3\pi^+ 2\pi^-$  | ( 7.9 ±0.8 )          | $\times 10^{-3}$ |        |
| $\Gamma_{77}$ | $2\pi^+ \pi^- 2\pi^0$  | —                     |                  |        |
| $\Gamma_{78}$ | $\eta \rho^+$  | [d] ( 8.9 ±0.8 ) %    |                  |        |
| $\Gamma_{79}$ | $\eta \pi^+ \pi^0$   | ( 9.5 ±0.5 ) %        |                  |        |
| $\Gamma_{80}$ | $\eta (\pi^+ \pi^0)_{P\text{-wave}}$                                   | ( 5.1 ±3.1 )          | $\times 10^{-3}$ |        |
| $\Gamma_{81}$ | $a_0(980)^{+0} \pi^{0+},$<br>$a_0(980)^{+0} \rightarrow \eta \pi^{+0}$ | ( 2.2 ±0.4 ) %        |                  |        |
| $\Gamma_{82}$ | $\omega \pi^+ \pi^0$   | [d] ( 2.8 ±0.7 ) %    |                  |        |
| $\Gamma_{83}$ | $3\pi^+ 2\pi^- \pi^0$  | ( 4.9 ±3.2 ) %        |                  |        |
| $\Gamma_{84}$ | $\omega 2\pi^+ \pi^-$  | [d] ( 1.6 ±0.5 ) %    |                  |        |
| $\Gamma_{85}$ | $\eta'(958) \pi^+$   | [c,d] ( 3.94±0.25 ) % |                  |        |
| $\Gamma_{86}$ | $3\pi^+ 2\pi^- 2\pi^0$   | —                     |                  |        |
| $\Gamma_{87}$ | $\omega \eta \pi^+$  | [d] $< 2.13$ %        |                  | CL=90% |
| $\Gamma_{88}$ | $\eta'(958) \rho^+$  | [c,d] ( 5.8 ±1.5 ) %  |                  |        |
| $\Gamma_{89}$ | $\eta'(958) \pi^+ \pi^0$   | ( 5.6 ±0.8 ) %        |                  |        |
| $\Gamma_{90}$ | $\eta'(958) \pi^+ \pi^0$ nonresonant                                   | $< 5.1$ %             |                  | CL=90% |

### Modes with one or three $K$ 's

|                |  |                   |                  |  |
|----------------|--|-------------------|------------------|--|
| $\Gamma_{91}$  | $K^+ \pi^0$  | ( 7.4 ±0.6 )      | $\times 10^{-4}$ |  |
| $\Gamma_{92}$  | $K_S^0 \pi^+$  | ( 1.10±0.05 )     | $\times 10^{-3}$ |  |
| $\Gamma_{93}$  | $K^+ \eta$   | [d] ( 1.60±0.11 ) | $\times 10^{-3}$ |  |
| $\Gamma_{94}$  | $K^+ \omega$   | [d] ( 8.7 ±2.5 )  | $\times 10^{-4}$ |  |
| $\Gamma_{95}$  | $K^+ \eta'(958)$                                       | [d] ( 2.65±0.25 ) | $\times 10^{-3}$ |  |
| $\Gamma_{96}$  | $K^+ \pi^+ \pi^-$                                      | ( 6.5 ±0.4 )      | $\times 10^{-3}$ |  |
| $\Gamma_{97}$  | $K^+ \rho^0$   | ( 2.5 ±0.4 )      | $\times 10^{-3}$ |  |
| $\Gamma_{98}$  | $K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-$     | ( 6.9 ±2.4 )      | $\times 10^{-4}$ |  |
| $\Gamma_{99}$  | $K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$       | ( 1.41±0.24 )     | $\times 10^{-3}$ |  |
| $\Gamma_{100}$ | $K^*(1410)^0 \pi^+, K^{*0} \rightarrow$<br>$K^+ \pi^-$ | ( 1.23±0.28 )     | $\times 10^{-3}$ |  |
| $\Gamma_{101}$ | $K^*(1430)^0 \pi^+, K^{*0} \rightarrow$<br>$K^+ \pi^-$ | ( 5.0 ±3.5 )      | $\times 10^{-4}$ |  |

|                |                                      |         |                                  |        |
|----------------|--------------------------------------|---------|----------------------------------|--------|
| $\Gamma_{102}$ | $K^+ \pi^+ \pi^-$ nonresonant        |         | $(1.03 \pm 0.34) \times 10^{-3}$ |        |
| $\Gamma_{103}$ | $K^0 \pi^+ \pi^0$                    |         | $(1.00 \pm 0.18) \%$             |        |
| $\Gamma_{104}$ | $K_S^0 2\pi^+ \pi^-$                 |         | $(3.0 \pm 1.1) \times 10^{-3}$   |        |
| $\Gamma_{105}$ | $K^+ \omega \pi^0$                   | $[d] <$ | $8.2 \times 10^{-3}$             | CL=90% |
| $\Gamma_{106}$ | $K^+ \omega \pi^+ \pi^-$             | $[d] <$ | $5.4 \times 10^{-3}$             | CL=90% |
| $\Gamma_{107}$ | $K^+ \omega \eta$                    | $[d] <$ | $7.9 \times 10^{-3}$             | CL=90% |
| $\Gamma_{108}$ | $2K^+ K^-$                           |         | $(2.16 \pm 0.20) \times 10^{-4}$ |        |
| $\Gamma_{109}$ | $\phi K^+, \phi \rightarrow K^+ K^-$ |         | $(8.8 \pm 2.0) \times 10^{-5}$   |        |

### Doubly Cabibbo-suppressed modes

|                |  |  |                                  |  |
|----------------|--|--|----------------------------------|--|
| $\Gamma_{110}$ | $2K^+ \pi^-$                                   |  | $(1.28 \pm 0.04) \times 10^{-4}$ |  |
| $\Gamma_{111}$ | $K^+ K^*(892)^0, K^{*0} \rightarrow K^+ \pi^-$ |  | $(6.0 \pm 3.4) \times 10^{-5}$   |  |

### Baryon-antibaryon mode

|                |                       |     |                                  |        |
|----------------|-----------------------|-----|----------------------------------|--------|
| $\Gamma_{112}$ | $p \bar{n}$           |     | $(1.22 \pm 0.11) \times 10^{-3}$ |        |
| $\Gamma_{113}$ | $p \bar{p} e^+ \nu_e$ | $<$ | $2.0 \times 10^{-4}$             | CL=90% |

### $\Delta C = 1$ weak neutral current (C1) modes, Lepton family number (LF), or Lepton number (L) violating modes

|                |  |   |                      |        |
|----------------|--|---|----------------------|--------|
| $\Gamma_{114}$ | $\pi^+ e^+ e^-$                        | $[i] <$   | $1.3 \times 10^{-5}$ | CL=90% |
| $\Gamma_{115}$ | $\pi^+ \phi, \phi \rightarrow e^+ e^-$ | $[j] (6 \begin{smallmatrix} +8 \\ -4 \end{smallmatrix}) \times$ | $10^{-6}$            |        |
| $\Gamma_{116}$ | $\pi^+ \mu^+ \mu^-$                    | $[i] <$   | $4.1 \times 10^{-7}$ | CL=90% |
| $\Gamma_{117}$ | $K^+ e^+ e^-$                          | C1 $<$  | $3.7 \times 10^{-6}$ | CL=90% |
| $\Gamma_{118}$ | $K^+ \mu^+ \mu^-$                      | C1 $<$  | $2.1 \times 10^{-5}$ | CL=90% |
| $\Gamma_{119}$ | $K^*(892)^+ \mu^+ \mu^-$               | C1 $<$  | $1.4 \times 10^{-3}$ | CL=90% |
| $\Gamma_{120}$ | $\pi^+ e^+ \mu^-$                      | LF $<$  | $1.2 \times 10^{-5}$ | CL=90% |
| $\Gamma_{121}$ | $\pi^+ e^- \mu^+$                      | LF $<$  | $2.0 \times 10^{-5}$ | CL=90% |
| $\Gamma_{122}$ | $K^+ e^+ \mu^-$                        | LF $<$  | $1.4 \times 10^{-5}$ | CL=90% |
| $\Gamma_{123}$ | $K^+ e^- \mu^+$                        | LF $<$  | $9.7 \times 10^{-6}$ | CL=90% |
| $\Gamma_{124}$ | $\pi^- 2e^+$                           | L $<$   | $4.1 \times 10^{-6}$ | CL=90% |
| $\Gamma_{125}$ | $\pi^- 2\mu^+$                         | L $<$   | $1.2 \times 10^{-7}$ | CL=90% |
| $\Gamma_{126}$ | $\pi^- e^+ \mu^+$                      | L $<$   | $8.4 \times 10^{-6}$ | CL=90% |
| $\Gamma_{127}$ | $K^- 2e^+$                             | L $<$   | $5.2 \times 10^{-6}$ | CL=90% |
| $\Gamma_{128}$ | $K^- 2\mu^+$                           | L $<$   | $1.3 \times 10^{-5}$ | CL=90% |
| $\Gamma_{129}$ | $K^- e^+ \mu^+$                        | L $<$   | $6.1 \times 10^{-6}$ | CL=90% |
| $\Gamma_{130}$ | $K^*(892)^- 2\mu^+$                    | L $<$   | $1.4 \times 10^{-3}$ | CL=90% |

[a] This is the purely  $e^+$  semileptonic branching fraction: the  $e^+$  fraction from  $\tau^+$  decays has been subtracted off. The sum of our (non- $\tau$ )  $e^+$  exclusive fractions — an  $e^+ \nu_e$  with an  $\eta, \eta', \phi, K^0$ , or  $K^{*0}$  — is  $5.99 \pm 0.31 \%$ .

[b] This fraction includes  $\eta$  from  $\eta'$  decays.

- [c] The sum of our exclusive  $\eta'$  fractions —  $\eta' e^+ \nu_e$ ,  $\eta' \mu^+ \nu_\mu$ ,  $\eta' \pi^+$ ,  $\eta' \rho^+$ , and  $\eta' K^+$  — is  $11.8 \pm 1.6\%$ .
- [d] This branching fraction includes all the decay modes of the final-state resonance.
- [e] A test for  $u\bar{u}$  or  $d\bar{d}$  content in the  $D_s^+$ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and  $\omega$ - $\phi$  mixing is an unlikely explanation for any fraction above about  $2 \times 10^{-4}$ .
- [f] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
- [g] We decouple the  $D_s^+ \rightarrow \phi \pi^+$  branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the  $D_s^+ \rightarrow \phi \pi^+$ ,  $\phi \rightarrow K^+ K^-$  branching fraction obtained from the Dalitz-plot analysis of  $D_s^+ \rightarrow K^+ K^- \pi^+$ . That is, the ratio of these two branching fractions is not exactly the  $\phi \rightarrow K^+ K^-$  branching fraction 0.491.
- [h] This is the average of a model-independent and a  $K$ -matrix parametrization of the  $\pi^+ \pi^-$   $S$ -wave and is a sum over several  $f_0$  mesons.
- [i] This mode is not a useful test for a  $\Delta C=1$  weak neutral current because both quarks must change flavor in this decay.
- [j] This is *not* a test for the  $\Delta C=1$  weak neutral current, but leads to the  $\pi^+ \ell^+ \ell^-$  final state.

---

### CONSTRAINED FIT INFORMATION

An overall fit to 13 branching ratios uses 18 measurements and one constraint to determine 10 parameters. The overall fit has a  $\chi^2 = 6.8$  for 9 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

|          |          |          |          |          |          |          |          |          |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| $x_{38}$ | 51       |          |          |          |          |          |          |          |
| $x_{50}$ | 22       | 29       |          |          |          |          |          |          |
| $x_{52}$ | 28       | 30       | 11       |          |          |          |          |          |
| $x_{54}$ | 23       | 25       | 13       | 38       |          |          |          |          |
| $x_{64}$ | 34       | 50       | 19       | 19       | 17       |          |          |          |
| $x_{74}$ | -7       | -7       | -15      | 1        | -7       | -7       |          |          |
| $x_{75}$ | 0        | 0        | -1       | 0        | 0        | 0        | 4        |          |
| $x_{96}$ | 5        | 12       | -6       | 7        | 1        | 3        | 16       | 1        |
|          | $x_{35}$ | $x_{38}$ | $x_{50}$ | $x_{52}$ | $x_{54}$ | $x_{64}$ | $x_{74}$ | $x_{75}$ |

---

See the related review(s):

[D<sub>s</sub><sup>+</sup> Branching Fractions](#)**D<sub>s</sub><sup>+</sup> BRANCHING RATIOS**

A number of older, now obsolete results have been omitted. They may be found in earlier editions.

**Inclusive modes****Γ(e<sup>+</sup> semileptonic)/Γ<sub>total</sub>****Γ<sub>1</sub>/Γ**

This is the purely e<sup>+</sup> semileptonic branching fraction: the e<sup>+</sup> fraction from τ<sup>+</sup> decays has been subtracted off. The sum of our (non-τ) e<sup>+</sup> exclusive fractions — an e<sup>+</sup>ν<sub>e</sub> with an η, η', φ, K<sup>0</sup>, or K\*<sup>0</sup> — is 5.99 ± 0.31 %.

| VALUE (%)             | EVTS     | DOCUMENT ID        | TECN | COMMENT  |
|-----------------------|----------|--------------------|------|--|
| <b>6.52±0.39±0.15</b> | 536 ± 29 | <sup>1</sup> ASNER | 10   | CLEO e <sup>+</sup> e <sup>-</sup> at 3774 MeV |

<sup>1</sup>Using the D<sub>s</sub><sup>+</sup> and D<sup>0</sup> lifetimes, ASNER 10 finds that the ratio of the D<sub>s</sub><sup>+</sup> and D<sup>0</sup> semileptonic widths is 0.828 ± 0.051 ± 0.025.

**Γ(π<sup>+</sup> anything)/Γ<sub>total</sub>****Γ<sub>2</sub>/Γ**

Events with two π<sup>+</sup>'s count twice, etc. But π<sup>+</sup>'s from K<sub>S</sub><sup>0</sup> → π<sup>+</sup>π<sup>-</sup> are not included.

| VALUE (%)            | DOCUMENT ID | TECN | COMMENT  |
|----------------------|-------------|------|--|
| <b>119.3±1.2±0.7</b> | DOBBS       | 09   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |

**Γ(π<sup>-</sup> anything)/Γ<sub>total</sub>****Γ<sub>3</sub>/Γ**

Events with two π<sup>-</sup>'s count twice, etc. But π<sup>-</sup>'s from K<sub>S</sub><sup>0</sup> → π<sup>+</sup>π<sup>-</sup> are not included.

| VALUE (%)           | DOCUMENT ID | TECN | COMMENT  |
|---------------------|-------------|------|--|
| <b>43.2±0.9±0.3</b> | DOBBS       | 09   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |

**Γ(π<sup>0</sup> anything)/Γ<sub>total</sub>****Γ<sub>4</sub>/Γ**

Events with two π<sup>0</sup>'s count twice, etc. But π<sup>0</sup>'s from K<sub>S</sub><sup>0</sup> → 2π<sup>0</sup> are not included.

| VALUE (%)            | DOCUMENT ID | TECN | COMMENT  |
|----------------------|-------------|------|--|
| <b>123.4±3.8±5.3</b> | DOBBS       | 09   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |

**Γ(K<sup>-</sup> anything)/Γ<sub>total</sub>****Γ<sub>5</sub>/Γ**

| VALUE (%)           | DOCUMENT ID | TECN | COMMENT  |
|---------------------|-------------|------|--|
| <b>18.7±0.5±0.2</b> | DOBBS       | 09   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |

**Γ(K<sup>+</sup> anything)/Γ<sub>total</sub>****Γ<sub>6</sub>/Γ**

| VALUE (%)           | DOCUMENT ID | TECN | COMMENT  |
|---------------------|-------------|------|--|
| <b>28.9±0.6±0.3</b> | DOBBS       | 09   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |

**Γ(K<sub>S</sub><sup>0</sup> anything)/Γ<sub>total</sub>****Γ<sub>7</sub>/Γ**

| VALUE (%)           | DOCUMENT ID | TECN | COMMENT  |
|---------------------|-------------|------|--|
| <b>19.0±1.0±0.4</b> | DOBBS       | 09   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |



$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ This ratio includes  $\eta$  particles from  $\eta'$  decays.

| <u>VALUE (%)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|---|-------------|--------------------|-------------|----------------------|
| <b>29.9±2.2±1.7</b>   |             | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |                    |             |                      |
| 23.5±3.1±2.0  | 674 ± 91    | HUANG              | 06B CLEO    | See DOBBS 09         |

 $\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

| <u>VALUE (%)</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|--------------------|--------------------|-------------|----------------------|
| <b>6.1±1.4±0.3</b> | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

 $\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

| <u>VALUE (%)</u>  | <u>EVTS</u>                         | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                  |
|---|-------------------------------------|--------------------|-------------|---------------------------------|
| <b>10.3±1.4 OUR AVERAGE</b>   | Error includes scale factor of 1.1. |                    |             |                                 |
| 8.8±1.8±0.5   | 68                                  | ABLIKIM            | 15Z BES3    | 482 pb <sup>-1</sup> , 4009 MeV |
| 11.7±1.7±0.7  |                                     | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV            |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |                    |             |                                 |
| 8.7±1.9±0.8   | 68                                  | HUANG              | 06B CLEO    | See DOBBS 09                    |

 $\Gamma(f_0(980) \text{ anything}, f_0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|------------------|------------|--------------------|-------------|----------------------|
| <b>&lt;1.3</b>   | 90         | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

 $\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

| <u>VALUE (%)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|---|-------------|--------------------|-------------|----------------------|
| <b>15.7±0.8±0.6</b>   |             | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |                    |             |                      |
| 16.1±1.2±1.1  | 398 ± 27    | HUANG              | 06B CLEO    | See DOBBS 09         |

 $\Gamma(K^+K^- \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

| <u>VALUE (%)</u>    | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|---------------------|--------------------|-------------|----------------------|
| <b>15.8±0.6±0.3</b> | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

 $\Gamma(K_S^0 K^+ \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

| <u>VALUE (%)</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|--------------------|--------------------|-------------|----------------------|
| <b>5.8±0.5±0.1</b> | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

 $\Gamma(K_S^0 K^- \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

| <u>VALUE (%)</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|--------------------|--------------------|-------------|----------------------|
| <b>1.9±0.4±0.1</b> | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

 $\Gamma(2K_S^0 \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

| <u>VALUE (%)</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|--------------------|--------------------|-------------|----------------------|
| <b>1.7±0.3±0.1</b> | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

 $\Gamma(2K^+ \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|------------------|------------|--------------------|-------------|----------------------|
| <b>&lt;0.26</b>  | 90         | DOBBS              | 09 CLEO     | $e^+e^-$ at 4170 MeV |

| $\Gamma(2K^- \text{ anything})/\Gamma_{\text{total}}$ |     |             |      |         | $\Gamma_{18}/\Gamma$ |
|---|-----|-------------|------|---------|----------------------|
| VALUE (%)   | CL% | DOCUMENT ID | TECN | COMMENT |                      |
| <0.06   | 90  | DOBBS       | 09   | CLEO    | $e^+e^-$ at 4170 MeV |

———— Leptonic and semileptonic modes ————

See the related review(s):

[Leptonic Decays of Charged Pseudoscalar Mesons](#)

| $\Gamma(e^+ \nu_e)/\Gamma_{\text{total}}$ |     |                     |      |         | $\Gamma_{19}/\Gamma$                     |
|---|-----|---------------------|------|---------|--|
| VALUE                                     | CL% | DOCUMENT ID         | TECN | COMMENT |  |
| <0.83 × 10 <sup>-4</sup>                  | 90  | <sup>1</sup> ZUPANC | 13   | BELL    | $e^+e^-$ at $\Upsilon(4S), \Upsilon(5S)$ |
| <2.3 × 10 <sup>-4</sup>                   | 90  | DEL-AMO-SA..10J     | BABR |         | $e^+e^-$ , 10.58 GeV                     |
| <1.2 × 10 <sup>-4</sup>                   | 90  | ALEXANDER           | 09   | CLEO    | $e^+e^-$ at 4170 MeV                     |
| <1.3 × 10 <sup>-4</sup>                   | 90  | PEDLAR              | 07A  | CLEO    | See ALEXANDER 09                         |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>ZUPANC 13 also gives the limit as  $< 1.0 \times 10^{-4}$  at 95% CL.

| $\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ |      |             |      |         | $\Gamma_{20}/\Gamma$ |
|---|------|-------------|------|---------|----------------------|
| VALUE (units 10 <sup>-3</sup> )               | EVTS | DOCUMENT ID | TECN | COMMENT |                      |

See the note on "Decay Constants of Charged Pseudoscalar Mesons."

|   |          |                              |      |      |  |
|---|----------|------------------------------|------|------|--|
| <b>5.49 ± 0.16 OUR AVERAGE</b>  |          |                              |      |      |  |
| 5.49 ± 0.16 ± 0.15  | 1.1k     | ABLIKIM                      | 19E  | BES3 | $e^+e^-$ at 4178 MeV                     |
| 4.95 ± 0.67 ± 0.26  | 69       | <sup>1</sup> ABLIKIM         | 160  | BES3 | $e^+e^-$ at 4.009 GeV                    |
| 5.31 ± 0.28 ± 0.20  | 492 ± 26 | <sup>2</sup> ZUPANC          | 13   | BELL | $e^+e^-$ at $\Upsilon(4S), \Upsilon(5S)$ |
| 6.02 ± 0.38 ± 0.34  | 275 ± 17 | <sup>3</sup> DEL-AMO-SA..10J | BABR |      | $e^+e^-$ , 10.58 GeV                     |
| 5.65 ± 0.45 ± 0.17  | 235 ± 14 | ALEXANDER                    | 09   | CLEO | $e^+e^-$ at 4170 MeV                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |          |                              |      |      |  |
| 6.44 ± 0.76 ± 0.57  | 169 ± 18 | <sup>4</sup> WIDHALM         | 08   | BELL | See ZUPANC 13                            |
| 5.94 ± 0.66 ± 0.31  | 88       | <sup>5</sup> PEDLAR          | 07A  | CLEO | See ALEXANDER 09                         |
| 6.8 ± 1.1 ± 1.8   | 553      | <sup>6</sup> HEISTER         | 02I  | ALEP | Z decays                                 |

<sup>1</sup>ABLIKIM 160 value is constrained by the Standard Model ratio of  $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.76$ ; the unconstrained value is  $(0.517 \pm 0.075 \pm 0.021)\%$ . The constrained value is used to obtain the decay constant,  $f_{D_s^+} = (241.0 \pm 16.3 \pm 6.6)$  MeV.

<sup>2</sup>ZUPANC 13 uses both  $\mu^+ \nu$  and  $\tau^+ \nu$  events to get  $f_{D_s} = (255.5 \pm 4.2 \pm 5.1)$  MeV.

<sup>3</sup>DEL-AMO-SANCHEZ 10J uses  $\mu^+ \nu_\mu$  and  $\tau^+ \nu_\tau$  events together to get  $f_{D_s} = (258.6 \pm 6.4 \pm 7.5)$  MeV.

<sup>4</sup>WIDHALM 08 gets  $f_{D_s} = (275 \pm 16 \pm 12)$  MeV from the branching fraction.

<sup>5</sup>PEDLAR 07A also fits  $\mu^+$  and  $\tau^+$  events together and gets an effective  $\mu^+ \nu_\mu$  branching fraction of  $(6.38 \pm 0.59 \pm 0.33) \times 10^{-3}$

<sup>6</sup>This HEISTER 02I result is not actually an independent measurement of the absolute  $\mu^+ \nu_\mu$  branching fraction, but is in fact based on our  $\phi\pi^+$  branching fraction of 3.6 ± 0.9%, so it cannot be included in our overall fit. HEISTER 02I combines its  $D_s^+ \rightarrow \tau^+ \nu_\tau$  and  $\mu^+ \nu_\mu$  branching fractions to get  $f_{D_s} = (285 \pm 19 \pm 40)$  MeV.

### $\Gamma(\mu^+ \nu_\mu)/\Gamma(\phi\pi^+)$

$\Gamma_{20}/\Gamma_{39}$

See the note on “Decay Constants of Charged Pseudoscalar Mesons” above.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |              |                         |          |                              |
|-----------------------------|--------------|-------------------------|----------|------------------------------|
| $0.143 \pm 0.018 \pm 0.006$ | $489 \pm 55$ | <sup>1</sup> AUBERT     | 07V BABR | $e^+ e^- \approx \gamma(4S)$ |
| $0.23 \pm 0.06 \pm 0.04$    | 18           | <sup>2</sup> ALEXANDROV | 00 BEAT  | $\pi^-$ nucleus, 350 GeV     |
| $0.173 \pm 0.023 \pm 0.035$ | 182          | <sup>3</sup> CHADHA     | 98 CLE2  | $e^+ e^- \approx \gamma(4S)$ |
| $0.245 \pm 0.052 \pm 0.074$ | 39           | <sup>4</sup> ACOSTA     | 94 CLE2  | See CHADHA 98                |

<sup>1</sup> AUBERT 07V gets  $f_{D_s^+} = (283 \pm 17 \pm 16)$  MeV, using  $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = (4.71 \pm 0.46)\%$ .

<sup>2</sup> ALEXANDROV 00 uses  $f_D^2/f_{D_s}^2 = 0.82 \pm 0.09$  from a lattice-gauge-theory calculation to get the relative numbers of  $D^+ \rightarrow \mu^+ \nu_\mu$  and  $D_s^+ \rightarrow \mu^+ \nu_\mu$  events. The present result leads to  $f_{D_s} = (323 \pm 44 \pm 36)$  MeV.

<sup>3</sup> CHADHA 98 obtains  $f_{D_s} = (280 \pm 19 \pm 28 \pm 34)$  MeV from this measurement, using  $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = 0.036 \pm 0.009$ .

<sup>4</sup> ACOSTA 94 obtains  $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$  MeV from this measurement, using  $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = 0.037 \pm 0.009$ .

### $\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$

$\Gamma_{21}/\Gamma$

See the note on “Decay Constants of Charged Pseudoscalar Mesons” above.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

#### **5.48 ± 0.23 OUR AVERAGE**

|                                 |              |                              |          |   |
|---------------------------------|--------------|------------------------------|----------|---|
| $4.83 \pm 0.65 \pm 0.26$        | 33           | <sup>1</sup> ABLIKIM         | 160 BES3 | $e^+ e^-$ at 4.009 GeV                                      |
| $5.70 \pm 0.21^{+0.31}_{-0.30}$ | 2.2k         | <sup>2</sup> ZUPANC          | 13 BELL  | $e^+ e^-$ at $\gamma(4S)$ , $\gamma(5S)$                    |
| $4.96 \pm 0.37 \pm 0.57$        | $748 \pm 53$ | <sup>3</sup> DEL-AMO-SA..10J | BABR     | $e^- \bar{\nu}_e \nu_\tau$ , $\mu^- \bar{\nu}_\mu \nu_\tau$ |
| $6.42 \pm 0.81 \pm 0.18$        | $126 \pm 16$ | <sup>4</sup> ALEXANDER       | 09 CLEO  | $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$                   |
| $5.52 \pm 0.57 \pm 0.21$        | $155 \pm 17$ | <sup>4</sup> NAIK            | 09A CLEO | $\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau$                  |
| $5.30 \pm 0.47 \pm 0.22$        | $181 \pm 16$ | <sup>4</sup> ONYISI          | 09 CLEO  | $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$               |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                          |     |                       |          |  |
|--------------------------|-----|-----------------------|----------|--|
| $6.17 \pm 0.71 \pm 0.34$ | 102 | <sup>5</sup> ECKLUND  | 08 CLEO  | See ONYISI 09                                |
| $8.0 \pm 1.3 \pm 0.4$    | 47  | <sup>5</sup> PEDLAR   | 07A CLEO | See ALEXANDER 09                             |
| $5.79 \pm 0.77 \pm 1.84$ | 881 | <sup>6</sup> HEISTER  | 02I ALEP | Z decays                                     |
| $7.0 \pm 2.1 \pm 2.0$    | 22  | <sup>7</sup> ABBIENDI | 01L OPAL | $D_s^{*+} \rightarrow \gamma D_s^+$ from Z's |
| $7.4 \pm 2.8 \pm 2.4$    | 16  | <sup>8</sup> ACCIARRI | 97F L3   | $D_s^{*+} \rightarrow \gamma D_s^+$ from Z's |

<sup>1</sup> ABLIKIM 160 value is constrained by the Standard Model ratio of  $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.76$ ; the unconstrained value is  $(3.28 \pm 1.83 \pm 0.37)\%$ .

<sup>2</sup> ZUPANC 13 uses both  $\mu^+ \nu$  and  $\tau^+ \nu$  events to get  $f_{D_s} = (255.5 \pm 4.2 \pm 5.1)$  MeV.

<sup>3</sup> DEL-AMO-SANCHEZ 10J (with a small correction; see LEES 15D) uses  $\mu^+ \nu_\mu$  and  $\tau^+ \nu_\tau$  events together to get  $f_{D_s} = (259.9 \pm 6.6 \pm 7.6)$  MeV.

<sup>4</sup> ALEXANDER 09, NAIK 09A, and ONYISI 09 use different  $\tau$  decay modes and are independent. The three papers combined give  $f_{D_s} = (259.7 \pm 7.8 \pm 3.4)$  MeV.

<sup>5</sup> ECKLUND 08 and PEDLAR 07A are independent: ECKLUND 08 uses  $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$  events, PEDLAR 07A uses  $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$  events.

<sup>6</sup> HEISTER 02I combines its  $D_s^+ \rightarrow \tau^+ \nu_\tau$  and  $\mu^+ \nu_\mu$  branching fractions to get  $f_{D_s} = (285 \pm 19 \pm 40)$  MeV.

<sup>7</sup> This ABBIENDI 01L value gives a decay constant  $f_{D_s}$  of  $(286 \pm 44 \pm 41)$  MeV.

<sup>8</sup> The second ACCIARRI 97F error here combines in quadrature systematic (0.016) and normalization (0.018) errors. The branching fraction gives  $f_{D_s} = (309 \pm 58 \pm 33 \pm 38)$  MeV.

**$\Gamma(\tau^+ \nu_\tau)/\Gamma(\mu^+ \nu_\mu)$   $\Gamma_{21}/\Gamma_{20}$**

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                  |          |                      |    |  |
|----------------------------------|----------|----------------------|----|--|
| $10.73 \pm 0.69^{+0.56}_{-0.53}$ | 2.2k/492 | <sup>1</sup> ZUPANC  | 13 | BELL $e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$ |
| $11.0 \pm 1.4 \pm 0.6$           | 102      | <sup>2</sup> ECKLUND | 08 | CLEO See ONYISI 09                             |

<sup>1</sup> This ZUPANC 13 ratio is not independent of the separate  $\tau\nu$  and  $\mu\nu$  fractions listed above.

<sup>2</sup> This ECKLUND 08 value also uses results from PEDLAR 07A, and it is not independent of other results in these Listings. Combined with earlier CLEO results, the decay constant  $f_{D_s}$  is  $274 \pm 10 \pm 5$  MeV.

**$\Gamma(\gamma e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                        |    |         |           |                         |
|------------------------|----|---------|-----------|-------------------------|
| $< 1.3 \times 10^{-4}$ | 90 | ABLIKIM | 19AD BES3 | for $E_\gamma > 10$ MeV |
|------------------------|----|---------|-----------|-------------------------|

**$\Gamma(K^+ K^- e^+ \nu_e)/\Gamma(K^+ K^- \pi^+)$   $\Gamma_{23}/\Gamma_{38}$**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |                     |           |                             |
|-----------------------------|---------------------|-----------|-----------------------------|
| $0.558 \pm 0.007 \pm 0.016$ | <sup>1</sup> AUBERT | 08AN BABR | $e^+ e^-$ at $\Upsilon(4S)$ |
|-----------------------------|---------------------|-----------|-----------------------------|

<sup>1</sup> This AUBERT 08AN ratio is only for the  $K^+ K^-$  mass in the range 1.01–to–1.03 GeV in the numerator and 1.0095–to–1.0295 GeV in the denominator.

**$\Gamma(\phi e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$**

See the end of the  $D_s^+$  Listings for measurements of  $D_s^+ \rightarrow \phi e^+ \nu_e$  form factors. Unseen decay modes of the  $\phi$  are included.


| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

**2.39 ± 0.16 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

|                          |     |         |           |                             |
|--------------------------|-----|---------|-----------|-----------------------------|
| $2.26 \pm 0.45 \pm 0.09$ | 26  | ABLIKIM | 18A BES3  | $e^+ e^-$ at 4.009 GeV      |
| $2.14 \pm 0.17 \pm 0.08$ | 207 | HIETALA | 15        | Uses CLEO data              |
| $2.61 \pm 0.03 \pm 0.17$ | 25k | AUBERT  | 08AN BABR | $e^+ e^-$ at $\Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                          |     |         |    |                     |
|--------------------------|-----|---------|----|---------------------|
| $2.36 \pm 0.23 \pm 0.13$ | 106 | ECKLUND | 09 | CLEO See HIETALA 15 |
| $2.29 \pm 0.37 \pm 0.11$ | 45  | YELTON  | 09 | CLEO See ECKLUND 09 |



### $\Gamma(\phi e^+ \nu_e) / \Gamma(\phi \pi^+)$

$\Gamma_{24} / \Gamma_{39}$

As noted in the comment column, most of these measurements use  $\phi \mu^+ \nu_\mu$  events in addition to or instead of  $\phi e^+ \nu_e$  events.

| VALUE   | EVTS | DOCUMENT ID | TECN     | COMMENT  |
|---|------|-------------|----------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |      |             |          |  |
| $0.540 \pm 0.033 \pm 0.048$   | 793  | LINK        | 02J FOCs | Uses $\phi \mu^+ \nu_\mu$                      |
| $0.54 \pm 0.05 \pm 0.04$  | 367  | BUTLER      | 94 CLE2  | Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$ |
| $0.58 \pm 0.17 \pm 0.07$  | 97   | FRABETTI    | 93G E687 | Uses $\phi \mu^+ \nu_\mu$                      |
| $0.57 \pm 0.15 \pm 0.15$  | 104  | ALBRECHT    | 91 ARG   | Uses $\phi e^+ \nu_e$                          |
| $0.49 \pm 0.10 \begin{smallmatrix} +0.10 \\ -0.14 \end{smallmatrix}$          | 54   | ALEXANDER   | 90B CLEO | Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$ |

### $\Gamma(\phi \mu^+ \nu_\mu) / \Gamma_{\text{total}}$

$\Gamma_{25} / \Gamma$

| VALUE (%)                                  | EVTS | DOCUMENT ID | TECN     | COMMENT                |
|--|------|-------------|----------|------------------------|
| <b><math>1.94 \pm 0.53 \pm 0.09</math></b> | 22   | ABLIKIM     | 18A BES3 | $e^+ e^-$ at 4.009 GeV |

### $\Gamma(\eta e^+ \nu_e) / \Gamma_{\text{total}}$

$\Gamma_{27} / \Gamma$

Unseen decay modes of the  $\eta$  are included.

| VALUE (%)   | EVTS | DOCUMENT ID          | TECN     | COMMENT                |
|---|------|----------------------|----------|------------------------|
| <b><math>2.32 \pm 0.08</math> OUR AVERAGE</b>                                 |      |                      |          |                        |
| $2.323 \pm 0.063 \pm 0.063$   | 1.8k | ABLIKIM              | 19S BES3 | $e^+ e^-$ at 4178 MeV  |
| $2.30 \pm 0.31 \pm 0.08$  | 63   | ABLIKIM              | 16T BES3 | $e^+ e^-$ at 4.009 GeV |
| $2.28 \pm 0.14 \pm 0.19$  | 358  | <sup>1</sup> HIETALA | 15       | Uses CLEO data         |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |      |                      |          |                        |
| $2.48 \pm 0.29 \pm 0.13$  | 82   | YELTON               | 09 CLEO  | See HIETALA 15         |

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\eta e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$   $\Gamma_{27}/\Gamma_{24}$

Unseen decay modes of the  $\eta$  and the  $\phi$  are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.24 ± 0.12 ± 0.15      440      <sup>1</sup> BRANDENB... 95      CLE2      See HIETALA 15

<sup>1</sup> BRANDENBURG 95 uses both  $e^+$  and  $\mu^+$  events and makes a phase-space adjustment to use the  $\mu^+$  events as  $e^+$  events.

$\Gamma(\eta'(958) e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$

Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

**0.80 ± 0.07 OUR AVERAGE**

0.824 ± 0.073 ± 0.027      261      ABLIKIM      19S      BES3       $e^+ e^-$  at 4178 MeV

0.93 ± 0.30 ± 0.05      14      ABLIKIM      16T      BES3       $e^+ e^-$  at 4009 MeV

0.68 ± 0.15 ± 0.06      20      <sup>1</sup> HIETALA      15      Uses CLEO data

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.91 ± 0.33 ± 0.05      7.5      YELTON      09      CLEO      See HIETALA 15

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\eta'(958) e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$   $\Gamma_{28}/\Gamma_{24}$

Unseen decay modes of the resonances are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.43 ± 0.11 ± 0.07      29      <sup>1</sup> BRANDENB... 95      CLE2      See HIETALA 15

<sup>1</sup> BRANDENBURG 95 uses both  $e^+$  and  $\mu^+$  events and makes a phase-space adjustment to use the  $\mu^+$  events as  $e^+$  events.

$[\Gamma(\eta e^+ \nu_e) + \Gamma(\eta'(958) e^+ \nu_e)]/\Gamma(\phi e^+ \nu_e)$   $\Gamma_{26}/\Gamma_{24} = (\Gamma_{27} + \Gamma_{28})/\Gamma_{24}$

Unseen decay modes of the resonances are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.67 ± 0.17 ± 0.17      <sup>1</sup> BRANDENB... 95      CLE2      See HIETALA 15

<sup>1</sup> This BRANDENBURG 95 data is redundant with data in previous blocks.

$\Gamma(\eta \mu^+ \nu_\mu)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

**2.42 ± 0.46 ± 0.11**      44      ABLIKIM      18A      BES3       $e^+ e^-$  at 4.009 GeV

$\Gamma(\eta'(958) \mu^+ \nu_\mu)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

**1.06 ± 0.54 ± 0.07**      10      ABLIKIM      18A      BES3       $e^+ e^-$  at 4.009 GeV

$\Gamma(\omega e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

A test for  $u\bar{u}$  or  $d\bar{d}$  content in the  $D_s^+$ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and  $\omega - \phi$  mixing is an unlikely explanation for any fraction above about  $2 \times 10^{-4}$ .

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------|-----|-------------|------|---------|
|-----------|-----|-------------|------|---------|

**<0.20**      90      MARTIN      11      CLEO       $e^+ e^-$  at 4170 MeV

$\Gamma(K^0 e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

| VALUE (%)  | EVTS               | DOCUMENT ID              | TECN | COMMENT               |
|--|--------------------|--------------------------|------|-----------------------|
| <b>0.34 ± 0.04</b>   | <b>OUR AVERAGE</b> |                          |      |                       |
| 0.325 ± 0.038 ± 0.016  | 117                | <sup>1</sup> ABLIKIM 19D | BES3 | $e^+ e^-$ at 4178 MeV |
| 0.39 ± 0.08 ± 0.03   | 42                 | HIETALA 15               |      | Uses CLEO data        |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                |                    |                          |      |                       |
| 0.37 ± 0.10 ± 0.02   | 14                 | YELTON 09                | CLEO | See HIETALA 15        |
| <sup>1</sup> $K^0$ reconstructed via $K^0 \rightarrow K_S^0 \rightarrow \pi^+ \pi^-$ decays. |                    |                          |      |                       |

$\Gamma(K^*(892)^0 e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

Unseen decay modes of the  $K^*(892)^0$  are included.

| VALUE (%)   | EVTS               | DOCUMENT ID                         | TECN | COMMENT                |
|---|--------------------|-------------------------------------|------|------------------------|
| <b>0.215 ± 0.028</b>  | <b>OUR AVERAGE</b> | Error includes scale factor of 1.1. |      |                        |
| 0.237 ± 0.026 ± 0.020   | 155                | ABLIKIM 19D                         | BES3 | $e^+ e^-$ at 4178 MeV  |
| 0.18 ± 0.04 ± 0.01  | 32                 | <sup>1</sup> HIETALA 15             |      | $e^+ e^-$ at 4.170 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |                                     |      |                        |
| 0.18 ± 0.07 ± 0.01  | 7.5                | YELTON 09                           | CLEO | See HIETALA 15         |
| <sup>1</sup> Uses CLEO data, but not authored by the CLEO collaboration       |                    |                                     |      |                        |

$\Gamma(f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

| VALUE (%)  | EVTS | DOCUMENT ID             | TECN | COMMENT        |
|--|------|-------------------------|------|----------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • •  |      |                         |      |                |
| 0.13 ± 0.03 ± 0.01   | 42   | <sup>1</sup> HIETALA 15 |      | Uses CLEO data |
| 0.20 ± 0.03 ± 0.01   | 44   | ECKLUND 09              | CLEO | See HIETALA 15 |
| 0.13 ± 0.04 ± 0.01   | 13   | YELTON 09               | CLEO | See ECKLUND 09 |
| <sup>1</sup> HIETALA 15 uses a tighter cut on the reconstructed $\pi^+ \pi^-$ mass ( $\pm 60$ MeV around the $f^0$ ) than ECKLUND 09. It finds that applying the same tight cut to both analyses gives consistent results. |      |                         |      |                |

————— Hadronic modes with a  $K\bar{K}$  pair —————

$\Gamma(K^+ K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

| VALUE (%)  | EVTS               | DOCUMENT ID                         | TECN | COMMENT               |
|--|--------------------|-------------------------------------|------|-----------------------|
| <b>1.46 ± 0.04</b>   | <b>OUR FIT</b>     | Error includes scale factor of 1.1. |      |                       |
| <b>1.46 ± 0.05</b>   | <b>OUR AVERAGE</b> | Error includes scale factor of 1.2. |      |                       |
| 1.425 ± 0.038 ± 0.031  | 1.8k               | ABLIKIM 19A                         | BES3 | $e^+ e^-$ at 4178 MeV |
| 1.52 ± 0.05 ± 0.03   |                    | ONYISI 13                           | CLEO | $e^+ e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • •      |                    |                                     |      |                       |
| 1.49 ± 0.07 ± 0.05   |                    | <sup>1</sup> ALEXANDER 08           | CLEO | See ONYISI 13         |
| <sup>1</sup> ALEXANDER 08 uses single- and double-tagged events in an overall fit. |                    |                                     |      |                       |

$\Gamma(K^+ K_S^0)/\Gamma(K^+ K^- \pi^+)$   $\Gamma_{35}/\Gamma_{38}$

| VALUE (units $10^{-2}$ )   | EVTS | DOCUMENT ID | TECN | COMMENT                     |
|----------------------------|------|-------------|------|-----------------------------|
| <b>27.55 ± 0.18 ± 0.50</b> | 40k  | ABLIKIM 20R | BES3 | $e^+ e^-$ , 4178 ~ 4226 MeV |

$\Gamma(K^+ K_L^0)/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$

| VALUE (%)                    | EVTS | DOCUMENT ID | TECN | COMMENT               |
|------------------------------|------|-------------|------|-----------------------|
| <b>1.485 ± 0.039 ± 0.046</b> | 2.3k | ABLIKIM 19A | BES3 | $e^+ e^-$ at 4178 MeV |

$\Gamma(K^+ \bar{K}^0)/\Gamma_{\text{total}}$

$\Gamma_{37}/\Gamma$

| VALUE (%)             | EVTS | DOCUMENT ID         | TECN | COMMENT                                       |
|-----------------------|------|---------------------|------|---|
| <b>2.95±0.11±0.09</b> | 2.0k | <sup>1</sup> ZUPANC | 13   | BELL $e^+e^-$ at $\Upsilon(4S), \Upsilon(5S)$ |

<sup>1</sup> ZUPANC 13 finds the  $\bar{K}^0$  from its missing-mass squared, not from  $K_S^0 \rightarrow \pi^+ \pi^-$ .

The DCS ( $D_s^+ \rightarrow K^+ K^0$ ) contribution to this fraction is estimated to be an order of magnitude below the statistical uncertainty.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$

$\Gamma_{38}/\Gamma$

| VALUE (%)                    | EVTS  | DOCUMENT ID | TECN | COMMENT |
|------------------------------|---|-------------|------|---------|
| <b>5.39±0.15 OUR FIT</b>     | Error includes scale factor of 1.2.                         |             |      |         |
| <b>5.44±0.18 OUR AVERAGE</b> | Error includes scale factor of 1.3. See the ideogram below. |             |      |         |

5.55±0.14±0.13

ONYISI 13 CLEO  $e^+e^-$  at 4.17 GeV

5.06±0.15±0.21

4.1k ZUPANC 13 BELL  $e^+e^-$  at  $\Upsilon(4S), \Upsilon(5S)$

5.78±0.20±0.30

DEL-AMO-SA..10J BABR  $e^+e^-$ , 10.58 GeV


• • • We do not use the following data for averages, fits, limits, etc. • • •

5.50±0.23±0.16

<sup>1</sup> ALEXANDER 08 CLEO See ONYISI 13

<sup>1</sup> ALEXANDER 08 uses single- and double-tagged events in an overall fit.

WEIGHTED AVERAGE  
5.44±0.18 (Error scaled by 1.3)



$\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$

$\Gamma_{39}/\Gamma$

The results here are model-independent. For earlier, model-dependent results, see our PDG 06 edition. We decouple the  $D_s^+ \rightarrow \phi\pi^+$  branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the  $D_s^+ \rightarrow \phi\pi^+, \phi \rightarrow K^+ K^-$  branching fraction obtained from the Dalitz-plot analysis



of  $D_s^+ \rightarrow K^+ K^- \pi^+$ . That is, the ratio of these two branching fractions is not exactly the  $\phi \rightarrow K^+ K^-$  branching fraction 0.491.

| VALUE (%)   | EVTS     | DOCUMENT ID         | TECN     | COMMENT                        |
|---|----------|---------------------|----------|--------------------------------|
| <b>4.5 ± 0.4 OUR AVERAGE</b>  |          |                     |          |                                |
| 4.62 ± 0.36 ± 0.51  |          | <sup>1</sup> AUBERT | 06N BABR | $e^+ e^-$ at $\Upsilon(4S)$    |
| 4.81 ± 0.52 ± 0.38  | 212 ± 19 | <sup>2</sup> AUBERT | 05V BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| 3.59 ± 0.77 ± 0.48  |          | <sup>3</sup> ARTUSO | 96 CLE2  | $e^+ e^-$ at $\Upsilon(4S)$    |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |          |                     |          |                                |
| 3.9 <sup>+5.1</sup> <sub>-1.9</sub> <sup>+1.8</sup> <sub>-1.1</sub>           |          | <sup>4</sup> BAI    | 95C BES  | $e^+ e^-$ 4.03 GeV             |

<sup>1</sup> This AUBERT 06N measurement uses  $\bar{B}^0 \rightarrow D_s^{(*)-} D^{(*)+}$  and  $B^- \rightarrow D_s^{(*)-} D^{(*)0}$  decays, including some from other papers. However, the result is independent of AUBERT 05V.

<sup>2</sup> AUBERT 05V uses the ratio of  $B^0 \rightarrow D^{*-} D_s^{*+}$  events seen in two different ways, in both of which the  $D^{*-} \rightarrow \bar{D}^0 \pi^-$  decay is fully reconstructed: (1) The  $D_s^{*+} \rightarrow D_s^+ \gamma$ ,  $D_s^+ \rightarrow \phi \pi^+$  decay is fully reconstructed. (2) The number of events in the  $D_s^+$  peak in the missing mass spectrum against the  $D^{*-} \gamma$  is measured.

<sup>3</sup> ARTUSO 96 uses partially reconstructed  $\bar{B}^0 \rightarrow D^{*+} D_s^{*-}$  decays to get a model-independent value for  $\Gamma(D_s^- \rightarrow \phi \pi^-) / \Gamma(D^0 \rightarrow K^- \pi^+)$  of  $0.92 \pm 0.20 \pm 0.11$ .

<sup>4</sup> BAI 95C uses  $e^+ e^- \rightarrow D_s^+ D_s^-$  events in which one or both of the  $D_s^\pm$  are observed to obtain the first model-independent measurement of the  $D_s^+ \rightarrow \phi \pi^+$  branching fraction, without assumptions about  $\sigma(D_s^\pm)$ . However, with only two “doubly-tagged” events, the statistical error is very large.

### $\Gamma(\phi \pi^+, \phi \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{40} / \Gamma_{38}$

This is the “fit fraction” from the Dalitz-plot analysis. We decouple the  $D_s^+ \rightarrow \phi \pi^+$  branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the  $D_s^+ \rightarrow \phi \pi^+$ ,  $\phi \rightarrow K^+ K^-$  branching fraction obtained from the Dalitz-plot analysis of  $D_s^+ \rightarrow K^+ K^- \pi^+$ . That is, the ratio of these two branching fractions is not exactly the  $\phi \rightarrow K^+ K^-$  branching fraction 0.491.

| VALUE (units $10^{-2}$ )  | DOCUMENT ID     | TECN     | COMMENT                    |
|---|-----------------|----------|----------------------------|
| <b>41.6 ± 0.8 OUR AVERAGE</b>   |                 |          |                            |
| 41.4 ± 0.8 ± 0.5  | DEL-AMO-SA..11G | BABR     | Dalitz fit, 96k ± 369 evts |
| 42.2 ± 1.6 ± 0.3  | MITCHELL        | 09A CLEO | Dalitz fit, 12k evts       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                 |          |                            |
| 39.6 ± 3.3 ± 4.7  | FRABETTI        | 95B E687 | Dalitz fit, 701 evts       |

### $\Gamma(K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{41} / \Gamma_{38}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE (units $10^{-2}$ )  | DOCUMENT ID     | TECN     | COMMENT                    |
|---|-----------------|----------|----------------------------|
| <b>47.8 ± 0.6 OUR AVERAGE</b>   |                 |          |                            |
| 47.9 ± 0.5 ± 0.5  | DEL-AMO-SA..11G | BABR     | Dalitz fit, 96k ± 369 evts |
| 47.4 ± 1.5 ± 0.4  | MITCHELL        | 09A CLEO | Dalitz fit, 12k evts       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                 |          |                            |
| 47.8 ± 4.6 ± 4.0  | FRABETTI        | 95B E687 | Dalitz fit, 701 evts       |

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{42}/\Gamma_{38}$

This is the "fit fraction" from the Dalitz-plot analysis.

| <u>VALUE (units <math>10^{-2}</math>)</u>                                     | <u>DOCUMENT ID</u>                  | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------------------------------|-------------|----------------------------|
| <b>21 ± 6 OUR AVERAGE</b>   | Error includes scale factor of 3.5. |             |                            |
| 16.4 ± 0.7 ± 2.0  | DEL-AMO-SA..11G                     | BABR        | Dalitz fit, 96k ± 369 evts |
| 28.2 ± 1.9 ± 1.8  | MITCHELL 09A                        | CLEO        | Dalitz fit, 12k evts       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |             |                            |
| 11.0 ± 3.5 ± 2.6  | FRABETTI 95B                        | E687        | Dalitz fit, 701 evts       |

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{43}/\Gamma_{38}$

This is the "fit fraction" from the Dalitz-plot analysis.

| <u>VALUE (units <math>10^{-2}</math>)</u> | <u>DOCUMENT ID</u>                  | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------------------------------|-------------|----------------------------|
| <b>1.3 ± 0.8 OUR AVERAGE</b>              | Error includes scale factor of 3.9. |             |                            |
| 1.1 ± 0.1 ± 0.2                           | DEL-AMO-SA..11G                     | BABR        | Dalitz fit, 96k ± 369 evts |
| 4.3 ± 0.6 ± 0.5                           | MITCHELL 09A                        | CLEO        | Dalitz fit, 12k evts       |

$\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{44}/\Gamma_{38}$

This is the "fit fraction" from the Dalitz-plot analysis.

| <u>VALUE (units <math>10^{-2}</math>)</u>                                     | <u>DOCUMENT ID</u>                  | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------------------------------|-------------|----------------------------|
| <b>1.2 ± 0.5 OUR AVERAGE</b>  | Error includes scale factor of 3.8. |             |                            |
| 1.1 ± 0.1 ± 0.1   | DEL-AMO-SA..11G                     | BABR        | Dalitz fit, 96k ± 369 evts |
| 3.4 ± 0.5 ± 0.3   | MITCHELL 09A                        | CLEO        | Dalitz fit, 12k evts       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |             |                            |
| 3.4 ± 2.3 ± 3.5   | FRABETTI 95B                        | E687        | Dalitz fit, 701 evts       |

$\Gamma(K^+\bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^-\pi^+)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{45}/\Gamma_{38}$

This is the "fit fraction" from the Dalitz-plot analysis.

| <u>VALUE (units <math>10^{-2}</math>)</u>                                     | <u>DOCUMENT ID</u>                  | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------------------------------|-------------|----------------------------|
| <b>3.4 ± 0.7 OUR AVERAGE</b>  | Error includes scale factor of 1.2. |             |                            |
| 2.4 ± 0.3 ± 1.0   | DEL-AMO-SA..11G                     | BABR        | Dalitz fit, 96k ± 369 evts |
| 3.9 ± 0.5 ± 0.5   | MITCHELL 09A                        | CLEO        | Dalitz fit, 12k evts       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |             |                            |
| 9.3 ± 3.2 ± 3.2   | FRABETTI 95B                        | E687        | Dalitz fit, 701 evts       |

$\Gamma(K^+K_S^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$

| <u>VALUE (%)</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|---------------------------|--------------------|-------------|----------------------|
| <b>1.52 ± 0.09 ± 0.20</b> | ONYISI 13          | CLEO        | $e^+e^-$ at 4.17 GeV |

$\Gamma(2K_S^0\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$

| <u>VALUE (%)</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>       |
|---------------------------|--------------------|-------------|----------------------|
| <b>0.77 ± 0.05 ± 0.03</b> | ONYISI 13          | CLEO        | $e^+e^-$ at 4.17 GeV |

$\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(\phi\pi^+)$   $\Gamma_{49}/\Gamma_{39}$

Unseen decay modes of the resonances are included.

| <u>VALUE</u>              | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|---------------------------|--------------------|-------------|-----------------|
| <b>1.20 ± 0.21 ± 0.13</b> | CHEN 89            | CLEO        | $e^+e^-$ 10 GeV |

$\Gamma(K^+K^-\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$

| <u>VALUE (%)</u>          | <u>DOCUMENT ID</u>                  | <u>TECN</u> | <u>COMMENT</u>       |
|---------------------------|-------------------------------------|-------------|----------------------|
| <b>6.2 ± 0.6 OUR FIT</b>  | Error includes scale factor of 1.1. |             |                      |
| <b>6.37 ± 0.21 ± 0.56</b> | ONYISI 13                           | CLEO        | $e^+e^-$ at 4.17 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.65 \pm 0.29 \pm 0.40$  <sup>1</sup> ALEXANDER 08 CLEO See ONYISI 13

<sup>1</sup> ALEXANDER 08 uses single- and double-tagged events in an overall fit.

| $\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$ |             |                    |             |                | $\Gamma_{51}/\Gamma_{39}$ |
|--|-------------|--------------------|-------------|----------------|---------------------------|
| <u>VALUE</u>                           | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |                           |
| $1.86 \pm 0.26^{+0.29}_{-0.40}$        | 253         | AVERY              | 92          | CLE2           | $e^+e^- \simeq 10.5$ GeV  |

| $\Gamma(K_S^0 K^- 2\pi^+)/\Gamma_{\text{total}}$ |  |                    |             |                | $\Gamma_{52}/\Gamma$ |
|--|--|--------------------|-------------|----------------|----------------------|
| <u>VALUE (%)</u>                                 |  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |                      |
| <b><math>1.65 \pm 0.10</math> OUR FIT</b>        |  |                    |             |                |                      |
| <b><math>1.69 \pm 0.07 \pm 0.08</math></b>       |  | ONYISI             | 13          | CLEO           | $e^+e^-$ at 4.17 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.64 \pm 0.10 \pm 0.07$  <sup>1</sup> ALEXANDER 08 CLEO See ONYISI 13

<sup>1</sup> ALEXANDER 08 uses single- and double-tagged events in an overall fit.

| $\Gamma(K^*(892)^+ \bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$ |  |                    |             |                | $\Gamma_{53}/\Gamma_{39}$ |
|---|--|--------------------|-------------|----------------|---------------------------|
| <u>VALUE</u>  |  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |                           |
| Unseen decay modes of the resonances are included.      |  |                    |             |                |                           |
| <b><math>1.6 \pm 0.4 \pm 0.4</math></b>                 |  | ALBRECHT           | 92B         | ARG            | $e^+e^- \simeq 10.4$ GeV  |

| $\Gamma(K^+ K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ |  |                    |             |                | $\Gamma_{54}/\Gamma$ |
|---|--|--------------------|-------------|----------------|----------------------|
| <u>VALUE (%)</u>                                      |  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |                      |
| <b><math>0.99 \pm 0.08</math> OUR FIT</b>             |  |                    |             |                |                      |
| <b><math>1.03 \pm 0.06 \pm 0.08</math></b>            |  | ONYISI             | 13          | CLEO           | $e^+e^-$ at 4.17 GeV |

| $\Gamma(K^+ K_S^0 \pi^+ \pi^-)/\Gamma(K_S^0 K^- 2\pi^+)$ |             |                    |             |                | $\Gamma_{54}/\Gamma_{52}$                  |
|--|-------------|--------------------|-------------|----------------|--|
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |  |
| <b><math>0.60 \pm 0.05</math> OUR FIT</b>                |             |                    |             |                |  |
| <b><math>0.586 \pm 0.052 \pm 0.043</math></b>            | 476         | LINK               | 01C         | FOCS           | $\gamma A, \bar{E}_\gamma \approx 180$ GeV |

| $\Gamma(K^+ K^- 2\pi^+ \pi^-)/\Gamma(K^+ K^- \pi^+)$ |             |                    |             |                | $\Gamma_{55}/\Gamma_{38}$                   |
|--|-------------|--------------------|-------------|----------------|---|
| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |   |
| <b><math>0.160 \pm 0.027</math> OUR AVERAGE</b>      |             |                    |             |                |   |
| $0.150 \pm 0.019 \pm 0.025$                          | 240         | LINK               | 03D         | FOCS           | $\gamma A, \bar{E}_\gamma \approx 180$ GeV  |
| $0.188 \pm 0.036 \pm 0.040$                          | 75          | FRABETTI           | 97C         | E687           | $\gamma Be, \bar{E}_\gamma \approx 200$ GeV |

| $\Gamma(\phi 2\pi^+ \pi^-)/\Gamma(\phi\pi^+)$   |             |                    |             |                | $\Gamma_{56}/\Gamma_{39}$                   |
|---|-------------|--------------------|-------------|----------------|---|
| <u>VALUE</u>                                    | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |   |
| <b><math>0.269 \pm 0.027</math> OUR AVERAGE</b> |             |                    |             |                |   |
| $0.249 \pm 0.024 \pm 0.021$                     | 136         | LINK               | 03D         | FOCS           | $\gamma A, \bar{E}_\gamma \approx 180$ GeV  |
| $0.28 \pm 0.06 \pm 0.01$                        | 40          | FRABETTI           | 97C         | E687           | $\gamma Be, \bar{E}_\gamma \approx 200$ GeV |
| $0.58 \pm 0.21 \pm 0.10$                        | 21          | FRABETTI           | 92          | E687           | $\gamma Be$                                 |
| $0.42 \pm 0.13 \pm 0.07$                        | 19          | ANJOS              | 88          | E691           | Photoproduction                             |
| $1.11 \pm 0.37 \pm 0.28$                        | 62          | ALBRECHT           | 85D         | ARG            | $e^+e^- 10$ GeV                             |

| $\Gamma(K^+ K^- \rho^0 \pi^+ \text{non-}\phi)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ |            |                    |             |                | $\Gamma_{60}/\Gamma_{55}$                  |
|---|------------|--------------------|-------------|----------------|--|
| <u>VALUE</u>  | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |  |
| <b><math>&lt; 0.03</math></b>   | 90         | LINK               | 03D         | FOCS           | $\gamma A, \bar{E}_\gamma \approx 180$ GeV |

| $\Gamma(\phi\rho^0\pi^+, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-2\pi^+\pi^-)$ |             | $\Gamma_{57}/\Gamma_{55}$ |  |
|--|-------------|---------------------------|--|
| VALUE  | DOCUMENT ID | TECN                      | COMMENT                                      |
| <b>0.75±0.06±0.04</b>  | LINK        | 03D FOCS                  | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

| $\Gamma(\phi a_1(1260)^+, \phi \rightarrow K^+K^-, a_1^+ \rightarrow \rho^0\pi^+)/\Gamma(K^+K^-\pi^+)$ |             | $\Gamma_{58}/\Gamma_{38}$ |  |
|--|-------------|---------------------------|--|
| VALUE  | DOCUMENT ID | TECN                      | COMMENT                                      |
| <b>0.137±0.019±0.011</b>   | LINK        | 03D FOCS                  | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

| $\Gamma(K^+K^-2\pi^+\pi^- \text{ nonresonant})/\Gamma(K^+K^-2\pi^+\pi^-)$ |             | $\Gamma_{61}/\Gamma_{55}$ |  |
|---|-------------|---------------------------|--|
| VALUE   | DOCUMENT ID | TECN                      | COMMENT                                      |
| <b>0.10±0.06±0.05</b>   | LINK        | 03D FOCS                  | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

| $\Gamma(\phi 2\pi^+\pi^- \text{ non-}\rho, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-2\pi^+\pi^-)$ |             | $\Gamma_{59}/\Gamma_{55}$ |  |
|--|-------------|---------------------------|--|
| VALUE  | DOCUMENT ID | TECN                      | COMMENT                                      |
| <b>0.21±0.05±0.06</b>  | LINK        | 03D FOCS                  | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

| $\Gamma(2K_S^0 2\pi^+\pi^-)/\Gamma(K_S^0 K^- 2\pi^+)$ |         | $\Gamma_{62}/\Gamma_{52}$ |          |  |
|---|---------|---------------------------|----------|--|
| VALUE   | EVTS    | DOCUMENT ID               | TECN     | COMMENT                                      |
| <b>0.051±0.015±0.015</b>                              | 37 ± 10 | LINK                      | 04D FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

————— Pionic modes —————

| $\Gamma(\pi^+\pi^0)/\Gamma(K^+K_S^0)$   |     | $\Gamma_{63}/\Gamma_{35}$ |      |                           |
|---|-----|---------------------------|------|---------------------------|
| VALUE (units $10^{-2}$ )  | CL% | DOCUMENT ID               | TECN | COMMENT                   |
| <b>&lt;2.3</b>  | 90  | MENDEZ                    | 10   | CLEO $e^+e^-$ at 4170 MeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                           |      |                           |
| <4.1  | 90  | ADAMS                     | 07A  | CLEO See MENDEZ 10        |

| $\Gamma(2\pi^+\pi^-)/\Gamma_{\text{total}}$  |                                     | $\Gamma_{64}/\Gamma$ |                           |
|--|-------------------------------------|----------------------|---------------------------|
| VALUE (%)  | DOCUMENT ID                         | TECN                 | COMMENT                   |
| <b>1.08±0.04 OUR FIT</b>   | Error includes scale factor of 1.1. |                      |                           |
| <b>1.11±0.04±0.04</b>  | ONYISI                              | 13                   | CLEO $e^+e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • •      |                                     |                      |                           |
| 1.11±0.07±0.04   | <sup>1</sup> ALEXANDER              | 08                   | CLEO See ONYISI 13        |
| <sup>1</sup> ALEXANDER 08 uses single- and double-tagged events in an overall fit. |                                     |                      |                           |

| $\Gamma(2\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$                                     |                 | $\Gamma_{64}/\Gamma_{38}$ |          |                               |
|---|-----------------|---------------------------|----------|-------------------------------|
| VALUE   | EVTS            | DOCUMENT ID               | TECN     | COMMENT                       |
| <b>0.201±0.007 OUR FIT</b>  |                 |                           |          |                               |
| <b>0.199±0.004±0.009</b>  | $\approx 10.5k$ | AUBERT                    | 09O BABR | $e^+e^- \approx 10.6$ GeV     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                 |                           |          |                               |
| 0.265±0.041±0.031   | 98              | FRABETTI                  | 97D E687 | $\gamma$ Be $\approx 200$ GeV |

| $\Gamma(\rho^0\pi^+)/\Gamma(2\pi^+\pi^-)$                                     |     | $\Gamma_{65}/\Gamma_{64}$ |          |                                  |
|---|-----|---------------------------|----------|----------------------------------|
| VALUE   | CL% | DOCUMENT ID               | TECN     | COMMENT                          |
| <b>0.018±0.005±0.010</b>  |     | AUBERT                    | 09O BABR | Dalitz fit, $\approx 10.5k$ evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                           |          |                                  |
| not seen  |     | LINK                      | 04 FOCS  | Dalitz fit, $1475 \pm 50$ evts   |
| 0.058±0.023±0.037   |     | AITALA                    | 01A E791 | Dalitz fit, 848 evts             |
| <0.073  | 90  | FRABETTI                  | 97D E687 | $\gamma$ Be $\approx 200$ GeV    |

$\Gamma(\pi^+(\pi^+\pi^-)_{S\text{-wave}})/\Gamma(2\pi^+\pi^-)$   $\Gamma_{66}/\Gamma_{64}$

This is the “fit fraction” from the Dalitz-plot analysis. See also KLEMPY 08, which uses 568  $D_S^+ \rightarrow 3\pi$  decays (over 280 background events) from FNAL E791 to study various parametrizations of the decay amplitudes. The emphasis there is more on S-wave  $\pi\pi$  decay products — 20 different solutions are given — than on  $D_S^+$  fit fractions.

| VALUE                            | DOCUMENT ID         | TECN     | COMMENT                    |
|----------------------------------|---------------------|----------|----------------------------|
| <b>0.833 ± 0.020 OUR AVERAGE</b> |                     |          |                            |
| 0.830 ± 0.009 ± 0.019            | <sup>1</sup> AUBERT | 090 BABR | Dalitz fit, ≈ 10.5k evts   |
| 0.8704 ± 0.0560 ± 0.0438         | <sup>2</sup> LINK   | 04 FOCS  | Dalitz fit, 1475 ± 50 evts |

<sup>1</sup>AUBERT 090 gives the amplitude and phase of the  $\pi^+\pi^-$  S-wave in 29  $\pi^+\pi^-$  invariant-mass bins.

<sup>2</sup>LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full  $\pi-\pi$  S-wave isoscalar scattering amplitude to describe the  $\pi^+\pi^-$  S-wave component of the  $\pi^+\pi^+\pi^-$  state. The fit fraction given above is a sum over five  $f_0$  mesons, the  $f_0(980)$ ,  $f_0(1300)$ ,  $f_0(1200-1600)$ ,  $f_0(1500)$ , and  $f_0(1750)$ . See LINK 04 for details and discussion.

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{67}/\Gamma_{64}$

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full  $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$  fit fraction.

| VALUE   | DOCUMENT ID | TECN     | COMMENT               |
|---|-------------|----------|-----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |          |                       |
| 0.565 ± 0.043 ± 0.047   | AITALA      | 01A E791 | Dalitz fit, 848 evts  |
| 1.074 ± 0.140 ± 0.043   | FRABETTI    | 97D E687 | $\gamma$ Be ≈ 200 GeV |

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{68}/\Gamma_{64}$

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full  $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$  fit fraction.

| VALUE   | DOCUMENT ID | TECN     | COMMENT              |
|---|-------------|----------|----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |          |                      |
| 0.324 ± 0.077 ± 0.017   | AITALA      | 01A E791 | Dalitz fit, 848 evts |

$\Gamma(f_0(1500)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{69}/\Gamma_{64}$

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full  $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$  fit fraction.

| VALUE   | DOCUMENT ID           | TECN     | COMMENT               |
|---|-----------------------|----------|-----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |          |                       |
| 0.274 ± 0.114 ± 0.019   | <sup>1</sup> FRABETTI | 97D E687 | $\gamma$ Be ≈ 200 GeV |

<sup>1</sup>FRABETTI 97D calls this mode  $S(1475)\pi^+$ , but finds the mass and width of this  $S(1475)$  to be in excellent agreement with those of the  $f_0(1500)$ .

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{70}/\Gamma_{64}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE   | DOCUMENT ID | TECN     | COMMENT                    |
|---|-------------|----------|----------------------------|
| <b>0.101 ± 0.018 OUR AVERAGE</b>  |             |          |                            |
| 0.101 ± 0.015 ± 0.011   | AUBERT      | 090 BABR | Dalitz fit, ≈ 10.5k evts   |
| 0.0974 ± 0.0449 ± 0.0294  | LINK        | 04 FOCS  | Dalitz fit, 1475 ± 50 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |          |                            |
| 0.197 ± 0.033 ± 0.006   | AITALA      | 01A E791 | Dalitz fit, 848 evts       |
| 0.123 ± 0.056 ± 0.018   | FRABETTI    | 97D E687 | $\gamma$ Be ≈ 200 GeV      |

$\Gamma(\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-) / \Gamma(2\pi^+ \pi^-)$   $\Gamma_{71} / \Gamma_{64}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE   | DOCUMENT ID | TECN | COMMENT                         |
|---|-------------|------|---------------------------------|
| <b>0.027 ± 0.018 OUR AVERAGE</b>  |             |      |                                 |
| 0.023 ± 0.008 ± 0.017   | AUBERT      | 090  | BABR Dalitz fit, ≈ 10.5k evts   |
| 0.0656 ± 0.0343 ± 0.0440  | LINK        | 04   | FOCS Dalitz fit, 1475 ± 50 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |                                 |
| 0.044 ± 0.021 ± 0.002   | AITALA      | 01A  | E791 Dalitz fit, 848 evts       |

$\Gamma(\pi^+ 2\pi^0) / \Gamma_{\text{total}}$   $\Gamma_{72} / \Gamma$

| VALUE (%)                 | EVTS    | DOCUMENT ID | TECN | COMMENT                    |
|---------------------------|---------|-------------|------|----------------------------|
| <b>0.65 ± 0.13 ± 0.03</b> | 72 ± 16 | NAIK        | 09A  | CLEO $e^+ e^-$ at 4170 MeV |

$\Gamma(2\pi^+ \pi^- \pi^0) / \Gamma(\phi \pi^+)$   $\Gamma_{73} / \Gamma_{39}$

| VALUE   | CL% | DOCUMENT ID | TECN | COMMENT              |
|---|-----|-------------|------|----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |      |                      |
| < 3.3   | 90  | ANJOS       | 89E  | E691 Photoproduction |

$\Gamma(\eta \pi^+) / \Gamma_{\text{total}}$   $\Gamma_{74} / \Gamma$

Unseen decay modes of the  $\eta$  are included.

| VALUE (%)                      | EVTS | DOCUMENT ID | TECN | COMMENT                             |
|--------------------------------|------|-------------|------|-------------------------------------|
| <b>1.68 ± 0.10 OUR FIT</b>     |      |             |      | Error includes scale factor of 1.2. |
| <b>1.71 ± 0.08 OUR AVERAGE</b> |      |             |      |                                     |

|  |      |                        |    |  |
|--|------|------------------------|----|--|
| 1.67 ± 0.08 ± 0.06   |      | ONYISI                 | 13 | CLEO $e^+ e^-$ at 4.17 GeV                     |
| 1.82 ± 0.14 ± 0.07   | 0.8k | ZUPANC                 | 13 | BELL $e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • •      |      |                        |    |  |
| 1.58 ± 0.11 ± 0.18   |      | <sup>1</sup> ALEXANDER | 08 | CLEO See ONYISI 13                             |
| <sup>1</sup> ALEXANDER 08 uses single- and double-tagged events in an overall fit. |      |                        |    |  |

$\Gamma(\eta \pi^+) / \Gamma(K^+ K_S^0)$   $\Gamma_{74} / \Gamma_{35}$

Unseen decay modes of the  $\eta$  are included.

| VALUE   | EVTS      | DOCUMENT ID | TECN | COMMENT                             |
|---|-----------|-------------|------|-------------------------------------|
| <b>1.15 ± 0.08 OUR FIT</b>  |           |             |      | Error includes scale factor of 1.3. |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |           |             |      |                                     |
| 1.236 ± 0.043 ± 0.063   | 2587 ± 89 | MENDEZ      | 10   | CLEO See ONYISI 13                  |

$\Gamma(\eta \pi^+) / \Gamma(\phi \pi^+)$   $\Gamma_{74} / \Gamma_{39}$

Unseen decay modes of the resonances are included.

| VALUE   | EVTS | DOCUMENT ID | TECN | COMMENT                             |
|---|------|-------------|------|-------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |             |      |                                     |
| 0.48 ± 0.03 ± 0.04  | 920  | JESSOP      | 98   | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |
| 0.54 ± 0.09 ± 0.06  | 165  | ALEXANDER   | 92   | CLE2 See JESSOP 98                  |

$\Gamma(\eta \pi^+) / \Gamma(K^+ K^- \pi^+)$   $\Gamma_{74} / \Gamma_{38}$

| VALUE (units $10^{-2}$ )   | EVTS  | DOCUMENT ID | TECN | COMMENT                          |
|----------------------------|-------|-------------|------|----------------------------------|
| <b>31.94 ± 0.33 ± 0.49</b> | 19.5k | ABLIKIM     | 20R  | BES3 $e^+ e^-$ , 4178 ~ 4226 MeV |

$\Gamma(\omega \pi^+) / \Gamma_{\text{total}}$   $\Gamma_{75} / \Gamma$

Unseen decay modes of the  $\omega$  are included.

| VALUE (%)                        | EVTS    | DOCUMENT ID | TECN | COMMENT                     |
|----------------------------------|---------|-------------|------|-----------------------------|
| <b>0.192 ± 0.030 OUR FIT</b>     |         |             |      |                             |
| <b>0.181 ± 0.032 OUR AVERAGE</b> |         |             |      |                             |
| 0.177 ± 0.032 ± 0.013            | 65 ± 12 | ABLIKIM     | 19AH | BES3 $e^+ e^-$ at 4.178 GeV |

0.21 ± 0.09 ± 0.01 6 ± 2.4 GE 09A CLEO e<sup>+</sup>e<sup>-</sup> at 4170 MeV $\Gamma(\omega\pi^+)/\Gamma(\eta\pi^+)$   $\Gamma_{75}/\Gamma_{74}$ 

Unseen decay modes of the resonances are included.

| VALUE                        | EVTS | DOCUMENT ID | TECN | COMMENT   |
|------------------------------|------|-------------|------|---|
| <b>0.114 ± 0.018 OUR FIT</b> |      |             |      |   |
| <b>0.16 ± 0.04 ± 0.03</b>    |      | BALEST      | 97   | CLE2 e <sup>+</sup> e <sup>-</sup> ≈ $\Upsilon(4S)$ |

 $\Gamma(3\pi^+2\pi^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{76}/\Gamma_{38}$ 

| VALUE                            | EVTS | DOCUMENT ID | TECN | COMMENT  |
|----------------------------------|------|-------------|------|--|
| <b>0.146 ± 0.014 OUR AVERAGE</b> |      |             |      |  |
| 0.145 ± 0.011 ± 0.010            | 671  | LINK        | 03D  | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| 0.158 ± 0.042 ± 0.031            | 37   | FRABETTI    | 97C  | E687 $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |

 $\Gamma(\eta\rho^+)/\Gamma_{\text{total}}$   $\Gamma_{78}/\Gamma$ Unseen decay modes of the  $\eta$  are included.

| VALUE (%)              | EVTS     | DOCUMENT ID | TECN | COMMENT                         |
|------------------------|----------|-------------|------|---------------------------------|
| <b>8.9 ± 0.6 ± 0.5</b> | 328 ± 22 | NAIK        | 09A  | CLEO $\eta \rightarrow 2\gamma$ |

 $\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$   $\Gamma_{78}/\Gamma_{39}$ 

Unseen decay modes of the resonances are included.

| VALUE   | EVTS | DOCUMENT ID | TECN | COMMENT   |
|---|------|-------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |             |      |   |
| 2.98 ± 0.20 ± 0.39  | 447  | JESSOP      | 98   | CLE2 e <sup>+</sup> e <sup>-</sup> ≈ $\Upsilon(4S)$ |
| 2.86 ± 0.38 <sup>+0.36</sup> <sub>-0.38</sub>                                 | 217  | AVERY       | 92   | CLE2 See JESSOP 98                                  |

 $\Gamma(\eta\rho^+)/\Gamma(\eta\pi^+\pi^0)$   $\Gamma_{78}/\Gamma_{79}$ 

| VALUE (units 10 <sup>-2</sup> ) | EVTS | DOCUMENT ID | TECN | COMMENT                                  |
|---------------------------------|------|-------------|------|--|
| <b>78.3 ± 5.0 ± 2.1</b>         | 1.2k | ABLIKIM     | 19BE | BES3 $\eta\pi^+\pi^0$ amplitude analysis |

 $\Gamma(\eta\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{79}/\Gamma$ 

| VALUE (%)                    | EVTS | DOCUMENT ID | TECN | COMMENT   |
|------------------------------|------|-------------|------|---|
| <b>9.5 ± 0.5 OUR AVERAGE</b> |      |             |      |   |
| 9.50 ± 0.28 ± 0.41           | 2.6k | ABLIKIM     | 19BE | BES3 e <sup>+</sup> e <sup>-</sup> at 4.178 GeV |
| 9.2 ± 0.4 ± 1.1              |      | ONYISI      | 13   | CLEO e <sup>+</sup> e <sup>-</sup> at 4.17 GeV  |

 $\Gamma(\eta(\pi^+\pi^0)_{P\text{-wave}})/\Gamma(\eta\pi^+\pi^0)$   $\Gamma_{80}/\Gamma_{79}$ 

| VALUE (units 10 <sup>-2</sup> ) | EVTS | DOCUMENT ID | TECN | COMMENT                                  |
|---------------------------------|------|-------------|------|--|
| <b>5.4 ± 2.1 ± 2.5</b>          | 1.2k | ABLIKIM     | 19BE | BES3 $\eta\pi^+\pi^0$ amplitude analysis |

 $\Gamma(a_0(980)^+\pi^0, a_0(980)^+ \rightarrow \eta\pi^+\pi^0)/\Gamma(\eta\pi^+\pi^0)$   $\Gamma_{81}/\Gamma_{79}$ 

| VALUE (units 10 <sup>-2</sup> ) | EVTS | DOCUMENT ID          | TECN | COMMENT                                  |
|---------------------------------|------|----------------------|------|--|
| <b>23.2 ± 2.3 ± 3.3</b>         | 1.2k | <sup>1</sup> ABLIKIM | 19BE | BES3 $\eta\pi^+\pi^0$ amplitude analysis |

<sup>1</sup> Coherent sum of  $D_s^+ \rightarrow a_0^+\pi^0 \rightarrow \eta\pi^+\pi^0$  and  $D_s^+ \rightarrow a_0^0\pi^+ \rightarrow \eta\pi^+\pi^0$ . ABLIKIM 19BE find  $a_0(980)^0 - f(980)$  mixing effects negligibly small in this  $D_s^+ \rightarrow \eta\pi^+\pi^0$  Dalitz plot analysis.

$\Gamma(\omega\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{82}/\Gamma$

Unseen decay modes of the  $\omega$  are included.

| VALUE (%)                 | EVTS     | DOCUMENT ID | TECN     | COMMENT              |
|---------------------------|----------|-------------|----------|----------------------|
| <b>2.78 ± 0.65 ± 0.25</b> | 34 ± 7.9 | GE          | 09A CLEO | $e^+e^-$ at 4170 MeV |

$\Gamma(3\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{83}/\Gamma$

| VALUE  | DOCUMENT ID | TECN | COMMENT         |
|--|-------------|------|-----------------|
| <b>0.049<sup>+0.033</sup><sub>-0.030</sub></b> | BARLAG 92C  | ACCM | $\pi^-$ 230 GeV |

$\Gamma(\omega 2\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{84}/\Gamma$

Unseen decay modes of the  $\omega$  are included.

| VALUE (%)                 | EVTS     | DOCUMENT ID | TECN     | COMMENT              |
|---------------------------|----------|-------------|----------|----------------------|
| <b>1.58 ± 0.45 ± 0.09</b> | 29 ± 8.2 | GE          | 09A CLEO | $e^+e^-$ at 4170 MeV |

$\Gamma(\eta'(958)\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$

Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE (%)                 | DOCUMENT ID | TECN | COMMENT              |
|---------------------------|-------------|------|----------------------|
| <b>3.94 ± 0.15 ± 0.20</b> | ONYISI 13   | CLEO | $e^+e^-$ at 4.17 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                    |                           |      |               |
|--------------------|---------------------------|------|---------------|
| 3.77 ± 0.25 ± 0.30 | <sup>1</sup> ALEXANDER 08 | CLEO | See ONYISI 13 |
|--------------------|---------------------------|------|---------------|

<sup>1</sup>ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+K_S^0)$   $\Gamma_{85}/\Gamma_{35}$

Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE                        | EVTS      | DOCUMENT ID | TECN | COMMENT       |
|------------------------------|-----------|-------------|------|---------------|
| <b>2.654 ± 0.088 ± 0.139</b> | 1436 ± 47 | MENDEZ 10   | CLEO | See ONYISI 13 |

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$   $\Gamma_{85}/\Gamma_{39}$

Unseen decay modes of the resonances are included.

| VALUE  | EVTS | DOCUMENT ID  | TECN | COMMENT                       |
|--|------|--------------|------|-------------------------------|
| <b>1.03 ± 0.06 ± 0.07</b>                      | 537  | JESSOP 98    | CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| <b>1.20 ± 0.15 ± 0.11</b>                      | 281  | ALEXANDER 92 | CLE2 | See JESSOP 98                 |
| <b>2.5 ± 1.0<sup>+1.5</sup><sub>-0.4</sub></b> | 22   | ALVAREZ 91   | NA14 | Photoproduction               |
| <b>2.5 ± 0.5 ± 0.3</b>                         | 215  | ALBRECHT 90D | ARG  | $e^+e^- \approx 10.4$ GeV     |

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{85}/\Gamma_{38}$

| VALUE (units 10 <sup>-2</sup> ) | EVTS | DOCUMENT ID | TECN | COMMENT                    |
|---------------------------------|------|-------------|------|----------------------------|
| <b>69.4 ± 0.8 ± 3.8</b>         | 9.9k | ABLIKIM 20R | BES3 | $e^+e^-$ , 4178 ~ 4226 MeV |

$\Gamma(\omega\eta\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$

Unseen decay modes of the  $\omega$  and  $\eta$  are included.

| VALUE                              | CL% | DOCUMENT ID | TECN     | COMMENT              |
|------------------------------------|-----|-------------|----------|----------------------|
| <b>&lt; 2.13 × 10<sup>-2</sup></b> | 90  | GE          | 09A CLEO | $e^+e^-$ at 4170 MeV |



| $\Gamma(\eta'(958)\rho^+)/\Gamma_{\text{total}}$ |      |             |      | $\Gamma_{88}/\Gamma$                 |
|--|------|-------------|------|--------------------------------------|
| VALUE (%)  | EVTS | DOCUMENT ID | TECN | COMMENT                              |
| <b>5.8±1.4±0.4</b>                               |      | ABLIKIM     | 15Z  | BES3 482 pb <sup>-1</sup> , 4009 MeV |

| $\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$       |      |             |      | $\Gamma_{88}/\Gamma_{39}$ |
|--|------|-------------|------|---------------------------|
| VALUE  | EVTS | DOCUMENT ID | TECN | COMMENT                   |
| Unseen decay modes of the resonances are included. |      |             |      |                           |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|   |     |                     |    |   |
|---|-----|---------------------|----|---|
| 2.78±0.28±0.30                              | 137 | <sup>1</sup> JESSOP | 98 | CLE2 e <sup>+</sup> e <sup>-</sup> ≈ $\Upsilon(4S)$ |
| 3.44±0.62 <sup>+0.44</sup> <sub>-0.46</sub> | 68  | AVERY               | 92 | CLE2 See JESSOP 98                                  |

<sup>1</sup>This JESSOP 98 fraction, when combined with other  $\eta'$  fractions, greatly overshoots the inclusive  $\eta'$  fraction. See the measurement just above, which fits nicely.

| $\Gamma(\eta'(958)\pi^+\pi^0)/\Gamma_{\text{total}}$ |      |             |      | $\Gamma_{89}/\Gamma$                           |
|--|------|-------------|------|--|
| VALUE (%)  | EVTS | DOCUMENT ID | TECN | COMMENT  |
| <b>5.6±0.5±0.6</b>                                   |      | ONYISI      | 13   | CLEO e <sup>+</sup> e <sup>-</sup> at 4.17 GeV |

| $\Gamma(\eta'(958)\pi^+\pi^0\text{nonresonant})/\Gamma_{\text{total}}$ |     |             |      | $\Gamma_{90}/\Gamma$                 |
|--|-----|-------------|------|--------------------------------------|
| VALUE  | CL% | DOCUMENT ID | TECN | COMMENT                              |
| <b>&lt;5.1 × 10<sup>-2</sup></b>                                       | 90  | ABLIKIM     | 15Z  | BES3 482 pb <sup>-1</sup> , 4009 MeV |

### ———— Modes with one or three K's ————

| $\Gamma(K^+\pi^0)/\Gamma(K^+K_S^0)$   |          |             |      | $\Gamma_{91}/\Gamma_{35}$                      |
|---|----------|-------------|------|--|
| VALUE (units 10 <sup>-2</sup> )   | EVTS     | DOCUMENT ID | TECN | COMMENT  |
| <b>4.2±1.4±0.2</b>  | 202 ± 70 | MENDEZ      | 10   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |          |             |      |  |
| 5.5±1.3±0.7   | 141 ± 34 | ADAMS       | 07A  | CLEO See MENDEZ 10                             |

| $\Gamma(K^+\pi^0)/\Gamma(K^+K^-\pi^+)$ |      |             |      | $\Gamma_{91}/\Gamma_{38}$                            |
|--|------|-------------|------|--|
| VALUE (units 10 <sup>-3</sup> )        | EVTS | DOCUMENT ID | TECN | COMMENT  |
| <b>13.73±0.90±0.33</b>                 | 2.3k | ABLIKIM     | 20R  | BES3 e <sup>+</sup> e <sup>-</sup> , 4178 ~ 4226 MeV |

| $\Gamma(K_S^0\pi^+)/\Gamma(K^+K_S^0)$   |           |             |      | $\Gamma_{92}/\Gamma_{35}$                            |
|---|-----------|-------------|------|--|
| VALUE (units 10 <sup>-2</sup> )   | EVTS      | DOCUMENT ID | TECN | COMMENT  |
| <b>8.12±0.28 OUR AVERAGE</b>  |           |             |      |  |
| 8.5 ±0.7 ±0.2   | 393 ± 33  | MENDEZ      | 10   | CLEO e <sup>+</sup> e <sup>-</sup> at 4170 MeV       |
| 8.03±0.24±0.19  | 17.6k±481 | WON         | 09   | BELL e <sup>+</sup> e <sup>-</sup> at $\Upsilon(4S)$ |
| 10.4 ±2.4 ±1.4  | 113 ± 26  | LINK        | 08   | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV    |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |           |             |      |  |
| 8.2 ±0.9 ±0.2   | 206 ± 22  | ADAMS       | 07A  | CLEO See MENDEZ 10                                   |

| $\Gamma(K_S^0\pi^+)/\Gamma(K^+K^-\pi^+)$ |      |             |      | $\Gamma_{92}/\Gamma_{38}$                            |
|--|------|-------------|------|--|
| VALUE (units 10 <sup>-3</sup> )          | EVTS | DOCUMENT ID | TECN | COMMENT  |
| <b>20.35±0.62±0.42</b>                   | 2.7k | ABLIKIM     | 20R  | BES3 e <sup>+</sup> e <sup>-</sup> , 4178 ~ 4226 MeV |

$\Gamma(K^+\eta)/\Gamma(K^+K_S^0)$   $\Gamma_{93}/\Gamma_{35}$

Unseen decay modes of the  $\eta$  are included.

| VALUE (units $10^{-2}$ ) | EVTS     | DOCUMENT ID | TECN | COMMENT                   |
|--------------------------|----------|-------------|------|---------------------------|
| <b>11.8±2.2±0.6</b>      | 222 ± 41 | MENDEZ      | 10   | CLEO $e^+e^-$ at 4170 MeV |

$\Gamma(K^+\eta)/\Gamma(\eta\pi^+)$   $\Gamma_{93}/\Gamma_{74}$

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|             |          |       |     |                    |
|-------------|----------|-------|-----|--------------------|
| 8.9±1.5±0.4 | 113 ± 18 | ADAMS | 07A | CLEO See MENDEZ 10 |
|-------------|----------|-------|-----|--------------------|

$\Gamma(K^+\eta)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{93}/\Gamma_{38}$

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

|                       |      |         |     |                                 |
|-----------------------|------|---------|-----|---------------------------------|
| <b>2.97±0.18±0.06</b> | 1.8k | ABLIKIM | 20R | BES3 $e^+e^-$ , 4178 ~ 4226 MeV |
|-----------------------|------|---------|-----|---------------------------------|

$\Gamma(K^+\omega)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$

Unseen decay modes of the  $\omega$  are included.

| VALUE (units $10^{-4}$ ) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|------|---------|
|--------------------------|-----|------|-------------|------|---------|

|                    |    |                      |      |      |                       |
|--------------------|----|----------------------|------|------|-----------------------|
| <b>8.7±2.4±0.8</b> | 29 | <sup>1</sup> ABLIKIM | 19AH | BES3 | $e^+e^-$ at 4.178 GeV |
|--------------------|----|----------------------|------|------|-----------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|     |    |    |     |      |                      |
|-----|----|----|-----|------|----------------------|
| <24 | 90 | GE | 09A | CLEO | $e^+e^-$ at 4170 MeV |
|-----|----|----|-----|------|----------------------|

<sup>1</sup>Evidence for mode at 4.4 $\sigma$ .

$\Gamma(K^+\eta'(958))/\Gamma(K^+K_S^0)$   $\Gamma_{95}/\Gamma_{35}$

Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

|                     |         |        |    |                           |
|---------------------|---------|--------|----|---------------------------|
| <b>11.8±3.6±0.7</b> | 56 ± 17 | MENDEZ | 10 | CLEO $e^+e^-$ at 4170 MeV |
|---------------------|---------|--------|----|---------------------------|

$\Gamma(K^+\eta'(958))/\Gamma(\eta'(958)\pi^+)$   $\Gamma_{95}/\Gamma_{85}$

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|             |        |       |     |                    |
|-------------|--------|-------|-----|--------------------|
| 4.2±1.3±0.3 | 28 ± 9 | ADAMS | 07A | CLEO See MENDEZ 10 |
|-------------|--------|-------|-----|--------------------|

$\Gamma(K^+\eta'(958))/\Gamma(K^+K^-\pi^+)$   $\Gamma_{95}/\Gamma_{38}$

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

|                       |     |         |     |                                 |
|-----------------------|-----|---------|-----|---------------------------------|
| <b>4.91±0.31±0.31</b> | 675 | ABLIKIM | 20R | BES3 $e^+e^-$ , 4178 ~ 4226 MeV |
|-----------------------|-----|---------|-----|---------------------------------|

$\Gamma(K^+\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

**0.65 ± 0.04 OUR FIT**

|                          |        |    |                           |
|--------------------------|--------|----|---------------------------|
| <b>0.654±0.033±0.025</b> | ONYISI | 13 | CLEO $e^+e^-$ at 4.17 GeV |
|--------------------------|--------|----|---------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                    |                        |    |                    |
|--------------------|------------------------|----|--------------------|
| 0.69 ± 0.05 ± 0.03 | <sup>1</sup> ALEXANDER | 08 | CLEO See ONYISI 13 |
|--------------------|------------------------|----|--------------------|

<sup>1</sup>ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(K^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{96}/\Gamma_{38}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

**0.121±0.008 OUR FIT** Error includes scale factor of 1.1.

|                          |          |      |     |   |
|--------------------------|----------|------|-----|---|
| <b>0.127±0.007±0.014</b> | 567 ± 31 | LINK | 04F | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |
|--------------------------|----------|------|-----|---|

$$\Gamma(K^+\rho^0)/\Gamma(K^+\pi^+\pi^-) \quad \Gamma_{97}/\Gamma_{96}$$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                       | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------|-------------|------|---------------------------|
| <b>0.3883±0.0531±0.0261</b> | LINK        | 04F  | FOCS Dalitz fit, 567 evts |

$$\Gamma(K^+\rho(1450)^0, \rho^0 \rightarrow \pi^+\pi^-)/\Gamma(K^+\pi^+\pi^-) \quad \Gamma_{98}/\Gamma_{96}$$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                       | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------|-------------|------|---------------------------|
| <b>0.1062±0.0351±0.0104</b> | LINK        | 04F  | FOCS Dalitz fit, 567 evts |

$$\Gamma(K^*(892)^0\pi^+, K^{*0} \rightarrow K^+\pi^-)/\Gamma(K^+\pi^+\pi^-) \quad \Gamma_{99}/\Gamma_{96}$$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                       | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------|-------------|------|---------------------------|
| <b>0.2164±0.0321±0.0114</b> | LINK        | 04F  | FOCS Dalitz fit, 567 evts |

$$\Gamma(K^*(1410)^0\pi^+, K^{*0} \rightarrow K^+\pi^-)/\Gamma(K^+\pi^+\pi^-) \quad \Gamma_{100}/\Gamma_{96}$$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                       | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------|-------------|------|---------------------------|
| <b>0.1882±0.0403±0.0122</b> | LINK        | 04F  | FOCS Dalitz fit, 567 evts |

$$\Gamma(K^*(1430)^0\pi^+, K^{*0} \rightarrow K^+\pi^-)/\Gamma(K^+\pi^+\pi^-) \quad \Gamma_{101}/\Gamma_{96}$$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                       | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------|-------------|------|---------------------------|
| <b>0.0765±0.0500±0.0170</b> | LINK        | 04F  | FOCS Dalitz fit, 567 evts |

$$\Gamma(K^+\pi^+\pi^- \text{ nonresonant})/\Gamma(K^+\pi^+\pi^-) \quad \Gamma_{102}/\Gamma_{96}$$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                       | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------|-------------|------|---------------------------|
| <b>0.1588±0.0492±0.0153</b> | LINK        | 04F  | FOCS Dalitz fit, 567 evts |

$$\Gamma(K^0\pi^+\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{103}/\Gamma$$

| VALUE (%)             | EVTS   | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------|--------|-------------|------|---------------------------|
| <b>1.00±0.18±0.04</b> | 44 ± 8 | NAIK        | 09A  | CLEO $e^+e^-$ at 4170 MeV |

$$\Gamma(K_S^0 2\pi^+\pi^-)/\Gamma(K_S^0 K^- 2\pi^+) \quad \Gamma_{104}/\Gamma_{52}$$

| VALUE                 | EVTS     | DOCUMENT ID | TECN | COMMENT   |
|-----------------------|----------|-------------|------|---|
| <b>0.18±0.04±0.05</b> | 179 ± 36 | LINK        | 08   | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$$\Gamma(K^+\omega\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{105}/\Gamma$$
Unseen decay modes of the  $\omega$  are included.

| VALUE (%)       | CL% | DOCUMENT ID | TECN | COMMENT                   |
|-----------------|-----|-------------|------|---------------------------|
| <b>&lt;0.82</b> | 90  | GE          | 09A  | CLEO $e^+e^-$ at 4170 MeV |

$$\Gamma(K^+\omega\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{106}/\Gamma$$
Unseen decay modes of the  $\omega$  are included.

| VALUE (%)       | CL% | DOCUMENT ID | TECN | COMMENT                   |
|-----------------|-----|-------------|------|---------------------------|
| <b>&lt;0.54</b> | 90  | GE          | 09A  | CLEO $e^+e^-$ at 4170 MeV |

$$\Gamma(K^+\omega\eta)/\Gamma_{\text{total}} \quad \Gamma_{107}/\Gamma$$
Unseen decay modes of the  $\omega$  and  $\eta$  are included.

| VALUE (%)       | CL% | DOCUMENT ID | TECN | COMMENT                   |
|-----------------|-----|-------------|------|---------------------------|
| <b>&lt;0.79</b> | 90  | GE          | 09A  | CLEO $e^+e^-$ at 4170 MeV |

$\Gamma(2K^+K^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{108}/\Gamma_{38}$

| VALUE (units $10^{-3}$ )  | EVTS         | DOCUMENT ID     | TECN | COMMENT                          |
|---|--------------|-----------------|------|----------------------------------|
| <b><math>4.0 \pm 0.3 \pm 0.2</math></b>                                       | $748 \pm 60$ | DEL-AMO-SA..11G | BABR | $e^+e^- \approx \Upsilon(4S)$    |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |              |                 |      |                                  |
| $8.95 \pm 2.12^{+2.24}_{-2.31}$   | 31           | LINK            | 02I  | FOCS $\gamma A, \approx 180$ GeV |

$\Gamma(\phi K^+, \phi \rightarrow K^+K^-)/\Gamma(2K^+K^-)$   $\Gamma_{109}/\Gamma_{108}$

| VALUE                                      | DOCUMENT ID     | TECN | COMMENT                       |
|--|-----------------|------|-------------------------------|
| <b><math>0.41 \pm 0.08 \pm 0.03</math></b> | DEL-AMO-SA..11G | BABR | $e^+e^- \approx \Upsilon(4S)$ |

———— Doubly Cabibbo-suppressed modes ————

$\Gamma(2K^+\pi^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{110}/\Gamma_{38}$

| VALUE (units $10^{-3}$ )                        | EVTS         | DOCUMENT ID     | TECN | COMMENT                         |
|---|--------------|-----------------|------|---------------------------------|
| <b><math>2.371 \pm 0.034</math> OUR AVERAGE</b> |              |                 |      |                                 |
| $2.372 \pm 0.024 \pm 0.025$                     | 67k          | AAIJ            | 19G  | LHCB $pp$ at 8 TeV              |
| $2.3 \pm 0.3 \pm 0.2$                           | $356 \pm 52$ | DEL-AMO-SA..11G | BABR | $e^+e^- \approx \Upsilon(4S)$   |
| $2.29 \pm 0.28 \pm 0.12$                        | $281 \pm 34$ | KO              | 09   | BELL $e^+e^-$ at $\Upsilon(4S)$ |
| $5.2 \pm 1.7 \pm 1.1$                           | $27 \pm 9$   | LINK            | 05k  | FOCS $<0.78\%$ , CL = 90%       |

$\Gamma(K^+K^*(892)^0, K^{*0} \rightarrow K^+\pi^-)/\Gamma(2K^+\pi^-)$   $\Gamma_{111}/\Gamma_{110}$

| VALUE                                      | DOCUMENT ID     | TECN | COMMENT                       |
|--|-----------------|------|-------------------------------|
| <b><math>0.47 \pm 0.22 \pm 0.15</math></b> | DEL-AMO-SA..11G | BABR | $e^+e^- \approx \Upsilon(4S)$ |

———— Baryon-antibaryon mode ————

$\Gamma(p\bar{n})/\Gamma_{\text{total}}$   $\Gamma_{112}/\Gamma$

This is the only baryonic mode allowed kinematically.

| VALUE (units $10^{-3}$ )                      | EVTS           | DOCUMENT ID | TECN    | COMMENT                                       |
|---|----------------|-------------|---------|---|
| <b><math>1.22 \pm 0.11</math> OUR AVERAGE</b> |                |             |         |   |
| $1.21 \pm 0.10 \pm 0.05$                      | $193 \pm 17$   | ABLIKIM     | 19oBES3 | $e^+e^-, E_{\text{cm}} = 4178$ MeV            |
| $1.30 \pm 0.36^{+0.12}_{-0.16}$               | $13.0 \pm 3.6$ | ATHAR       | 08      | CLEO $e^+e^-, E_{\text{cm}} \approx 4170$ MeV |

$\Gamma(p\bar{p}e^+\nu_e)/\Gamma_{\text{total}}$   $\Gamma_{113}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                   |
|---|-----|-------------|------|---------------------------|
| <b><math>&lt; 2.0 \times 10^{-4}</math></b> | 90  | ABLIKIM     | 19BD | BES3 $e^+e^-$ at 4178 MeV |

———— Rare or forbidden modes ————

$\Gamma(\pi^+e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{114}/\Gamma$

This mode is not a useful test for a  $\Delta C=1$  weak neutral current because both quarks must change flavor in this decay.

| VALUE   | CL% | DOCUMENT ID        | TECN | COMMENT                            |
|---|-----|--------------------|------|------------------------------------|
| <b><math>&lt; 13 \times 10^{-6}</math></b>                                    | 90  | LEES               | 11G  | BABR $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                    |      |                                    |
| $< 2.2 \times 10^{-5}$  | 90  | <sup>1</sup> RUBIN | 10   | CLEO $e^+e^-$ at 4170 MeV          |
| $< 27 \times 10^{-5}$   | 90  | AITALA             | 99G  | E791 $\pi^- N$ 500 GeV             |

<sup>1</sup>This RUBIN 10 limit is for the  $e^+e^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block.

$\Gamma(\pi^+ \phi, \phi \rightarrow e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{115}/\Gamma$

This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ e^+ e^-$  final state.

| VALUE                                | EVTS | DOCUMENT ID | TECN | COMMENT                    |
|--------------------------------------|------|-------------|------|----------------------------|
| $(6_{-4}^{+8} \pm 1) \times 10^{-6}$ | 3    | RUBIN       | 10   | CLEO $e^+ e^-$ at 4170 MeV |

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{116}/\Gamma$

This mode is not a useful test for a  $\Delta C=1$  weak neutral current because both quarks must change flavor in this decay.

| VALUE   | CL% | DOCUMENT ID | TECN | COMMENT   |
|---|-----|-------------|------|---|
| $< 4.1 \times 10^{-7}$  | 90  | AAIJ        | 13AF | LHCB $pp$ at 7 TeV                              |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |      |   |
| $< 4.3 \times 10^{-5}$  | 90  | LEES        | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$             |
| $< 2.6 \times 10^{-5}$  | 90  | LINK        | 03F  | FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV |
| $< 1.4 \times 10^{-4}$  | 90  | AITALA      | 99G  | E791 $\pi^- N$ 500 GeV                          |
| $< 4.3 \times 10^{-4}$  | 90  | KODAMA      | 95   | E653 $\pi^-$ emulsion 600 GeV                   |

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{117}/\Gamma$

A test for the  $\Delta C=1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE   | CL% | DOCUMENT ID | TECN | COMMENT                             |
|---|-----|-------------|------|-------------------------------------|
| $< 3.7 \times 10^{-6}$  | 90  | LEES        | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |      |                                     |
| $< 5.2 \times 10^{-5}$  | 90  | RUBIN       | 10   | CLEO $e^+ e^-$ at 4170 MeV          |
| $< 1.6 \times 10^{-3}$  | 90  | AITALA      | 99G  | E791 $\pi^- N$ 500 GeV              |

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{118}/\Gamma$

A test for the  $\Delta C=1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE   | CL% | DOCUMENT ID | TECN | COMMENT   |
|---|-----|-------------|------|---|
| $< 21 \times 10^{-6}$   | 90  | LEES        | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$             |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |      |   |
| $< 3.6 \times 10^{-5}$  | 90  | LINK        | 03F  | FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV |
| $< 1.4 \times 10^{-4}$  | 90  | AITALA      | 99G  | E791 $\pi^- N$ 500 GeV                          |
| $< 5.9 \times 10^{-4}$  | 90  | KODAMA      | 95   | E653 $\pi^-$ emulsion 600 GeV                   |

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{119}/\Gamma$

A test for the  $\Delta C=1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE                  | CL% | DOCUMENT ID | TECN | COMMENT                       |
|------------------------|-----|-------------|------|-------------------------------|
| $< 1.4 \times 10^{-3}$ | 90  | KODAMA      | 95   | E653 $\pi^-$ emulsion 600 GeV |

$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{120}/\Gamma$

A test of lepton-family-number conservation.

| VALUE                 | CL% | DOCUMENT ID | TECN | COMMENT                             |
|-----------------------|-----|-------------|------|-------------------------------------|
| $< 12 \times 10^{-6}$ | 90  | LEES        | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{121}/\Gamma$

A test of lepton-family-number conservation.

| VALUE                 | CL% | DOCUMENT ID | TECN | COMMENT                             |
|-----------------------|-----|-------------|------|-------------------------------------|
| $< 20 \times 10^{-6}$ | 90  | LEES        | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{122}/\Gamma$

A test of lepton-family-number conservation.

| VALUE                | CL% | DOCUMENT ID | TECN     | COMMENT                        |
|----------------------|-----|-------------|----------|--------------------------------|
| $<14 \times 10^{-6}$ | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{123}/\Gamma$

A test of lepton-family-number conservation.

| VALUE                 | CL% | DOCUMENT ID | TECN     | COMMENT                        |
|-----------------------|-----|-------------|----------|--------------------------------|
| $<9.7 \times 10^{-6}$ | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\pi^- 2e^+)/\Gamma_{\text{total}}$   $\Gamma_{124}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                        |
|---|-----|-------------|----------|--------------------------------|
| $< 4.1 \times 10^{-6}$  | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |                                |
| $< 1.8 \times 10^{-5}$  | 90  | RUBIN       | 10 CLEO  | $e^+ e^-$ at 4170 MeV          |
| $< 69 \times 10^{-5}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV              |

$\Gamma(\pi^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{125}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN      | COMMENT                                    |
|---|-----|-------------|-----------|--|
| $<1.2 \times 10^{-7}$   | 90  | AAIJ        | 13AF LHCb | $pp$ at 7 TeV                              |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |           |  |
| $<1.4 \times 10^{-5}$   | 90  | LEES        | 11G BABR  | $e^+ e^- \approx \Upsilon(4S)$             |
| $<2.9 \times 10^{-5}$   | 90  | LINK        | 03F FOCS  | $\gamma A, \bar{E}_\gamma \approx 180$ GeV |
| $<8.2 \times 10^{-5}$   | 90  | AITALA      | 99G E791  | $\pi^- N$ 500 GeV                          |
| $<4.3 \times 10^{-4}$   | 90  | KODAMA      | 95 E653   | $\pi^-$ emulsion 600 GeV                   |

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{126}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                        |
|---|-----|-------------|----------|--------------------------------|
| $<8.4 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |                                |
| $<7.3 \times 10^{-4}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV              |

$\Gamma(K^- 2e^+)/\Gamma_{\text{total}}$   $\Gamma_{127}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                        |
|---|-----|-------------|----------|--------------------------------|
| $< 5.2 \times 10^{-6}$  | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |                                |
| $< 1.7 \times 10^{-5}$  | 90  | RUBIN       | 10 CLEO  | $e^+ e^-$ at 4170 MeV          |
| $< 63 \times 10^{-5}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV              |

$\Gamma(K^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{128}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                    |
|---|-----|-------------|----------|--|
| $<1.3 \times 10^{-5}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$             |
| $<1.3 \times 10^{-5}$   | 90  | LINK        | 03F FOCS | $\gamma A, \bar{E}_\gamma \approx 180$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |  |
| $<1.8 \times 10^{-4}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV                          |
| $<5.9 \times 10^{-4}$   | 90  | KODAMA      | 95 E653  | $\pi^-$ emulsion 600 GeV                   |

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{129}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                        |
|---|-----|-------------|----------|--------------------------------|
| $<6.1 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |             |          |                                |
| $<6.8 \times 10^{-4}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV              |

$\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{130}/\Gamma$

A test of lepton-number conservation.

| VALUE                 | CL% | DOCUMENT ID | TECN    | COMMENT                  |
|-----------------------|-----|-------------|---------|--------------------------|
| $<1.4 \times 10^{-3}$ | 90  | KODAMA      | 95 E653 | $\pi^-$ emulsion 600 GeV |

$D_s^+ - D_s^-$  CP-VIOLATING DECAY-RATE ASYMMETRIES

This is the difference between  $D_s^+$  and  $D_s^-$  partial widths for the decay to state  $f$ , divided by the sum of the widths:

$$A_{CP}(f) = [\Gamma(D_s^+ \rightarrow f) - \Gamma(D_s^- \rightarrow \bar{f})] / [\Gamma(D_s^+ \rightarrow f) + \Gamma(D_s^- \rightarrow \bar{f})].$$

$A_{CP}(\mu^\pm \nu)$  in  $D_s^+ \rightarrow \mu^+ \nu$ ,  $D_s^- \rightarrow \mu^- \bar{\nu}_\mu$

| VALUE (%)     | DOCUMENT ID  | TECN | COMMENT               |
|---------------|--------------|------|-----------------------|
| $4.8 \pm 6.1$ | ALEXANDER 09 | CLEO | $e^+ e^-$ at 4170 MeV |

$A_{CP}(K^\pm K_S^0)$  in  $D_s^\pm \rightarrow K^\pm K_S^0$

| VALUE (%)   | EVTS | DOCUMENT ID       | TECN     | COMMENT                        |
|---|------|-------------------|----------|--------------------------------|
| <b>0.09 ± 0.26 OUR AVERAGE</b>  |      |                   |          |                                |
| $0.6 \pm 2.8 \pm 0.6$   | 1.8k | ABLIKIM           | 19AMBES3 | $e^+ e^-$ at 4178 MeV          |
| $-0.05 \pm 0.23 \pm 0.24$   | 288k | <sup>1</sup> LEES | 13E BABR | $e^+ e^-$ at $\Upsilon(4S)$    |
| $2.6 \pm 1.5 \pm 0.6$   |      | ONYISI            | 13 CLEO  | $e^+ e^-$ at 4.17 GeV          |
| $0.12 \pm 0.36 \pm 0.22$  |      | KO                | 10 BELL  | $e^+ e^- \approx \Upsilon(4S)$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |      |                   |          |                                |
| $4.7 \pm 1.8 \pm 0.9$   | 4.0k | MENDEZ            | 10 CLEO  | See ONYISI 13                  |
| $4.9 \pm 2.1 \pm 0.9$   |      | ALEXANDER         | 08 CLEO  | See MENDEZ 10                  |

<sup>1</sup>LEES 13E finds that after subtracting the contribution due to  $K^0 - \bar{K}^0$  mixing, the CP asymmetry is  $(+0.28 \pm 0.23 \pm 0.24)\%$ .

$A_{CP}(K^\pm K_L^0)$  in  $D_s^\pm \rightarrow K^\pm K_L^0$

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN     | COMMENT               |
|--------------------------|------|-------------|----------|-----------------------|
| $-1.1 \pm 2.6 \pm 0.6$   | 2.3k | ABLIKIM     | 19AMBES3 | $e^+ e^-$ at 4178 MeV |

$A_{CP}(K^+ K^- \pi^\pm)$  in  $D_s^\pm \rightarrow K^+ K^- \pi^\pm$

| VALUE (%)              | DOCUMENT ID | TECN    | COMMENT               |
|------------------------|-------------|---------|-----------------------|
| $-0.5 \pm 0.8 \pm 0.4$ | ONYISI      | 13 CLEO | $e^+ e^-$ at 4.17 GeV |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

|                       |           |         |               |
|-----------------------|-----------|---------|---------------|
| $0.3 \pm 1.1 \pm 0.8$ | ALEXANDER | 08 CLEO | See ONYISI 13 |
|-----------------------|-----------|---------|---------------|

$A_{CP}(\phi \pi^\pm)$  in  $D_s^\pm \rightarrow \phi \pi^\pm$

| VALUE (%)                 | DOCUMENT ID | TECN   | COMMENT                |
|---------------------------|-------------|--------|------------------------|
| $-0.38 \pm 0.26 \pm 0.08$ | ABAZOV      | 14B D0 | $p\bar{p}$ at 1.96 TeV |

**$A_{CP}(K^\pm K_S^0 \pi^0)$  in  $D_s^\pm \rightarrow K^\pm K_S^0 \pi^0$**

| <u>VALUE (%)</u>                         | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|--|--------------------|-------------|-----------------------|
| <b><math>-1.6 \pm 6.0 \pm 1.1</math></b> | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |

**$A_{CP}(2K_S^0 \pi^\pm)$  in  $D_s^\pm \rightarrow 2K_S^0 \pi^\pm$**

| <u>VALUE (%)</u>                        | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|---|--------------------|-------------|-----------------------|
| <b><math>3.1 \pm 5.2 \pm 0.6</math></b> | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |

**$A_{CP}(K^+ K^- \pi^\pm \pi^0)$  in  $D_s^\pm \rightarrow K^+ K^- \pi^\pm \pi^0$**

| <u>VALUE (%)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|---|--------------------|-------------|-----------------------|
| <b><math>0.0 \pm 2.7 \pm 1.2</math></b>                                       | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |             |                       |
| $-5.9 \pm 4.2 \pm 1.2$  | ALEXANDER 08       | CLEO        | See ONYISI 13         |

**$A_{CP}(K^\pm K_S^0 \pi^+ \pi^-)$  in  $D_s^\pm \rightarrow K^\pm K_S^0 \pi^+ \pi^-$**

| <u>VALUE (%)</u>                         | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|--|--------------------|-------------|-----------------------|
| <b><math>-5.7 \pm 5.3 \pm 0.9</math></b> | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |

**$A_{CP}(K_S^0 K^\mp 2\pi^\pm)$  in  $D_s^\pm \rightarrow K_S^0 K^\mp 2\pi^\pm$**

| <u>VALUE (%)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|---|--------------------|-------------|-----------------------|
| <b><math>4.1 \pm 2.7 \pm 0.9</math></b>                                       | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |             |                       |
| $-0.7 \pm 3.6 \pm 1.1$  | ALEXANDER 08       | CLEO        | See ONYISI 13         |

**$A_{CP}(\pi^+ \pi^- \pi^\pm)$  in  $D_s^\pm \rightarrow \pi^+ \pi^- \pi^\pm$**

| <u>VALUE (%)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|---|--------------------|-------------|-----------------------|
| <b><math>-0.7 \pm 3.0 \pm 0.6</math></b>                                      | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |             |                       |
| $2.0 \pm 4.6 \pm 0.7$   | ALEXANDER 08       | CLEO        | See ONYISI 13         |

**$A_{CP}(\pi^\pm \eta)$  in  $D_s^\pm \rightarrow \pi^\pm \eta$**

| <u>VALUE (%)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|---|-------------|--------------------|-------------|-----------------------|
| <b><math>1.1 \pm 3.0 \pm 0.8</math></b>                                       |             | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |                    |             |                       |
| $-4.6 \pm 2.9 \pm 0.3$  | 2.5k        | MENDEZ 10          | CLEO        | See ONYISI 13         |
| $-8.2 \pm 5.2 \pm 0.8$  |             | ALEXANDER 08       | CLEO        | See MENDEZ 10         |

**$A_{CP}(\pi^\pm \eta')$  in  $D_s^\pm \rightarrow \pi^\pm \eta'$**

| <u>VALUE (%)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|---|-------------|--------------------|-------------|-----------------------|
| <b><math>-0.9 \pm 0.5</math> OUR AVERAGE</b>                                  |             |                    |             |                       |
| $-0.82 \pm 0.36 \pm 0.35$   | 152k        | AAIJ 17AF          | LHCB        | $pp$ at 7, 8 TeV      |
| $-2.2 \pm 2.2 \pm 0.6$  |             | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |                    |             |                       |
| $-6.1 \pm 3.0 \pm 0.3$  | 1.4k        | MENDEZ 10          | CLEO        | See ONYISI 13         |
| $-5.5 \pm 3.7 \pm 1.2$  |             | ALEXANDER 08       | CLEO        | See MENDEZ 10         |

**$A_{CP}(\eta \pi^\pm \pi^0)$  in  $D_s^\pm \rightarrow \eta \pi^\pm \pi^0$**

| <u>VALUE (%)</u>                         | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>        |
|--|--------------------|-------------|-----------------------|
| <b><math>-0.5 \pm 3.9 \pm 2.0</math></b> | ONYISI 13          | CLEO        | $e^+ e^-$ at 4.17 GeV |



**$A_{CP}(\eta' \pi^\pm \pi^0)$  in  $D_s^\pm \rightarrow \eta' \pi^\pm \pi^0$** 

| VALUE (%)                                | DOCUMENT ID | TECN | COMMENT                    |
|--|-------------|------|----------------------------|
| <b><math>-0.4 \pm 7.4 \pm 1.9</math></b> | ONYISI      | 13   | CLEO $e^+ e^-$ at 4.17 GeV |

 **$A_{CP}(K^\pm \pi^0)$  in  $D_s^\pm \rightarrow K^\pm \pi^0$** 

| VALUE (%)                                  | EVTS         | DOCUMENT ID | TECN | COMMENT                    |
|--|--------------|-------------|------|----------------------------|
| <b><math>-26.6 \pm 23.8 \pm 0.9</math></b> | $202 \pm 70$ | MENDEZ      | 10   | CLEO $e^+ e^-$ at 4170 MeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|            |       |     |      |               |
|------------|-------|-----|------|---------------|
| $2 \pm 29$ | ADAMS | 07A | CLEO | See MENDEZ 10 |
|------------|-------|-----|------|---------------|

 **$A_{CP}(\bar{K}^0 / K^0 \pi^\pm)$  in  $D_s^+ \rightarrow \bar{K}^0 \pi^+$ ,  $D_s^- \rightarrow K^0 \pi^-$** 

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

**0.4 ± 0.5 OUR AVERAGE**

|                          |      |                   |      |                                  |
|--------------------------|------|-------------------|------|----------------------------------|
| $0.38 \pm 0.46 \pm 0.17$ | 121k | <sup>1</sup> AAIJ | 14BD | LHCB $pp$ at 7, 8 TeV            |
| $0.3 \pm 2.0 \pm 0.3$    | 14k  | LEES              | 13E  | BABR $e^+ e^-$ at $\Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                          |     |      |     |                    |
|--------------------------|-----|------|-----|--------------------|
| $0.61 \pm 0.83 \pm 0.14$ | 26k | AAIJ | 13W | LHCB See AAIJ 14BD |
|--------------------------|-----|------|-----|--------------------|

<sup>1</sup>AAIJ 14BD reports its result as  $A_{CP}(D_s^\pm \rightarrow K_S^0 K^\pm)$  with  $CP$ -violation effects in the  $K^0 - \bar{K}^0$  system subtracted. It also measures  $A_{CP}(D^\pm \rightarrow \bar{K}^0 / K^0 K^\pm) + A_{CP}(D_s^\pm \rightarrow \bar{K}^0 / K^0 \pi^\pm) = (0.41 \pm 0.49 \pm 0.26)\%$ .

 **$A_{CP}(K_S^0 \pi^\pm)$  in  $D_s^\pm \rightarrow K_S^0 \pi^\pm$** 

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

**0.20 ± 0.18 OUR AVERAGE**

|                          |      |        |     |                                     |
|--------------------------|------|--------|-----|-------------------------------------|
| $0.16 \pm 0.17 \pm 0.05$ | 721k | AAIJ   | 19T | LHCB $pp$ at 7, 8, 13 TeV           |
| $0.6 \pm 2.0 \pm 0.3$    | 14k  | LEES   | 13E | BABR $e^+ e^-$ at $\Upsilon(4S)$    |
| $5.45 \pm 2.50 \pm 0.33$ |      | KO     | 10  | BELL $e^+ e^- \approx \Upsilon(4S)$ |
| $16.3 \pm 7.3 \pm 0.3$   | 0.4k | MENDEZ | 10  | CLEO $e^+ e^-$ at 4170 MeV          |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|             |       |     |      |               |
|-------------|-------|-----|------|---------------|
| $27 \pm 11$ | ADAMS | 07A | CLEO | See MENDEZ 10 |
|-------------|-------|-----|------|---------------|

 **$A_{CP}(K^\pm \pi^+ \pi^-)$  in  $D_s^\pm \rightarrow K^\pm \pi^+ \pi^-$** 

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

|   |        |    |                            |
|---|--------|----|----------------------------|
| <b><math>4.5 \pm 4.8 \pm 0.6</math></b> | ONYISI | 13 | CLEO $e^+ e^-$ at 4.17 GeV |
|---|--------|----|----------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                        |           |    |      |               |
|------------------------|-----------|----|------|---------------|
| $11.2 \pm 7.0 \pm 0.9$ | ALEXANDER | 08 | CLEO | See ONYISI 13 |
|------------------------|-----------|----|------|---------------|

 **$A_{CP}(K^\pm \eta)$  in  $D_s^\pm \rightarrow K^\pm \eta$** 

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

|  |              |        |    |                            |
|--|--------------|--------|----|----------------------------|
| <b><math>9.3 \pm 15.2 \pm 0.9</math></b> | $222 \pm 41$ | MENDEZ | 10 | CLEO $e^+ e^-$ at 4170 MeV |
|--|--------------|--------|----|----------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|              |       |     |      |               |
|--------------|-------|-----|------|---------------|
| $-20 \pm 18$ | ADAMS | 07A | CLEO | See MENDEZ 10 |
|--------------|-------|-----|------|---------------|

 **$A_{CP}(K^\pm \eta'(958))$  in  $D_s^\pm \rightarrow K^\pm \eta'(958)$** 

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

|  |             |        |    |                            |
|--|-------------|--------|----|----------------------------|
| <b><math>6.0 \pm 18.9 \pm 0.9</math></b> | $56 \pm 17$ | MENDEZ | 10 | CLEO $e^+ e^-$ at 4170 MeV |
|--|-------------|--------|----|----------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

–17 ±37 ADAMS 07A CLEO See MENDEZ 10

### CP VIOLATING ASYMMETRIES OF P-ODD (T-ODD) MOMENTS

#### $A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D_S^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$  is a parity-odd correlation of the  $K^+$ ,  $\pi^+$ , and  $\pi^-$  momenta for the  $D_S^+$ .  $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$  is the corresponding quantity for the  $D_S^-$ . Then

$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$ , and

$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]$ , and

$A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ .  $C_T$  and  $\bar{C}_T$  are commonly referred to as  $T$ -odd moments, because they are odd under  $T$  reversal. However, the  $T$ -conjugate process  $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D_S^\pm$  is not accessible, while the  $P$ -conjugate process is.

| VALUE (units $10^{-3}$ )  | EVTS        | DOCUMENT ID | TECN     | COMMENT  |
|---|-------------|-------------|----------|--|
| <b>–13.6 ± 7.7 ± 3.4</b>  | 29.8 ± 0.3k | LEES        | 11E BABR | $e^+ e^- \approx \Upsilon(4S)$                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |             |          |  |
| –36 ±67 ±23   | 508 ± 34    | LINK        | 05E FOCS | $\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$ |

### $D_S^+$ Semileptonic Form Factors and Decay Constants

#### $r_2 \equiv A_2(0)/A_1(0)$ in $D_S^+ \rightarrow \phi \ell^+ \nu_\ell$

| VALUE                                     | EVTS                                | DOCUMENT ID         | TECN      | COMMENT                              |
|---|-------------------------------------|---------------------|-----------|--------------------------------------|
| <b>0.84 ± 0.11 OUR AVERAGE</b>            | Error includes scale factor of 2.4. |                     |           |                                      |
| 0.816 ± 0.036 ± 0.030                     | 25 ± 0.5k                           | <sup>1</sup> AUBERT | 08AN BABR | $\phi e^+ \nu_e$                     |
| 0.713 ± 0.202 ± 0.284                     | 793                                 | LINK                | 04C FOCS  | $\phi \mu^+ \nu_\mu$                 |
| 1.57 ± 0.25 ± 0.19                        | 271                                 | AITALA              | 99D E791  | $\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$ |
| 1.4 ± 0.5 ± 0.3                           | 308                                 | AVERY               | 94B CLE2  | $\phi e^+ \nu_e$                     |
| 1.1 ± 0.8 ± 0.1                           | 90                                  | FRABETTI            | 94F E687  | $\phi \mu^+ \nu_\mu$                 |
| 2.1 <sup>+0.6</sup> <sub>–0.5</sub> ± 0.2 | 19                                  | KODAMA              | 93 E653   | $\phi \mu^+ \nu_\mu$                 |

<sup>1</sup>To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at  $m_A = 2.5 \text{ GeV}/c^2$  and  $m_V = 2.1 \text{ GeV}/c^2$ . A simultaneous fit to  $r_2$ ,  $r_V$ ,  $r_0$  (a significant  $s$ -wave contribution) and  $m_A$ , gives  $r_2 = 0.763 \pm 0.071 \pm 0.065$ .

#### $r_V \equiv V(0)/A_1(0)$ in $D_S^+ \rightarrow \phi \ell^+ \nu_\ell$

| VALUE                                     | EVTS      | DOCUMENT ID         | TECN      | COMMENT                              |
|---|-----------|---------------------|-----------|--------------------------------------|
| <b>1.80 ± 0.08 OUR AVERAGE</b>            |           |                     |           |                                      |
| 1.807 ± 0.046 ± 0.065                     | 25 ± 0.5k | <sup>1</sup> AUBERT | 08AN BABR | $\phi e^+ \nu_e$                     |
| 1.549 ± 0.250 ± 0.148                     | 793       | LINK                | 04C FOCS  | $\phi \mu^+ \nu_\mu$                 |
| 2.27 ± 0.35 ± 0.22                        | 271       | AITALA              | 99D E791  | $\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$ |
| 0.9 ± 0.6 ± 0.3                           | 308       | AVERY               | 94B CLE2  | $\phi e^+ \nu_e$                     |
| 1.8 ± 0.9 ± 0.2                           | 90        | FRABETTI            | 94F E687  | $\phi \mu^+ \nu_\mu$                 |
| 2.3 <sup>+1.1</sup> <sub>–0.9</sub> ± 0.4 | 19        | KODAMA              | 93 E653   | $\phi \mu^+ \nu_\mu$                 |

<sup>1</sup>To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at  $m_A = 2.5 \text{ GeV}/c^2$  and  $m_V = 2.1 \text{ GeV}/c^2$ . A simultaneous fit to  $r_2$ ,  $r_V$ ,  $r_0$  (a significant  $s$ -wave contribution) and  $m_A$ , gives  $r_V = 1.849 \pm 0.060 \pm 0.095$ .

$\Gamma_L/\Gamma_T$  in  $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$ 

| VALUE                        | EVTS | DOCUMENT ID           | TECN     | COMMENT              |
|------------------------------|------|-----------------------|----------|----------------------|
| <b>0.72±0.18 OUR AVERAGE</b> |      |                       |          |                      |
| 1.0 ±0.3 ±0.2                | 308  | AVERY                 | 94B CLE2 | $\phi e^+ \nu_e$     |
| 1.0 ±0.5 ±0.1                | 90   | <sup>1</sup> FRABETTI | 94F E687 | $\phi \mu^+ \nu_\mu$ |
| 0.54±0.21±0.10               | 19   | <sup>1</sup> KODAMA   | 93 E653  | $\phi \mu^+ \nu_\mu$ |

<sup>1</sup>FRABETTI 94F and KODAMA 93 evaluate  $\Gamma_L/\Gamma_T$  for a lepton mass of zero.

 $f_+(0) |V_{cs}|$  in  $D_s^+ \rightarrow \eta e^+ \nu_e$ 

| VALUE                       | EVTS | DOCUMENT ID | TECN     | COMMENT               |
|-----------------------------|------|-------------|----------|-----------------------|
| <b>0.4455±0.0053±0.0044</b> | 1.8k | ABLIKIM     | 19S BES3 | $e^+ e^-$ at 4178 MeV |

 $f_+(0) |V_{cs}|$  in  $D_s^+ \rightarrow \eta' e^+ \nu_e$ 

| VALUE                    | EVTS | DOCUMENT ID | TECN     | COMMENT               |
|--------------------------|------|-------------|----------|-----------------------|
| <b>0.477±0.049±0.011</b> | 261  | ABLIKIM     | 19S BES3 | $e^+ e^-$ at 4178 MeV |

 $f_+(0) |V_{cd}|$  in  $D_s^+ \rightarrow K^0 e^+ \nu_e$ 

| VALUE                    | EVTS | DOCUMENT ID          | TECN     | COMMENT           |
|--------------------------|------|----------------------|----------|-------------------|
| <b>0.162±0.019±0.003</b> | 117  | <sup>1</sup> ABLIKIM | 19D BES3 | $K_S^0 e^+ \nu_e$ |

<sup>1</sup>Using a two parameter fit in the z expansion.

 $r_V \equiv V(0)/A_1(0)$  in  $D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e$ 

| VALUE                 | EVTS | DOCUMENT ID | TECN     | COMMENT               |
|-----------------------|------|-------------|----------|-----------------------|
| <b>1.67±0.34±0.16</b> | 155  | ABLIKIM     | 19D BES3 | $e^+ e^-$ at 4178 MeV |

 $r_2 \equiv A_2(0)/A_1(0)$  in  $D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e$ 

| VALUE                 | EVTS | DOCUMENT ID | TECN     | COMMENT               |
|-----------------------|------|-------------|----------|-----------------------|
| <b>0.77±0.28±0.07</b> | 155  | ABLIKIM     | 19D BES3 | $e^+ e^-$ at 4178 MeV |

 $f_{D_s^+} |V_{cs}|$  in  $D_s^+ \rightarrow \mu^+ \nu_\mu$ 

| VALUE (MeV)          | EVTS | DOCUMENT ID | TECN     | COMMENT               |
|----------------------|------|-------------|----------|-----------------------|
| <b>246.2±3.6±3.5</b> | 1.1k | ABLIKIM     | 19E BES3 | $e^+ e^-$ at 4178 MeV |

 $D_s^\pm$  REFERENCES

|         |      |                |                          |                  |
|---------|------|----------------|--------------------------|------------------|
| ABLIKIM | 20R  | JHEP 2008 146  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| AAIJ    | 19G  | JHEP 1903 176  | R. Aaij <i>et al.</i>    | (LHCb Collab.)   |
| AAIJ    | 19T  | PRL 122 191803 | R. Aaij <i>et al.</i>    | (LHCb Collab.)   |
| ABLIKIM | 19AD | PR D99 072002  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AH | PR D99 091101  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AM | PR D99 112005  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19BD | PR D100 112008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19BE | PRL 123 112001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19D  | PRL 122 061801 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19E  | PRL 122 071802 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19O  | PR D99 031101  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19S  | PRL 122 121801 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18A  | PR D97 012006  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| AAIJ    | 17AF | PL B771 21     | R. Aaij <i>et al.</i>    | (LHCb Collab.)   |
| AAIJ    | 17AN | PRL 119 101801 | R. Aaij <i>et al.</i>    | (LHCb Collab.)   |
| ABLIKIM | 16O  | PR D94 072004  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 16T  | PR D94 112003  | M. Ablikim <i>et al.</i> | (BESIII Collab.) |

|               |      |                        |                                       |                         |
|---------------|------|------------------------|---------------------------------------|-------------------------|
| ABLIKIM       | 15Z  | PL B750 466            | M. Ablikim <i>et al.</i>              | (BESIII Collab.)        |
| HIETALA       | 15   | PR D92 012009          | J. Hietala <i>et al.</i>              | (MINN, LUTH, OXF)       |
| LEES          | 15D  | PR D91 019901 (errat.) | J.P. Lees <i>et al.</i>               | (BABAR Collab.)         |
| AAIJ          | 14BD | JHEP 1410 025          | R. Aaij <i>et al.</i>                 | (LHCb Collab.)          |
| ABAZOV        | 14B  | PRL 112 111804         | V.M. Abazov <i>et al.</i>             | (D0 Collab.)            |
| AAIJ          | 13AF | PL B724 203            | R. Aaij <i>et al.</i>                 | (LHCb Collab.)          |
| AAIJ          | 13V  | JHEP 1306 065          | R. Aaij <i>et al.</i>                 | (LHCb Collab.)          |
| AAIJ          | 13W  | JHEP 1306 112          | R. Aaij <i>et al.</i>                 | (LHCb Collab.)          |
| LEES          | 13E  | PR D87 052012          | J.P. Lees <i>et al.</i>               | (BABAR Collab.)         |
| ONYISI        | 13   | PR D88 032009          | P.U.E. Onyisi <i>et al.</i>           | (CLEO Collab.)          |
| ZUPANC        | 13   | JHEP 1309 139          | A. Zupanc <i>et al.</i>               | (BELLE Collab.)         |
| DEL-AMO-SA... | 11G  | PR D83 052001          | P. del Amo Sanchez <i>et al.</i>      | (BABAR Collab.)         |
| LEES          | 11E  | PR D84 031103          | J.P. Lees <i>et al.</i>               | (BABAR Collab.)         |
| LEES          | 11G  | PR D84 072006          | J.P. Lees <i>et al.</i>               | (BABAR Collab.)         |
| MARTIN        | 11   | PR D84 012005          | L. Martin <i>et al.</i>               | (CLEO Collab.)          |
| ASNER         | 10   | PR D81 052007          | D.M. Asner <i>et al.</i>              | (CLEO Collab.)          |
| DEL-AMO-SA... | 10J  | PR D82 091103          | P. del Amo Sanchez <i>et al.</i>      | (BABAR Collab.)         |
| Also          |      | PR D91 019901 (errat.) | J.P. Lees <i>et al.</i>               | (BABAR Collab.)         |
| KO            | 10   | PRL 104 181602         | B.R. Ko <i>et al.</i>                 | (BELLE Collab.)         |
| MENDEZ        | 10   | PR D81 052013          | H. Mendez <i>et al.</i>               | (CLEO Collab.)          |
| RUBIN         | 10   | PR D82 092007          | P. Rubin <i>et al.</i>                | (CLEO Collab.)          |
| ALEXANDER     | 09   | PR D79 052001          | J.P. Alexander <i>et al.</i>          | (CLEO Collab.)          |
| AUBERT        | 09O  | PR D79 032003          | B. Aubert <i>et al.</i>               | (BABAR Collab.)         |
| DOBBS         | 09   | PR D79 112008          | S. Dobbs <i>et al.</i>                | (CLEO Collab.)          |
| ECKLUND       | 09   | PR D80 052009          | K.M. Ecklund <i>et al.</i>            | (CLEO Collab.)          |
| GE            | 09A  | PR D80 051102          | J.Y. Ge <i>et al.</i>                 | (CLEO Collab.)          |
| KO            | 09   | PRL 102 221802         | B.R. Ko <i>et al.</i>                 | (BELLE Collab.)         |
| MITCHELL      | 09A  | PR D79 072008          | R.E. Mitchell <i>et al.</i>           | (CLEO Collab.)          |
| NAIK          | 09A  | PR D80 112004          | P. Naik <i>et al.</i>                 | (CLEO Collab.)          |
| ONYISI        | 09   | PR D79 052002          | P.U.E. Onyisi <i>et al.</i>           | (CLEO Collab.)          |
| WON           | 09   | PR D80 111101          | E. Won <i>et al.</i>                  | (BELLE Collab.)         |
| YELTON        | 09   | PR D80 052007          | J. Yelton <i>et al.</i>               | (CLEO Collab.)          |
| ALEXANDER     | 08   | PRL 100 161804         | J.P. Alexander <i>et al.</i>          | (CLEO Collab.)          |
| ATHAR         | 08   | PRL 100 181802         | S.B. Athar <i>et al.</i>              | (CLEO Collab.)          |
| AUBERT        | 08AN | PR D78 051101          | B. Aubert <i>et al.</i>               | (BABAR Collab.)         |
| ECKLUND       | 08   | PRL 100 161801         | K.M. Ecklund <i>et al.</i>            | (CLEO Collab.)          |
| KLEMPPT       | 08   | EPJ C55 39             | E. Klempt, M. Matveev, A.V. Sarantsev | (BONN+)                 |
| LINK          | 08   | PL B660 147            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| WIDHALM       | 08   | PRL 100 241801         | L. Widhalm <i>et al.</i>              | (BELLE Collab.)         |
| ADAMS         | 07A  | PRL 99 191805          | G.S. Adams <i>et al.</i>              | (CLEO Collab.)          |
| AUBERT        | 07V  | PRL 98 141801          | B. Aubert <i>et al.</i>               | (BABAR Collab.)         |
| PEDLAR        | 07A  | PR D76 072002          | T.K. Pedlar <i>et al.</i>             | (CLEO Collab.)          |
| Also          |      | PRL 99 071802          | M. Artuso <i>et al.</i>               | (CLEO Collab.)          |
| AUBERT        | 06N  | PR D74 031103          | B. Aubert <i>et al.</i>               | (BABAR Collab.)         |
| HUANG         | 06B  | PR D74 112005          | G.S. Huang <i>et al.</i>              | (CLEO Collab.)          |
| PDG           | 06   | JP G33 1               | W.-M. Yao <i>et al.</i>               | (PDG Collab.)           |
| AUBERT        | 05V  | PR D71 091104          | B. Aubert <i>et al.</i>               | (BABAR Collab.)         |
| LINK          | 05E  | PL B622 239            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 05J  | PRL 95 052003          | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 05K  | PL B624 166            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 04   | PL B585 200            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 04C  | PL B586 183            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 04D  | PL B586 191            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 04F  | PL B601 10             | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| ACOSTA        | 03D  | PR D68 072004          | D. Acosta <i>et al.</i>               | (FNAL CDF-II Collab.)   |
| ANISOVICH     | 03   | EPJ A16 229            | V.V. Anisovich <i>et al.</i>          |                         |
| LINK          | 03D  | PL B561 225            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 03F  | PL B572 21             | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| AUBERT        | 02G  | PR D65 091104          | B. Aubert <i>et al.</i>               | (BABAR Collab.)         |
| HEISTER       | 02I  | PL B528 1              | A. Heister <i>et al.</i>              | (ALEPH Collab.)         |
| LINK          | 02I  | PL B541 227            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| LINK          | 02J  | PL B541 243            | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| ABBIENDI      | 01L  | PL B516 236            | G. Abbiendi <i>et al.</i>             | (OPAL Collab.)          |
| AITALA        | 01A  | PRL 86 765             | E.M. Aitala <i>et al.</i>             | (FNAL E791 Collab.)     |
| IORI          | 01   | PL B523 22             | M. Iori <i>et al.</i>                 | (FNAL SELEX Collab.)    |
| LINK          | 01C  | PRL 87 162001          | J.M. Link <i>et al.</i>               | (FNAL FOCUS Collab.)    |
| ALEXANDROV    | 00   | PL B478 31             | Y. Alexandrov <i>et al.</i>           | (CERN BEATRICE Collab.) |
| AITALA        | 99   | PL B445 449            | E.M. Aitala <i>et al.</i>             | (FNAL E791 Collab.)     |
| AITALA        | 99D  | PL B450 294            | E.M. Aitala <i>et al.</i>             | (FNAL E791 Collab.)     |
| AITALA        | 99G  | PL B462 401            | E.M. Aitala <i>et al.</i>             | (FNAL E791 Collab.)     |
| BONVICINI     | 99   | PRL 82 4586            | G. Bonvicini <i>et al.</i>            | (CLEO Collab.)          |

|             |     |               |                                |                          |
|-------------|-----|---------------|--------------------------------|--------------------------|
| CHADHA      | 98  | PR D58 032002 | M. Chada <i>et al.</i>         | (CLEO Collab.)           |
| JESSOP      | 98  | PR D58 052002 | C.P. Jessop <i>et al.</i>      | (CLEO Collab.)           |
| ACCIARRI    | 97F | PL B396 327   | M. Acciarri <i>et al.</i>      | (L3 Collab.)             |
| BALEST      | 97  | PRL 79 1436   | R. Balest <i>et al.</i>        | (CLEO Collab.)           |
| FRABETTI    | 97C | PL B401 131   | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| FRABETTI    | 97D | PL B407 79    | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| ARTUSO      | 96  | PL B378 364   | M. Artuso <i>et al.</i>        | (CLEO Collab.)           |
| BAI         | 95C | PR D52 3781   | J.Z. Bai <i>et al.</i>         | (BES Collab.)            |
| BRANDENB... | 95  | PRL 75 3804   | G.W. Brandenburg <i>et al.</i> | (CLEO Collab.)           |
| FRABETTI    | 95B | PL B351 591   | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| KODAMA      | 95  | PL B345 85    | K. Kodama <i>et al.</i>        | (FNAL E653 Collab.)      |
| ACOSTA      | 94  | PR D49 5690   | D. Acosta <i>et al.</i>        | (CLEO Collab.)           |
| AVERY       | 94B | PL B337 405   | P. Avery <i>et al.</i>         | (CLEO Collab.)           |
| BROWN       | 94  | PR D50 1884   | D. Brown <i>et al.</i>         | (CLEO Collab.)           |
| BUTLER      | 94  | PL B324 255   | F. Butler <i>et al.</i>        | (CLEO Collab.)           |
| FRABETTI    | 94F | PL B328 187   | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| FRABETTI    | 93F | PRL 71 827    | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| FRABETTI    | 93G | PL B313 253   | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| KODAMA      | 93  | PL B309 483   | K. Kodama <i>et al.</i>        | (FNAL E653 Collab.)      |
| ALBRECHT    | 92B | ZPHY C53 361  | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)          |
| ALEXANDER   | 92  | PRL 68 1275   | J. Alexander <i>et al.</i>     | (CLEO Collab.)           |
| AVERY       | 92  | PRL 68 1279   | P. Avery <i>et al.</i>         | (CLEO Collab.)           |
| BARLAG      | 92C | ZPHY C55 383  | S. Barlag <i>et al.</i>        | (ACCMOR Collab.)         |
| Also        |     | ZPHY C48 29   | S. Barlag <i>et al.</i>        | (ACCMOR Collab.)         |
| FRABETTI    | 92  | PL B281 167   | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| ALBRECHT    | 91  | PL B255 634   | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)          |
| ALVAREZ     | 91  | PL B255 639   | M.P. Alvarez <i>et al.</i>     | (CERN NA14/2 Collab.)    |
| ALBRECHT    | 90D | PL B245 315   | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)          |
| ALEXANDER   | 90B | PRL 65 1531   | J. Alexander <i>et al.</i>     | (CLEO Collab.)           |
| BARLAG      | 90C | ZPHY C46 563  | S. Barlag <i>et al.</i>        | (ACCMOR Collab.)         |
| FRABETTI    | 90  | PL B251 639   | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)      |
| ANJOS       | 89E | PL B223 267   | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)      |
| CHEN        | 89  | PL B226 192   | W.Y. Chen <i>et al.</i>        | (CLEO Collab.)           |
| ALBRECHT    | 88  | PL B207 349   | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)          |
| ANJOS       | 88  | PRL 60 897    | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)      |
| RAAB        | 88  | PR D37 2391   | J.R. Raab <i>et al.</i>        | (FNAL E691 Collab.)      |
| BECKER      | 87B | PL B184 277   | H. Becker <i>et al.</i>        | (NA11 and NA32 Collabs.) |
| BLAYLOCK    | 87  | PRL 58 2171   | G.T. Blaylock <i>et al.</i>    | (Mark III Collab.)       |
| USHIDA      | 86  | PRL 56 1767   | N. Ushida <i>et al.</i>        | (FNAL E531 Collab.)      |
| ALBRECHT    | 85D | PL 153B 343   | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)          |
| DERRICK     | 85B | PRL 54 2568   | M. Derrick <i>et al.</i>       | (HRS Collab.)            |
| AIHARA      | 84D | PRL 53 2465   | H. Aihara <i>et al.</i>        | (TPC Collab.)            |
| ALTHOFF     | 84  | PL 136B 130   | M. Althoff <i>et al.</i>       | (TASSO Collab.)          |
| BAILEY      | 84  | PL 139B 320   | R. Bailey <i>et al.</i>        | (ACCMOR Collab.)         |
| CHEN        | 83C | PRL 51 634    | A. Chen <i>et al.</i>          | (CLEO Collab.)           |

### OTHER RELATED PAPERS

|         |    |            |                            |              |
|---------|----|------------|----------------------------|--------------|
| RICHMAN | 95 | RMP 67 893 | J.D. Richman, P.R. Burchat | (UCSB, STAN) |
|---------|----|------------|----------------------------|--------------|