

$J/\psi(1S)$

$$J^G(J^{PC}) = 0^-(1^{--})$$

 $J/\psi(1S)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3096.900±0.006 OUR AVERAGE				
3096.900±0.002±0.006		¹ ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	² ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.66 ±0.19 ±0.02	6.1k	⁴ AAIJ 15BI	LHCB	$pp \rightarrow J/\psi X$
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ±0.3		GRIBUSHIN 96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE 82	GOLI	185 $\pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$
3096.93 ±0.09	502	⁵ ZHOLENTZ 80	REDE	e^+e^-
3097.0 ±1		⁶ BRANDELIK 79C	DASP	e^+e^-

¹ Supersedes AULCHENKO 03.² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$. **$J/\psi(1S)$ WIDTH**

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
92.6 ± 1.7 OUR AVERAGE				
Error includes scale factor of 1.1.				
92.45± 1.40±1.48		¹ ANASHIN 20	KEDR	e^+e^-
96.1 ± 3.2	13k	² ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4 ± 8.9		BAI 95B	BES	e^+e^-
91 ±11 ±6		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 ^{+6.1} _{-5.8}		⁴ HSUEH 92	RVUE	See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
92.94± 1.83		^{5,6} ANASHIN 18A	KEDR	e^+e^-
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there² Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

³The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

⁴Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79c.

⁵Using $\Gamma(e^+e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ from PDG 16.

⁶Superseded by ANASHIN 20 that is based on the same dataset .

⁷Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

$J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 \pm 0.30) %	
Γ_3 ggg	(64.1 \pm 1.0) %	
Γ_4 γgg	(8.8 \pm 1.1) %	
Γ_5 e^+e^-	(5.971 \pm 0.032) %	
Γ_6 $e^+e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+\mu^-$	(5.961 \pm 0.033) %	

Decays involving hadronic resonances

Γ_8 $\rho\pi$	(1.69 \pm 0.15) %	S=2.4
Γ_9 $\rho^0\pi^0$	(5.6 \pm 0.7) $\times 10^{-3}$	
Γ_{10} $\rho(770)^\mp K^\pm K_S^0$	(1.9 \pm 0.4) $\times 10^{-3}$	
Γ_{11} $\rho(1450)\pi$		
Γ_{12} $\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	(2.3 \pm 0.7) $\times 10^{-3}$	
Γ_{13} $\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp$	(3.5 \pm 0.6) $\times 10^{-4}$	
Γ_{14} $\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	(2.7 \pm 0.6) $\times 10^{-4}$	
Γ_{15} $\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958)$	(3.3 \pm 0.7) $\times 10^{-6}$	
Γ_{16} $\rho(1700)\pi$		
Γ_{17} $\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	(1.7 \pm 1.1) $\times 10^{-4}$	
Γ_{18} $\rho(2150)\pi$		
Γ_{19} $\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	(8 \pm 40) $\times 10^{-6}$	
Γ_{20} $\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$		
Γ_{21} $a_2(1320)\rho$	(1.09 \pm 0.22) %	
Γ_{22} $\omega\pi^+\pi^+\pi^-\pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$	
Γ_{23} $\omega\pi^+\pi^-\pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$	
Γ_{24} $\omega\pi^+\pi^-$	(7.2 \pm 1.0) $\times 10^{-3}$	
Γ_{25} $\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$	
Γ_{26} $K^*(892)^0 \bar{K}^*(892)^0$	(2.3 \pm 0.6) $\times 10^{-4}$	
Γ_{27} $K^*(892)^\pm K^*(892)^\mp$	(1.00 \pm 0.22 \pm 0.40) $\times 10^{-3}$	
Γ_{28} $K^*(892)^\pm K^*(700)^\mp$	(1.1 \pm 1.0 \pm 0.6) $\times 10^{-3}$	
Γ_{29} $K_S^0 \pi^- K^*(892)^+ + \text{c.c.}$	(2.0 \pm 0.5) $\times 10^{-3}$	

Γ_{30}	$K_S^0 \pi^- K^*(892)^+ + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$	
Γ_{31}	$K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm_{-0.5}^{0.6}) \times 10^{-6}$	
Γ_{32}	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm_{-0.19}^{0.25}) \times 10^{-4}$	
Γ_{33}	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(1.10 \pm_{-0.14}^{0.60}) \times 10^{-5}$	
Γ_{34}	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(6.2 \pm_{-1.6}^{2.9}) \times 10^{-6}$	
Γ_{35}	$\eta K^*(892)^0 \bar{K}^*(892)^0$	$(1.15 \pm 0.26) \times 10^{-3}$	
Γ_{36}	$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$	
Γ_{37}	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$	
Γ_{38}	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$	
Γ_{39}	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$	
Γ_{40}	$K^*(1410) \bar{K} + \text{c.c.}$		
Γ_{41}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$	
Γ_{42}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(8 \pm 6) \times 10^{-5}$	
Γ_{43}	$K_2^*(1430) \bar{K} + \text{c.c.}$		
Γ_{44}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$	
Γ_{45}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(4.0 \pm 1.0) \times 10^{-4}$	
Γ_{46}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$	
Γ_{47}	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	
Γ_{48}	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow$ $K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(4 \pm 4) \times 10^{-4}$	
Γ_{49}	$K^*(892)^0 \bar{K}_2^*(1770)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$	
Γ_{50}	$\omega K^*(892) \bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	
Γ_{51}	$\bar{K} K^*(892) + \text{c.c.}$		
Γ_{52}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(5.0 \pm 0.5) \times 10^{-3}$	
Γ_{53}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm_{-1.0}^{0.8}) \times 10^{-3}$	S=2.9
Γ_{54}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm_{-0.20}^{0.13}) \times 10^{-3}$	
Γ_{55}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	
Γ_{56}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	
Γ_{57}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	

Γ_{58}	$K_1(1400)^\pm K^\mp$		$(3.8 \pm 1.4) \times 10^{-3}$	
Γ_{59}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$		$(7.7 \pm 1.6) \times 10^{-3}$	
Γ_{60}	$K^*(892)^\pm K^\mp \pi^0$		$(4.1 \pm 1.3) \times 10^{-3}$	
Γ_{61}	$K^*(892)^0 K_S^0 \pi^0$		$(7 \pm 4) \times 10^{-4}$	
Γ_{62}	$\omega \pi^0 \pi^0$		$(3.4 \pm 0.8) \times 10^{-3}$	
Γ_{63}	$\omega \pi^0 \eta$		$(3.4 \pm 1.7) \times 10^{-4}$	
Γ_{64}	$b_1(1235)^\pm \pi^\mp$	[b]	$(3.0 \pm 0.5) \times 10^{-3}$	
Γ_{65}	$\omega K^\pm K_S^0 \pi^\mp$	[b]	$(3.4 \pm 0.5) \times 10^{-3}$	
Γ_{66}	$b_1(1235)^0 \pi^0$		$(2.3 \pm 0.6) \times 10^{-3}$	
Γ_{67}	$\eta K^\pm K_S^0 \pi^\mp$	[b]	$(2.2 \pm 0.4) \times 10^{-3}$	
Γ_{68}	$\phi K^*(892) \bar{K} + \text{c.c.}$		$(2.18 \pm 0.23) \times 10^{-3}$	
Γ_{69}	$\omega K \bar{K}$		$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{70}	$\omega f_0(1710) \rightarrow \omega K \bar{K}$		$(4.8 \pm 1.1) \times 10^{-4}$	
Γ_{71}	$\phi 2(\pi^+ \pi^-)$		$(1.60 \pm 0.32) \times 10^{-3}$	
Γ_{72}	$\Delta(1232)^{++} \bar{p} \pi^-$		$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{73}	$\omega \eta$		$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6
Γ_{74}	$\omega \eta' \pi^+ \pi^-$		$(1.12 \pm 0.13) \times 10^{-3}$	
Γ_{75}	$\phi K \bar{K}$		$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3
Γ_{76}	$\phi K_S^0 K_S^0$		$(5.9 \pm 1.5) \times 10^{-4}$	
Γ_{77}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$		$(3.6 \pm 0.6) \times 10^{-4}$	
Γ_{78}	$\phi K^+ K^-$		$(8.3 \pm 1.1) \times 10^{-4}$	
Γ_{79}	$\phi f_2(1270)$		$(3.2 \pm 0.6) \times 10^{-4}$	
Γ_{80}	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$		$(1.10 \pm 0.29) \times 10^{-3}$	
Γ_{81}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.})$	[b]	$(1.16 \pm 0.05) \times 10^{-3}$	
Γ_{82}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$		$(1.07 \pm 0.08) \times 10^{-3}$	
Γ_{83}	$K^+ K^- f_2'(1525)$		$(1.06 \pm 0.35) \times 10^{-3}$	
Γ_{84}	$\phi f_2'(1525)$		$(8 \pm 4) \times 10^{-4}$	S=2.7
Γ_{85}	$\phi \pi^+ \pi^-$		$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7
Γ_{86}	$\phi \pi^0 \pi^0$		$(5.0 \pm 1.0) \times 10^{-4}$	
Γ_{87}	$\phi K^\pm K_S^0 \pi^\mp$	[b]	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{88}	$\omega f_1(1420)$		$(6.8 \pm 2.4) \times 10^{-4}$	
Γ_{89}	$\phi \eta$		$(7.4 \pm 0.8) \times 10^{-4}$	S=1.5
Γ_{90}	$\Xi^0 \Xi^0$		$(1.17 \pm 0.04) \times 10^{-3}$	
Γ_{91}	$\Xi(1530)^- \Xi^+ + \text{c.c.}$		$(3.18 \pm 0.08) \times 10^{-4}$	
Γ_{92}	$\rho K^- \bar{\Sigma}(1385)^0$		$(5.1 \pm 3.2) \times 10^{-4}$	
Γ_{93}	$\omega \pi^0$		$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4
Γ_{94}	$\omega \pi^0 \rightarrow \pi^+ \pi^- \pi^0$		$(1.7 \pm 0.8) \times 10^{-5}$	
Γ_{95}	$\phi \eta'(958)$		$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2
Γ_{96}	$\phi f_0(980)$		$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9
Γ_{97}	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$		$(2.60 \pm 0.34) \times 10^{-4}$	
Γ_{98}	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$		$(1.8 \pm 0.5) \times 10^{-4}$	
Γ_{99}	$\phi \eta \eta'$		$(2.32 \pm 0.17) \times 10^{-4}$	
Γ_{100}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$		$(4.5 \pm 1.0) \times 10^{-6}$	

Γ_{101}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0 p^0\pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$	
Γ_{102}	$\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$	
Γ_{103}	$\phi a_0(980)^0 \rightarrow \phi\eta\pi^0$	$(4.4 \pm 1.4) \times 10^{-6}$	
Γ_{104}	$\Xi(1530)^0 \Xi^0$	$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{105}	$\Sigma(1385)^- \bar{\Sigma}^+$ (or c.c.)	[b] $(3.1 \pm 0.5) \times 10^{-4}$	
Γ_{106}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{107}	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow$ $\phi\pi^0\pi^+\pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$	
Γ_{108}	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow$ $\phi\pi^0\pi^0\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$	
Γ_{109}	$\eta\pi^+\pi^-$	$(3.8 \pm 0.7) \times 10^{-4}$	
Γ_{110}	$\eta\rho$	$(1.93 \pm 0.23) \times 10^{-4}$	
Γ_{111}	$\omega\eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$	
Γ_{112}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{113}	$\rho\eta'(958)$	$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6
Γ_{114}	$a_2(1320)^\pm \pi^\mp$	[b] < 4.3	$\times 10^{-3}$ CL=90%
Γ_{115}	$K\bar{K}_2^*(1430) + \text{c.c.}$	< 4.0	$\times 10^{-3}$ CL=90%
Γ_{116}	$K_1(1270)^\pm K^\mp$	< 3.0	$\times 10^{-3}$ CL=90%
Γ_{117}	$K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$	
Γ_{118}	$K_S^0\pi^- K_2^*(1430)^+ + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$	
Γ_{119}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	< 2.9	$\times 10^{-3}$ CL=90%
Γ_{120}	$\phi\pi^0$	3×10^{-6} or 1×10^{-7}	
Γ_{121}	$\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$	
Γ_{122}	$\omega f_2'(1525)$	< 2.2	$\times 10^{-4}$ CL=90%
Γ_{123}	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9	$\times 10^{-6}$ CL=95%
Γ_{124}	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	< 6.2	$\times 10^{-5}$
Γ_{125}	$\phi X(1835) \rightarrow \phi p\bar{p}$	< 2.1	$\times 10^{-7}$ CL=90%
Γ_{126}	$\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-$	< 2.8	$\times 10^{-4}$ CL=90%
Γ_{127}	$\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-$	< 6.13	$\times 10^{-5}$ CL=90%
Γ_{128}	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow$ $\eta\phi\pi^+\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	
Γ_{129}	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0 \bar{K}^*(892)^0$	< 2.52	$\times 10^{-4}$ CL=90%
Γ_{130}	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	< 8.2	$\times 10^{-6}$ CL=90%
Γ_{131}	$\Delta(1232)^+ \bar{p}$	< 1	$\times 10^{-4}$ CL=90%
Γ_{132}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	< 4.1	$\times 10^{-6}$ CL=90%
Γ_{133}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	< 1.80	$\times 10^{-3}$ CL=90%
Γ_{134}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	< 1.1	$\times 10^{-5}$ CL=90%
Γ_{135}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 2.1	$\times 10^{-5}$ CL=90%
Γ_{136}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 1.6	$\times 10^{-5}$ CL=90%
Γ_{137}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 5.6	$\times 10^{-5}$ CL=90%
Γ_{138}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 1.1	$\times 10^{-5}$ CL=90%

Decays into stable hadrons

Γ_{139}	$2(\pi^+\pi^-\pi^0)$	$(3.73 \pm 0.32)\%$	S=1.4
Γ_{140}	$3(\pi^+\pi^-\pi^0)$	$(2.9 \pm 0.6)\%$	
Γ_{141}	$\pi^+\pi^-\pi^0$	$(2.10 \pm 0.08)\%$	S=1.6
Γ_{142}	$\pi^+\pi^-\pi^0\pi^0\pi^0$	$(2.71 \pm 0.29)\%$	
Γ_{143}	$\rho^\pm\pi^\mp\pi^0\pi^0$	$(1.41 \pm 0.22)\%$	
Γ_{144}	$\rho^+\rho^-\pi^0$	$(6.0 \pm 1.1) \times 10^{-3}$	
Γ_{145}	$\pi^+\pi^-\pi^0K^+K^-$	$(1.20 \pm 0.30)\%$	
Γ_{146}	$4(\pi^+\pi^-\pi^0)$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{147}	$\pi^+\pi^-K^+K^-$	$(6.86 \pm 0.28) \times 10^{-3}$	
Γ_{148}	$\pi^+\pi^-K_S^0K_L^0$	$(3.8 \pm 0.6) \times 10^{-3}$	
Γ_{149}	$\pi^+\pi^-K_S^0K_S^0$	$(1.68 \pm 0.19) \times 10^{-3}$	
Γ_{150}	$\pi^\pm\pi^0K^\mp K_S^0$	$(5.7 \pm 0.5) \times 10^{-3}$	
Γ_{151}	$K^+K^-K_S^0K_S^0$	$(4.2 \pm 0.7) \times 10^{-4}$	
Γ_{152}	$\pi^+\pi^-K^+K^-\eta$	$(4.7 \pm 0.7) \times 10^{-3}$	
Γ_{153}	$\pi^0\pi^0K^+K^-$	$(2.13 \pm 0.22) \times 10^{-3}$	
Γ_{154}	$\pi^0\pi^0K_S^0K_L^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{155}	$K\bar{K}\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{156}	$K^+K^-\pi^0$	$(2.88 \pm 0.12) \times 10^{-3}$	
Γ_{157}	$K_S^0K^\pm\pi^\mp$	$(5.6 \pm 0.5) \times 10^{-3}$	
Γ_{158}	$K_S^0K_L^0\pi^0$	$(2.06 \pm 0.26) \times 10^{-3}$	
Γ_{159}	$K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K_L^0\pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$	
Γ_{160}	$K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K_L^0\pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$	
Γ_{161}	$K_S^0K_L^0\eta$	$(1.45 \pm 0.33) \times 10^{-3}$	
Γ_{162}	$2(\pi^+\pi^-)$	$(3.57 \pm 0.30) \times 10^{-3}$	
Γ_{163}	$3(\pi^+\pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$	
Γ_{164}	$2(\pi^+\pi^-\pi^0)$	$(1.61 \pm 0.20)\%$	
Γ_{165}	$2(\pi^+\pi^-\eta)$	$(2.26 \pm 0.28) \times 10^{-3}$	
Γ_{166}	$3(\pi^+\pi^-\eta)$	$(7.2 \pm 1.5) \times 10^{-4}$	
Γ_{167}	$\pi^+\pi^-\pi^0\pi^0\eta$	$(2.3 \pm 0.5) \times 10^{-3}$	
Γ_{168}	$\rho^\pm\pi^\mp\pi^0\eta$	$(1.9 \pm 0.8) \times 10^{-3}$	
Γ_{169}	$\rho\bar{\rho}$	$(2.120 \pm 0.029) \times 10^{-3}$	
Γ_{170}	$\rho\bar{\rho}\pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1
Γ_{171}	$\rho\bar{\rho}\pi^+\pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{172}	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	[c] $(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
Γ_{173}	$\rho\bar{\rho}\eta$	$(2.00 \pm 0.12) \times 10^{-3}$	
Γ_{174}	$\rho\bar{\rho}\rho$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{175}	$\rho\bar{\rho}\omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3
Γ_{176}	$\rho\bar{\rho}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0
Γ_{177}	$\rho\bar{\rho}a_0(980) \rightarrow \rho\bar{\rho}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$	
Γ_{178}	$\rho\bar{\rho}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$	

Γ_{179}	$n\bar{n}$		$(2.09 \pm 0.16) \times 10^{-3}$	
Γ_{180}	$n\bar{n}\pi^+\pi^-$		$(4 \pm 4) \times 10^{-3}$	
Γ_{181}	$\Sigma^+\bar{\Sigma}^-$		$(1.50 \pm 0.24) \times 10^{-3}$	
Γ_{182}	$\Sigma^0\bar{\Sigma}^0$		$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4
Γ_{183}	$2(\pi^+\pi^-)K^+K^-$		$(3.1 \pm 1.3) \times 10^{-3}$	
Γ_{184}	$p\bar{n}\pi^-$		$(2.12 \pm 0.09) \times 10^{-3}$	
Γ_{185}	$nN(1440)$		seen	
Γ_{186}	$nN(1520)$		seen	
Γ_{187}	$nN(1535)$		seen	
Γ_{188}	$\Xi^-\bar{\Xi}^+$		$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4
Γ_{189}	$\Lambda\bar{\Lambda}$		$(1.89 \pm 0.09) \times 10^{-3}$	S=2.8
Γ_{190}	$\Lambda\bar{\Sigma}^-\pi^+$ (or c.c.)	[b]	$(8.3 \pm 0.7) \times 10^{-4}$	S=1.2
Γ_{191}	$pK^-\bar{\Lambda} + \text{c.c.}$		$(8.6 \pm 1.1) \times 10^{-4}$	
Γ_{192}	$2(K^+K^-)$		$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{193}	$pK^-\bar{\Sigma}^0$		$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{194}	K^+K^-		$(2.86 \pm 0.21) \times 10^{-4}$	
Γ_{195}	$K_S^0 K_L^0$		$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4
Γ_{196}	$\Lambda\bar{\Lambda}\pi^+\pi^-$		$(4.3 \pm 1.0) \times 10^{-3}$	
Γ_{197}	$\Lambda\bar{\Lambda}\eta$		$(1.62 \pm 0.17) \times 10^{-4}$	
Γ_{198}	$\Lambda\bar{\Lambda}\pi^0$		$(3.8 \pm 0.4) \times 10^{-5}$	
Γ_{199}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$		$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{200}	$\pi^+\pi^-$		$(1.47 \pm 0.14) \times 10^{-4}$	
Γ_{201}	$\Lambda\bar{\Sigma} + \text{c.c.}$		$(2.83 \pm 0.23) \times 10^{-5}$	
Γ_{202}	$K_S^0 K_S^0$		$< 1.4 \times 10^{-8}$	CL=95%

Radiative decays

Γ_{203}	3γ		$(1.16 \pm 0.22) \times 10^{-5}$	
Γ_{204}	4γ		$< 9 \times 10^{-6}$	CL=90%
Γ_{205}	5γ		$< 1.5 \times 10^{-5}$	CL=90%
Γ_{206}	$\gamma\pi^0\pi^0$		$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{207}	$\gamma\eta\pi^0$		$(2.14 \pm 0.31) \times 10^{-5}$	
Γ_{208}	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$		$< 2.5 \times 10^{-6}$	CL=95%
Γ_{209}	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$		$< 6.6 \times 10^{-6}$	CL=95%
Γ_{210}	$\gamma K_S^0 K_S^0$		$(8.1 \pm 0.4) \times 10^{-4}$	
Γ_{211}	$\gamma\eta_c(1S)$		$(1.7 \pm 0.4) \%$	S=1.5
Γ_{212}	$\gamma\eta_c(1S) \rightarrow 3\gamma$		$(3.8 \pm 1.3) \times 10^{-6}$	S=1.1
Γ_{213}	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta'$		$(4.9 \pm 0.8) \times 10^{-5}$	
Γ_{214}	$\gamma\pi^+\pi^-2\pi^0$		$(8.3 \pm 3.1) \times 10^{-3}$	
Γ_{215}	$\gamma\eta\pi\pi$		$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{216}	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$		$(6.2 \pm 2.4) \times 10^{-4}$	
Γ_{217}	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$		$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
Γ_{218}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$		$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
Γ_{219}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$		$(3.0 \pm 0.5) \times 10^{-4}$	
Γ_{220}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$		$< 8.2 \times 10^{-5}$	CL=95%

Γ_{221}	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	< 2.63	$\times 10^{-6}$	CL=90%
Γ_{222}	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	< 1.86	$\times 10^{-6}$	CL=90%
Γ_{223}	$\gamma\rho\rho$	(4.5 ± 0.8)	$\times 10^{-3}$	
Γ_{224}	$\gamma\rho\omega$	< 5.4	$\times 10^{-4}$	CL=90%
Γ_{225}	$\gamma\rho\phi$	< 8.8	$\times 10^{-5}$	CL=90%
Γ_{226}	$\gamma\eta'(958)$	(5.25 ± 0.07)	$\times 10^{-3}$	S=1.3
Γ_{227}	$\gamma 2\pi^+ 2\pi^-$	(2.8 ± 0.5)	$\times 10^{-3}$	S=1.9
Γ_{228}	$\gamma f_2(1270) f_2(1270)$	(9.5 ± 1.7)	$\times 10^{-4}$	
Γ_{229}	$\gamma f_2(1270) f_2(1270)$ (non resonant)	(8.2 ± 1.9)	$\times 10^{-4}$	
Γ_{230}	$\gamma K^+ K^- \pi^+ \pi^-$	(2.1 ± 0.6)	$\times 10^{-3}$	
Γ_{231}	$\gamma f_4(2050)$	(2.7 ± 0.7)	$\times 10^{-3}$	
Γ_{232}	$\gamma\omega\omega$	(1.61 ± 0.33)	$\times 10^{-3}$	
Γ_{233}	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	(1.7 ± 0.4)	$\times 10^{-3}$	S=1.3
Γ_{234}	$\gamma f_2(1270)$	(1.64 ± 0.12)	$\times 10^{-3}$	S=1.3
Γ_{235}	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm_{-0.22}^{0.60})$	$\times 10^{-5}$	
Γ_{236}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	(4.2 ± 1.5)	$\times 10^{-4}$	
Γ_{237}	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	(1.1 ± 0.4)	$\times 10^{-5}$	
Γ_{238}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm_{-0.60}^{0.24})$	$\times 10^{-5}$	
Γ_{239}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm_{-0.5}^{1.0})$	$\times 10^{-4}$	S=1.5
Γ_{240}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	(3.8 ± 0.5)	$\times 10^{-4}$	
Γ_{241}	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	(3.1 ± 1.0)	$\times 10^{-4}$	
Γ_{242}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm_{-0.7}^{1.2})$	$\times 10^{-4}$	
Γ_{243}	$\gamma\eta$	(1.108 ± 0.027)	$\times 10^{-3}$	
Γ_{244}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	(7.9 ± 1.3)	$\times 10^{-4}$	
Γ_{245}	$\gamma f_1(1285)$	(6.1 ± 0.8)	$\times 10^{-4}$	
Γ_{246}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	(4.5 ± 1.2)	$\times 10^{-4}$	
Γ_{247}	$\gamma f_2'(1525)$	$(5.7 \pm_{-0.5}^{0.8})$	$\times 10^{-4}$	S=1.5
Γ_{248}	$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm_{-0.5}^{0.7})$	$\times 10^{-5}$	
Γ_{249}	$\gamma f_2'(1525) \rightarrow \gamma\eta\eta$	(3.4 ± 1.4)	$\times 10^{-5}$	
Γ_{250}	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	(2.8 ± 1.8)	$\times 10^{-4}$	
Γ_{251}	$\gamma f_2(1910) \rightarrow \gamma\omega\omega$	(2.0 ± 1.4)	$\times 10^{-4}$	
Γ_{252}	$\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm_{-0.33}^{0.20})$	$\times 10^{-5}$	
Γ_{253}	$\gamma f_0(1710) \rightarrow \gamma\omega\phi$	(2.5 ± 0.6)	$\times 10^{-4}$	
Γ_{254}	$\gamma f_2(1810) \rightarrow \gamma\eta\eta$	$(5.4 \pm_{-2.4}^{3.5})$	$\times 10^{-5}$	
Γ_{255}	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892)\bar{K}^*(892)$	(7.0 ± 2.2)	$\times 10^{-4}$	
Γ_{256}	$\gamma K^*(892)\bar{K}^*(892)$	(4.0 ± 1.3)	$\times 10^{-3}$	
Γ_{257}	$\gamma\phi\phi$	(4.0 ± 1.2)	$\times 10^{-4}$	S=2.1

Γ_{258}	$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{259}	$\gamma \eta(2225)$	$(3.14 \pm_{-0.19}^{+0.50}) \times 10^{-4}$	
Γ_{260}	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
Γ_{261}	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$	
Γ_{262}	$\gamma \eta(1760) \rightarrow \gamma \gamma \gamma$	$< 4.80 \times 10^{-6}$	CL=90%
Γ_{263}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.7 \pm_{-0.8}^{+0.6}) \times 10^{-4}$	S=1.6
Γ_{264}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \pm_{-0.9}^{+1.5}) \times 10^{-5}$	
Γ_{265}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm_{-1.3}^{+2.0}) \times 10^{-5}$	
Γ_{266}	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$		
Γ_{267}	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	$< 3.56 \times 10^{-6}$	CL=90%
Γ_{268}	$\gamma X(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$	
Γ_{269}	$\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$	
Γ_{270}	$\gamma X(2370) \rightarrow \gamma \eta \eta \eta'$	$< 9.2 \times 10^{-6}$	CL=90%
Γ_{271}	$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \pm_{-0.8}^{+0.7}) \times 10^{-5}$	
Γ_{272}	$\gamma(K \bar{K} \pi) [J^{PC} = 0^{-+}]$	$(7 \pm 4) \times 10^{-4}$	S=2.1
Γ_{273}	$\gamma \pi^0$	$(3.56 \pm 0.17) \times 10^{-5}$	
Γ_{274}	$\gamma p \bar{p} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%
Γ_{275}	$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{276}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \pm_{-0.30}^{+0.60}) \times 10^{-4}$	
Γ_{277}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$	
Γ_{278}	$\gamma f_0(2200)$		
Γ_{279}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$	
Γ_{280}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm_{-0.50}^{+0.19}) \times 10^{-4}$	
Γ_{281}	$\gamma f_J(2220)$		
Γ_{282}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$	CL=90%
Γ_{283}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$	CL=90%
Γ_{284}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$	
Γ_{285}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$	
Γ_{286}	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \pm_{-2.2}^{+2.4}) \times 10^{-5}$	
Γ_{287}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm_{-1.5}^{+4.0}) \times 10^{-5}$	
Γ_{288}	$\gamma f_0(1500) \rightarrow \gamma \pi \pi$	$(1.09 \pm 0.24) \times 10^{-4}$	
Γ_{289}	$\gamma f_0(1500) \rightarrow \gamma \eta \eta$	$(1.7 \pm_{-1.4}^{+0.6}) \times 10^{-5}$	
Γ_{290}	$\gamma A \rightarrow \gamma \text{invisible}$	$[d] < 1.7 \times 10^{-6}$	CL=90%
Γ_{291}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	$[e] < 5 \times 10^{-6}$	CL=90%

Dalitz decays

Γ_{292}	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$
Γ_{293}	$\eta e^+ e^-$	$(1.43 \pm 0.07) \times 10^{-5}$

Γ_{294}	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$	
Γ_{295}	$\eta U \rightarrow \eta e^+ e^-$	< 9.11	$\times 10^{-7}$ CL=90%
Γ_{296}	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	< 2.0	$\times 10^{-7}$ CL=90%
Γ_{297}	$\phi e^+ e^-$	< 1.2	$\times 10^{-7}$ CL=90%

Weak decays

Γ_{298}	$D^- e^+ \nu_e + \text{c.c.}$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{299}	$\overline{D}^0 e^+ e^- + \text{c.c.}$	< 8.5	$\times 10^{-8}$ CL=90%
Γ_{300}	$D_s^- e^+ \nu_e + \text{c.c.}$	< 1.3	$\times 10^{-6}$ CL=90%
Γ_{301}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	< 1.8	$\times 10^{-6}$ CL=90%
Γ_{302}	$D^- \pi^+ + \text{c.c.}$	< 7.5	$\times 10^{-5}$ CL=90%
Γ_{303}	$\overline{D}^0 \overline{K}^0 + \text{c.c.}$	< 1.7	$\times 10^{-4}$ CL=90%
Γ_{304}	$\overline{D}^0 \overline{K}^{*0} + \text{c.c.}$	< 2.5	$\times 10^{-6}$ CL=90%
Γ_{305}	$D_s^- \pi^+ + \text{c.c.}$	< 1.3	$\times 10^{-4}$ CL=90%
Γ_{306}	$D_s^- \rho^+ + \text{c.c.}$	< 1.3	$\times 10^{-5}$ CL=90%

Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

Γ_{307}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$ CL=90%
Γ_{308}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$ CL=90%
Γ_{309}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$ CL=90%
Γ_{310}	$e^\pm \tau^\mp$	LF	< 8.3	$\times 10^{-6}$ CL=90%
Γ_{311}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$ CL=90%
Γ_{312}	$\Lambda_c^+ e^- + \text{c.c.}$		< 6.9	$\times 10^{-8}$ CL=90%

Other decays

Γ_{313}	invisible	< 7	$\times 10^{-4}$ CL=90%
----------------	-----------	-------	-------------------------

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta, p\bar{p}\omega, p\bar{p}\eta'$.

[d] For a narrow state A with mass less than 960 MeV.

[e] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

$J/\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

$81.37 \pm 1.36 \pm 1.30$	¹ ANASHIN	20	KEDR $e^+ e^-$
---	----------------------	----	----------------

••• We do not use the following data for averages, fits, limits, etc. •••

74.1 ± 8.1	BAI	95B	BES $e^+ e^-$
59 ± 24	BALDINI-...	75	FRAG $e^+ e^-$
59 ± 14	BOYARSKI	75	MRK1 $e^+ e^-$
50 ± 25	ESPOSITO	75B	FRAM $e^+ e^-$

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there

$\Gamma(e^+e^-)$ Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.53 ± 0.10 OUR AVERAGE				
5.550 ± 0.056 ± 0.089		1,2 ANASHIN	18A	KEDR e^+e^-
5.36 ^{+0.29} _{-0.28}		3 HSUEH	92	RVUE See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.58 ± 0.05 ± 0.08		4 ABLIKIM	16Q	BES3 3.773 $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.71 ± 0.16	13k	5 ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ± 0.19	7.8k	5 AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ± 0.39		BAI	95B	BES e^+e^-
4.72 ± 0.35		ALEXANDER	89	RVUE See Υ mini-review
4.4 ± 0.6		3 BRANDELIK	79C	DASP e^+e^-
4.6 ± 0.8		6 BALDINI-...	75	FRAG e^+e^-
4.8 ± 0.6		BOYARSKI	75	MRK1 e^+e^-
4.6 ± 1.0		ESPOSITO	75B	FRAM e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

⁴ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.

⁵ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

⁶ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

$\Gamma(\mu^+\mu^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
5.13 ± 0.52	BAI	95B	BES e^+e^-
4.8 ± 0.6	BOYARSKI	75	MRK1 e^+e^-
5 ± 1	ESPOSITO	75B	FRAM e^+e^-

$\Gamma(\gamma\gamma)$ Γ_{307}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C	DASP e^+e^-

$J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(l) in the e^+e^- annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_1\Gamma_5/\Gamma$**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.884 ± 0.048 ± 0.078	1,2 ANASHIN	18A	KEDR	e ⁺ e ⁻
-----------------------	-------------	-----	------	-------------------------------

4 ± 0.8	3 BALDINI-...	75	FRAG	e ⁺ e ⁻
---------	---------------	----	------	-------------------------------

3.9 ± 0.8	3 ESPOSITO	75B	FRAM	e ⁺ e ⁻
-----------	------------	-----	------	-------------------------------

¹ From the cross sections of e⁺e⁻ → e⁺e⁻ and e⁺e⁻ → hadrons near the J/ψ(1S) peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_5\Gamma_5/\Gamma$**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

333.1 ± 6.6 ± 4.0	1,2 ANASHIN	18A	KEDR	e ⁺ e ⁻
-------------------	-------------	-----	------	-------------------------------

332.3 ± 6.4 ± 4.8	ANASHIN	10	KEDR	3.097 e ⁺ e ⁻ → e ⁺ e ⁻
-------------------	---------	----	------	---

350 ± 20	BRANDELIK	79C	DASP	e ⁺ e ⁻
----------	-----------	-----	------	-------------------------------

320 ± 70	3 BALDINI-...	75	FRAG	e ⁺ e ⁻
----------	---------------	----	------	-------------------------------

340 ± 90	3 ESPOSITO	75B	FRAM	e ⁺ e ⁻
----------	------------	-----	------	-------------------------------

360 ± 100	3 FORD	75	SPEC	e ⁺ e ⁻
-----------	--------	----	------	-------------------------------

¹ From the cross sections of e⁺e⁻ → e⁺e⁻ and e⁺e⁻ → hadrons near the J/ψ(1S) peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_7\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

333 ± 4 OUR AVERAGE

333.4 ± 2.5 ± 4.4	ABLIKIM	16Q	BES3	3.773 e ⁺ e ⁻ → μ ⁺ μ ⁻ γ
-------------------	---------	-----	------	---

331.8 ± 5.2 ± 6.3	ANASHIN	10	KEDR	3.097 e ⁺ e ⁻ → μ ⁺ μ ⁻
-------------------	---------	----	------	---

338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO e ⁺ e ⁻ → μ ⁺ μ ⁻ γ
-------------------	-----	-------	-----	--

330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR e ⁺ e ⁻ → μ ⁺ μ ⁻ γ
-------------------	------	--------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90	DASP	75	DASP	e ⁺ e ⁻
----------	------	----	------	-------------------------------

380 ± 50	1 ESPOSITO	75B	FRAM	e ⁺ e ⁻
----------	------------	-----	------	-------------------------------

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(\rho(770)^\mp K^\pm K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{10}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

10.4 ± 1.0 ± 1.9	130	LEES	17D	BABR e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ
-------------------------	-----	------	-----	---

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{23}\Gamma_5/\Gamma$**

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------------------------	------	-------------	------	---------

2.2 ± 0.3 ± 0.2	170	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ π ⁰ γ
------------------------	-----	--------	-----	---

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

53.6±5.0±0.4 788 ¹ AUBERT 07AU BABR 10.6 e⁺e⁻ → ωπ⁺π⁻γ

¹ AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π⁺π⁻π⁰)] = 47.8 ± 3.1 ± 3.2 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{62}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

27.8±3.5±0.2 398 ¹ LEES 18E BABR 10.6 e⁺e⁻ → π⁺π⁻3π⁰γ

¹ LEES 18E reports [$\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π⁺π⁻π⁰)] = 24.8 ± 1.8 ± 2.5 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.28±0.34±0.07 47±12 ¹ LEES 12F BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28±0.40±0.11 25 ± 8 ^{1,2} AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

¹ Dividing by (2/3)² to take twice into account that B(K*⁰ → K⁺π⁻) = 2/3 B(K*⁰ → K⁻π⁺).

² Superseded by LEES 12F.

$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{27}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.80±0.48±0.32 1 ± 5 ¹ LEES 14H BABR e⁺e⁻ → π⁺π⁻K_S⁰K_S⁰γ

¹ Dividing by (1/4)² to take twice into account B(K*(892) → K_S⁰π) = 1/4.

$\Gamma(K_S^0\pi^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{29}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

11.0±2.8 OUR AVERAGE

9.2±1.2±3.2 64 ¹ LEES 17D BABR e⁺e⁻ → K_S⁰K[±]π[∓]π⁰γ

14.8±4.8±1.2 53 ² LEES 14H BABR e⁺e⁻ → π⁺π⁻K_S⁰K_S⁰γ

¹ Dividing by 1/2 to take into account B(K*(892)[±] → K[±]π[∓]) = 1/2.

² Dividing by 1/4 to take into account B(K*(892) → K_S⁰π) = 1/4.

$\Gamma(K_S^0\pi^- K^*(892)^+ + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{30}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.7±1.2±0.3 53 LEES 14H BABR e⁺e⁻ → π⁺π⁻K_S⁰K_S⁰γ

$\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{46}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

25.8±1.4±0.6 710 ^{1,2,3} LEES 12F BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1 317 ^{2,4} AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{47} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
18.6 ± 16.1 ± 0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 9.28 \pm 8.0 \pm 0.32 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{48} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.32 ± 2.00 ± 0.08	8 ± 8	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$\Gamma(K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{49} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.8 ± 0.4 ± 0.3	110 ± 14	¹ AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	

¹ Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{53} \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
29.0 ± 1.7 ± 1.3	AUBERT 08S	BABR	$10.6 e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$		

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{54} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.96 ± 0.85 ± 0.70	155	AUBERT 08S	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$	

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}}$					Γ_{54} / Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.69 ± 0.01^{+0.13}_{-0.20}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{55} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$16.76 \pm 1.70 \pm 1.00$	89	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{56} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$26.6 \pm 2.5 \pm 1.5$	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{57} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$17.70 \pm 1.70 \pm 1.00$	94	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{31} / \Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.28^{+0.16+0.59}_{-0.17-0.52}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{59} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$42.6 \pm 4.8 \pm 7.2$	99	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$.

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{60} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$22.8 \pm 2.8 \pm 6.8$	80	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$.

$$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{32} / \Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.04^{+0.25}_{-0.19}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{33} / \Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.1^{+0.6}_{-0.1}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{34} / \Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 0.7^{+2.8}_{-1.4}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{61} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.60 \pm 0.75 \pm 2.25$	34	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{67} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.3 ± 1.4 ± 0.4	44	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\omega K \bar{K}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{69} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70 ± 1.98 ± 0.03	24	¹ AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega K^+ K^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K \bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 3.3 \pm 1.3 \pm 1.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{71} \Gamma_5 / \Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.95 ± 0.19 ± 0.01	35	¹ AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$

¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+ \pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{76} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.25 ± 0.84 ± 0.03	29	¹ LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{78} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.59 ± 0.62 ± 0.05	163	¹ LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{79} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.79 ± 0.32^{+0.02}_{-0.06}	61	^{1,2,3} LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.08 ± 0.73 ^{+0.04} _{-0.14}	44	^{2,4} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
---	----	-----------------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

³ Using $\pi^+ \pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- f'_2(1525)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{83} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K \bar{K})$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{84} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.2±3.2±0.2	11	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K \bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = 7.2 \pm 2.8 \pm 0.3$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{85} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.47±0.35 OUR AVERAGE				

4.45±0.49±0.05	181	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
----------------	-----	-------------------	----------	---

4.50±0.48±0.05	254 ± 23	² SHEN	09 BELL	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
----------------	----------	-------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.3 ± 0.7 ± 0.1	103	³ AUBERT, BE 06D	BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
-----------------	-----	-----------------------------	------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{86}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.76±0.57±0.03	45	¹ LEES	12F BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
-----------------------	----	-------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.13±0.88±0.03	23	² AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
----------------	----	-----------------------------	------	--

¹LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{89}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

6.1±2.7±0.4	6	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
--------------------	---	---------------------	-----------	--

¹AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{97}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.44±0.19 OUR AVERAGE

1.40±0.25±0.02	57 ± 9	¹ LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
----------------	--------	-------------------	----------	--

1.48±0.27±0.09	60 ± 11	² SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
----------------	---------	-------------------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.02±0.24±0.01	20 ± 5	³ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
----------------	--------	---------------------	-----------	--

¹LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Multiplied by 2/3 to take into account the $\phi\pi^+\pi^-$ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.

³Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{98}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.98±0.26±0.01	16 ± 4	¹ LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
-----------------------	--------	-------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.95 \pm 0.40 \pm 0.01$ 7.0 ± 2.8 ² AUBERT 07AK BABR $10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.48 \pm 0.12 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{109} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.3 ± 0.4 OUR AVERAGE
 $2.34 \pm 0.43 \pm 0.16$ 49 LEES 18 BABR $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$
 $2.23 \pm 0.97 \pm 0.03$ 9 ¹ AUBERT 07AU BABR $10.6 e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$
¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+ \pi^- \pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (22.92 \pm 0.28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{118} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

20.1 ± 9.8 ± 0.5 35 ^{1,2} LEES 14H BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$
¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$.
² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{139} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

303 ± 5 ± 18 4990 AUBERT 07AU BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$
 $\Gamma(\pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{141} \Gamma_5 / \Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.122 ± 0.005 ± 0.008	AUBERT,B	04N	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{142} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

150.0 ± 4.0 ± 15.0 2.3k LEES 18E BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$
 $\Gamma(\rho^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{143} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0 ± 9.0 ± 8.0	1.2k	LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

$\Gamma(\rho^+ \rho^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{144} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0 ± 5.0 ± 3.3	529	LEES	18E BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ

$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{145} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0 ± 4.3 ± 6.4	768	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ

$\Gamma(\pi^+ \pi^- K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{147} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
37.94 ± 0.81 ± 1.10	3.1k	LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ± 1.3 ± 2.1 1.5k ¹ AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

33.6 ± 2.7 ± 2.7 233 ² AUBERT 05D BABR 10.6 e⁺e⁻ → K⁺K⁻π⁺π⁻γ

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$\Gamma(\pi^+ \pi^- K_S^0 K_L^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{148} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.8 ± 2.3 ± 2.1	248	LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _L ⁰ γ

$\Gamma(\pi^+ \pi^- K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{149} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3 ± 0.9 ± 0.5	133	LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _S ⁰ γ

$\Gamma(\pi^\pm \pi^0 K^\mp K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{150} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7 ± 1.9 ± 1.8	393	LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

$\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{151} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 ± 0.1	29	LEES	14H BABR	e ⁺ e ⁻ → K _S ⁰ K _S ⁰ K ⁺ K ⁻ γ

$\Gamma(\pi^+ \pi^- K^+ K^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{152} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9 ± 3.9 ± 0.1	73	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ

¹ AUBERT 07AU reports [Γ(J/ψ(1S) → π⁺π⁻K⁺K⁻η) × Γ(J/ψ(1S) → e⁺e⁻) / Γ_{total}] × [B(η → 2γ)] = 10.2 ± 1.3 ± 0.8 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0 \pi^0 K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{153} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.75 ± 0.81 ± 0.90	388	LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.6 ± 1.1 ± 1.3 203 ¹ AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁰π⁰K⁺K⁻γ

¹ Superseded by LEES 12F.

$$\Gamma(\pi^0 \pi^0 K_S^0 K_L^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{154} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.3±2.3±0.5	47	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{158} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4±1.3±0.6	182	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{159} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.9±0.4	106	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{160} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.7±0.1	37	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{161} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±1.8±0.4	45	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$

$$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{162} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4±0.9±0.4		LEES	12E	BABR $10.6 e^+ e^- \rightarrow 2\pi^+ 2\pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$19.5 \pm 1.4 \pm 1.3$	270	¹ AUBERT	05D	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \gamma$
------------------------	-----	---------------------	-----	---

¹ Superseded by LEES 12E.

$$\Gamma(3(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{163} \Gamma_5 / \Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT	06D	BABR $10.6 e^+ e^- \rightarrow 3(\pi^+ \pi^-) \gamma$

$$\Gamma(2(\pi^+ \pi^- \pi^0)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{164} \Gamma_5 / \Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9±0.5±1.0	761	AUBERT	06D	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^- \pi^0) \gamma$

$$\Gamma(2(\pi^+ \pi^-) \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{165} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \eta \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+ \pi^- \pi^0 \pi^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{167} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
12.8±1.8±2.0	203	LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$

$\Gamma(\omega\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{63}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.96±0.01	27	¹ LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta\gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{168}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta\gamma$


$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{169}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.9±0.6 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			

11.3±0.4±0.3	821	¹ LEES	13O BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
12.9±0.4±0.4	918	² LEES	13Y BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
9.7±1.7		³ ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$
••• We do not use the following data for averages, fits, limits, etc. •••				
12.0±0.6±0.5	438	⁴ AUBERT	06B BABR	$e^+e^- \rightarrow p\bar{p}\gamma$

- ¹ ISR photon reconstructed in the detector
- ² ISR photon undetected
- ³ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.
- ⁴ Superseded by LEES 130

WEIGHTED AVERAGE
11.9±0.6 (Error scaled by 1.8)



$\Gamma(\Sigma^0 \bar{\Sigma}^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{182} \Gamma_5 / \Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
6.4 ± 1.2 ± 0.6	AUBERT	07BD	BABR 10.6 e ⁺ e ⁻ → Σ ⁰ Σ ⁰ γ

$\Gamma(2(\pi^+ \pi^-) K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{183} \Gamma_5 / \Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.75 ± 0.23 ± 0.17	205	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ 2(π ⁺ π ⁻)γ

$\Gamma(\Lambda \bar{\Lambda}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{189} \Gamma_5 / \Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
10.7 ± 0.9 ± 0.7	AUBERT	07BD	BABR 10.6 e ⁺ e ⁻ → ΛΛ̄γ

$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{192} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.00 ± 0.33 ± 0.29	287 ± 24	LEES	12F	BABR 10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.11 ± 0.39 ± 0.30	156 ± 15	¹ AUBERT	07AK	BABR 10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ
4.0 ± 0.7 ± 0.6	38	² AUBERT	05D	BABR 10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{194} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.78 ± 0.11 ± 0.05	462	¹ LEES	15J	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.94 ± 0.11 ± 0.05	462	² LEES	15J	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.42 ± 0.23 ± 0.08	51	³ LEES	13Q	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78 ± 0.11 ± 0.05	462	¹ LEES	15J	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.94 ± 0.11 ± 0.05	462	² LEES	15J	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.42 ± 0.23 ± 0.08	51	³ LEES	13Q	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ

¹ sin φ > 0.

² sin φ < 0.

³ Interference with non-resonant K⁺K⁻ production not taken into account.

J/ψ(1S) BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) × Γ(e⁺e⁻)/Γ_{total} above.

$\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$ Γ_1 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.877 ± 0.005 OUR AVERAGE			
0.878 ± 0.005	BAI	95B	BES e ⁺ e ⁻
0.86 ± 0.02	BOYARSKI	75	MRK1 e ⁺ e ⁻

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}}$ Γ_2 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.135 ± 0.003	^{1,2} SETH	04	RVUE e ⁺ e ⁻
0.17 ± 0.02	¹ BOYARSKI	75	MRK1 e ⁺ e ⁻

¹Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

²Using $B(J/\psi \rightarrow \ell^+ \ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(g g g)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
64.1 ± 1.0	6 M	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- + \text{hadrons}$

¹Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+ \ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma \eta_C)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.79 ± 1.05	200 k	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

¹Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
13.7 ± 0.1 ± 0.7	6 M	BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.971 ± 0.032 OUR AVERAGE				
5.983 ± 0.007 ± 0.037	720k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945 ± 0.067 ± 0.042	15k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B	BES $e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.8 ± 1.3 ± 0.4	¹ ARMSTRONG	96	E760 $\bar{p} p \rightarrow e^+ e^- \gamma$

¹For $E_\gamma > 100$ MeV.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.961 ± 0.033 OUR AVERAGE				
5.973 ± 0.007 ± 0.038	770k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960 ± 0.065 ± 0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ± 0.06 ± 0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ± 0.33		BAI	95B	BES $e^+ e^-$
5.90 ± 0.15 ± 0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+e^-)/\Gamma(\mu^+\mu^-)$ Γ_5/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.0016 ± 0.0031 OUR AVERAGE			
1.0022 ± 0.0044 ± 0.0048	¹ AULCHENKO 14	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
1.0017 ± 0.0017 ± 0.0033	² ABLIKIM 13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
1.002 ± 0.021 ± 0.013	³ ANASHIN 10	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
0.997 ± 0.012 ± 0.006	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.011 ± 0.013 ± 0.016	BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
1.00 ± 0.07	BAI	95B BES	e^+e^-
1.00 ± 0.05	BOYARSKI 75	MRK1	e^+e^-
0.91 ± 0.15	ESPOSITO 75B	FRAM	e^+e^-
0.93 ± 0.10	FORD 75	SPEC	e^+e^-

¹ From 235.3k $J/\psi \rightarrow e^+e^-$ and 156.6k $J/\psi \rightarrow \mu^+\mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+e^-)/\Gamma_{total}$ and $\Gamma(\mu^+\mu^-)/\Gamma_{total}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{total}$ and $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{total}$.

————— **HADRONIC DECAYS** —————

$\Gamma(\rho\pi)/\Gamma_{total}$ Γ_8/Γ

<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.69 ± 0.15 OUR AVERAGE				
Error includes scale factor of 2.4. See the ideogram below.				
2.18 ± 0.19		^{1,2} AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
2.184 ± 0.005 ± 0.201	220k	^{2,3} BAI	04H BES	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
2.091 ± 0.021 ± 0.116		^{2,4} BAI	04H BES	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
1.21 ± 0.20		BAI	96D BES	$e^+e^- \rightarrow \rho\pi$
1.42 ± 0.01 ± 0.19		COFFMAN	88 MRK3	e^+e^-
1.3 ± 0.3	150	FRANKLIN	83 MRK2	e^+e^-
1.6 ± 0.4	183	ALEXANDER	78 PLUT	e^+e^-
1.33 ± 0.21		BRANDELIK	78B DASP	e^+e^-
1.0 ± 0.2	543	BARTEL	76 CNTR	e^+e^-
1.3 ± 0.3	153	JEAN-MARIE	76 MRK1	e^+e^-


¹ From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).

² Not independent of their $B(\pi^+\pi^-\pi^0)$.

³ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

⁴ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

WEIGHTED AVERAGE
 1.69 ± 0.15 (Error scaled by 2.4)



$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$

Γ_8/Γ_{141}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.142 \pm 0.011 \pm 0.026$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.331 ± 0.033	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.
² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

Γ_9/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
$0.328 \pm 0.005 \pm 0.027$	COFFMAN 88	MRK3	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.35 ± 0.08	ALEXANDER 78	PLUT	e^+e^-
0.32 ± 0.08	BRANDELIK 78B	DASP	e^+e^-
0.39 ± 0.11	BARTEL 76	CNTR	e^+e^-
0.37 ± 0.09	JEAN-MARIE 76	MRK1	e^+e^-

$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

Γ_{12}/Γ_{141}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$10.9 \pm 1.7 \pm 2.7$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.80 ± 0.27	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.
² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{13} / \Gamma_{157}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.3±0.8±0.6	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$ $\Gamma_{14} / \Gamma_{156}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±2.0±0.6	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+ \pi^- \eta'(958)) / \Gamma_{\text{total}}$ Γ_{15} / Γ

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
3.28±0.55±0.44	119	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

$\Gamma(\rho(1700)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{17} / \Gamma_{141}$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
8±2±5	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

22±6	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
------	-----	-------------------	----------	--

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{19} / \Gamma_{141}$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
4± 1±20	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

600±250	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
---------	-----	-------------------	----------	--

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{20} / \Gamma_{141}$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
4.0±0.8	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a Dalitz plot analysis in a Veneziano model.

$\Gamma(a_2(1320)\rho) / \Gamma_{\text{total}}$ Γ_{21} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±2.2 OUR AVERAGE				
11.7±0.7±2.5	7584	AUGUSTIN 89	DM2	$J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
8.4±4.5	36	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\omega \pi^+ \pi^+ \pi^- \pi^-) / \Gamma_{\text{total}}$ Γ_{22} / Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
85±34	140	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow 3(\pi^+ \pi^-) \pi^0$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 1.0	OUR AVERAGE			
7.0 ± 1.6	18058	AUGUSTIN 89	DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8 ± 1.6	215	BURMESTER 77D	PLUT	e^+e^-
6.8 ± 1.9	348	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.12 \pm 0.02 \pm 0.13$	14k	¹ ABLIKIM 19AC	BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$
¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.				

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3 ± 0.6	OUR AVERAGE			
$4.3 \pm 0.2 \pm 0.6$	5860	AUGUSTIN 89	DM2	e^+e^-
4.0 ± 1.6	70	BURMESTER 77D	PLUT	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9 ± 0.8	81	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5	90	VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.00 \pm 0.19^{+0.11}_{-0.32}$	323	ABLIKIM 10E	BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K^*(892)^\pm K^*(700)^\mp)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.09 \pm 0.18^{+0.94}_{-0.54}$	655	ABLIKIM 10E	BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(\eta K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM 10C	BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+K^-\pi^0)$ Γ_{41}/Γ_{156}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 1.1 \pm 0.7$	2k	¹ LEES 17C	BABR	$J/\psi \rightarrow K^+K^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{42}/Γ_{157}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.9$	4k	¹ LEES 17C	BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0) / \Gamma(K^+ K^- \pi^0)$ $\Gamma_{44} / \Gamma_{156}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 1.3 \pm 0.9$	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{45} / \Gamma_{157}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 1.2$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{46} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7 ± 2.6	40	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
---------------	----	-------------	------	---

$\Gamma(\omega K^*(892)\bar{K} + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{50} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

61 ± 9 OUR AVERAGE

$62.0 \pm 6.8 \pm 10.6$	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
$65.3 \pm 10.2 \pm 13.5$	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
$53 \pm 14 \pm 14$	530 ± 140	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{52} / \Gamma_{157}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$90.5 \pm 0.9 \pm 3.8$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^+ K^*(892)^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{53} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------


6.0 ^{+0.8} _{-1.0} OUR AVERAGE Error includes scale factor of 2.9. See the ideogram below.

$8.07 \pm 0.04 +0.38 -0.61$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
$4.57 \pm 0.17 \pm 0.70$	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$5.26 \pm 0.13 \pm 0.53$		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp,$ $K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

WEIGHTED AVERAGE
6.0±0.8-1.0 (Error scaled by 2.9)



$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$ $\Gamma_{54} / \Gamma_{156}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
92.4 ± 1.5 ± 3.4	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^0 \bar{K}^*(892)^0 + c.c.) / \Gamma_{total}$ Γ_{56} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE				
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
4.33 ± 0.12 ± 0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.7 ± 0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

$\Gamma(K_1(1400)^\pm K^\mp) / \Gamma_{total}$ Γ_{58} / Γ

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.8 ± 1.2	¹ BAI	99C BES	$e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + c.c.) / \Gamma_{total}$ Γ_{59} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

seen ¹ ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

¹ A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 0.3 \pm 0.7$	509	AUGUSTIN	89 DM2	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

$\Gamma(b_1(1235)^\pm\pi^\mp)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
30 ± 5 OUR AVERAGE				
31 ± 6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$
29 ± 7	87	BURMESTER	77D PLUT	e^+e^-

$\Gamma(\omega K^\pm K_S^0\pi^\mp)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
$37.7 \pm 0.8 \pm 5.8$	1972 ± 41	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$
$29.5 \pm 1.4 \pm 7.0$	879 ± 41	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(b_1(1235)^0\pi^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$23 \pm 3 \pm 5$	229	AUGUSTIN	89 DM2	e^+e^-

$\Gamma(\eta K^\pm K_S^0\pi^\mp)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$21.8 \pm 2.2 \pm 3.4$	232 ± 23	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\eta' K^{*0}\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.66 \pm 0.03 \pm 0.21$	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.48 ± 0.13 OUR AVERAGE			
$1.50 \pm 0.02 \pm 0.19$	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$
$1.47 \pm 0.03 \pm 0.17$	² ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.
² From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.51 \pm 0.09 \pm 0.21$	1.0k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.16 \pm 0.12 \pm 0.29$	1.1k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.3	OUR AVERAGE			
20.8 ± 2.7 ± 3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 ± 3.7 ± 4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7 ± 2.4 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19 ± 4	OUR AVERAGE			
19.8 ± 2.1 ± 3.9		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

¹ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

 $\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.8 ± 1.1 ± 0.3	^{1,2} FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.

 $\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
16.0 ± 1.0 ± 3.0	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

 $\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$ Γ_{72}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.58 ± 0.23 ± 0.40	332	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\omega \eta)/\Gamma_{\text{total}}$ Γ_{73}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.20	OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
2.352 ± 0.273	5k	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta$
1.44 ± 0.40 ± 0.14	13	² AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \omega \eta \gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.



$\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$						Γ_{75}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
17.7± 1.6 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.				
16.6± 1.9±1.2	163 ± 19	LEES	12F BABR	10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$		
21.4± 0.4±2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$		
48 $^{+20}_{-16}$ ±6	9.0 $^{+3.7}_{-3.0}$	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+K^-) K^+$		
14.6± 0.8±2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$		
18 ± 8	14	FELDMAN	77 MRK1	e^+e^-		



¹We have multiplied $K^+ K^-$ measurement by 2 to obtain $K\bar{K}$.

²Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

$\Gamma(\phi f_0(1710) \rightarrow \phi K\bar{K})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6±0.2±0.6	^{1,2} FALVARD	88	DM2 $J/\psi \rightarrow$ hadrons

¹Including interference with $f'_2(1525)$.

²Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

$\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
• • •		We do not use the following data for averages, fits, limits, etc. • • •		
< 0.45	90	FALVARD	88	DM2 $J/\psi \rightarrow$ hadrons
< 0.37	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10±0.09±0.28	233	EATON	84	MRK2 $e^+ e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.16 ±0.05 OUR AVERAGE				
1.096±0.012±0.071	43k	ABLIKIM	16L	BES3 $J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
1.258±0.014±0.078	53k	ABLIKIM	16L	BES3 $J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
1.23 ±0.07 ±0.30	0.8k	ABLIKIM	12P	BES2 $J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
1.50 ±0.08 ±0.38	1k	ABLIKIM	12P	BES2 $J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
1.00 ±0.04 ±0.21	0.6k	HENRARD	87	DM2 $e^+ e^- \rightarrow \Sigma^{*-}$
1.19 ±0.04 ±0.25	0.7k	HENRARD	87	DM2 $e^+ e^- \rightarrow \Sigma^{*+}$
0.86 ±0.18 ±0.22	56	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*-}$
1.03 ±0.24 ±0.25	68	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*+}$

$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.071±0.009±0.082	103k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ±4 OUR AVERAGE		Error includes scale factor of 2.7.		
12.3±0.6±2.0		^{1,2} FALVARD	88	DM2 $J/\psi \rightarrow$ hadrons
4.8±1.8	46	¹ GIDAL	81	MRK2 $J/\psi \rightarrow K^+ K^- K^+ K^-$

¹Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

²Including interference with $f_0(1710)$.

$\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94±0.15 OUR AVERAGE		Error includes scale factor of 1.7.		
1.09±0.02±0.13		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$

0.78±0.03±0.12 FALVARD 88 DM2 $J/\psi \rightarrow$ hadrons
 2.1 ±0.9 23 FELDMAN 77 MRK1 e^+e^-

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ **Γ_{87}/Γ**

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.2±0.8 OUR AVERAGE				
7.4±0.6±1.4	227 ± 19	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$
7.4±0.9±1.1		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
7 ±0.6±1.0	163 ± 15	BECKER	87 MRK3	$e^+e^- \rightarrow$ hadrons


$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ **Γ_{88}/Γ**

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8^{+1.9}_{-1.6}±1.7	111 ⁺³¹ ₋₂₆	BECKER	87 MRK3	$e^+e^- \rightarrow$ hadrons

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$ **Γ_{89}/Γ**

<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.74 ±0.08 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
0.898±0.024±0.089		ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow$ hadr
0.64 ±0.04 ±0.11	346	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.661±0.045±0.078		COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\eta$

WEIGHTED AVERAGE
 0.74±0.08 (Error scaled by 1.5)



$\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$ **Γ_{99}/Γ**

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.32±0.06±0.16	2.2k	¹ ABLIKIM	19AN BES3	$e^+e^- \rightarrow J/\psi \rightarrow$ hadrons

¹Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ **Γ_{90}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.17 ± 0.04 OUR AVERAGE				
1.165 ± 0.004 ± 0.043	135k	ABLIKIM	17E BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons
1.20 ± 0.12 ± 0.21	206	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Xi(1530)^- \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{91}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.318 ± 0.008 OUR AVERAGE				
0.317 ± 0.002 ± 0.008	70k	ABLIKIM	20 BES3	$e^+ e^- \rightarrow J/\psi$
0.59 ± 0.09 ± 0.12	75	HENRARD	87 DM2	$e^+ e^-$


$\Gamma(\rho K^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ **Γ_{92}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.51 ± 0.26 ± 0.18	89	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$ **Γ_{93}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.45 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.538 ± 0.012 ± 0.065	2090	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.482 ± 0.019 ± 0.064		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

WEIGHTED AVERAGE
0.45 ± 0.05 (Error scaled by 1.4)



¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega \pi^0 \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{94} / \Gamma_{141}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8 \pm 3 \pm 2$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model and significance 4.9σ .

$\Gamma(\phi \eta'(958)) / \Gamma_{\text{total}}$ Γ_{95} / Γ


VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ± 0.5 OUR AVERAGE			Error includes scale factor of 2.2. See the ideogram below.		

$5.10 \pm 0.03 \pm 0.32$	31k	ABLIKIM	16M	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$5.46 \pm 0.31 \pm 0.56$		ABLIKIM	05B	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$4.1 \pm 0.3 \pm 0.8$	167	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
$3.08 \pm 0.34 \pm 0.36$		COFFMAN	88	MRK3	$e^+ e^- \rightarrow K^+ K^- \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 13	90	VANNUCCI	77	MRK1	$e^+ e^-$
------	----	----------	----	------	-----------

WEIGHTED AVERAGE
 4.6 ± 0.5 (Error scaled by 2.2)



$\Gamma(\phi f_0(980)) / \Gamma_{\text{total}}$ Γ_{96} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		

$4.6 \pm 0.4 \pm 0.8$		¹ FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.6 ± 0.6	50	¹ GIDAL	81	MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.50±0.80±0.61	355	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.50±0.24	70	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.23±0.75±0.73	52	ABLIKIM	08F	BES $J/\psi \rightarrow \eta\phi f_0(980)$

$\Gamma(\phi a_0(980)^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.37±1.35	¹ ABLIKIM 18D	BES3	$J/\psi \rightarrow \phi\eta\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.0 ± 2.7 ± 2.5	² ABLIKIM 11D	BES3	$J/\psi \rightarrow \phi\eta\pi^0$

¹ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.

² Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.32±0.12±0.07	24 ± 9	HENRARD	87	DM2 $e^+ e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.}))/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31±0.05 OUR AVERAGE				
0.30±0.03±0.07	74 ± 8	HENRARD	87	DM2 $e^+ e^- \rightarrow \Sigma^{*-}$
0.34±0.04±0.07	77 ± 9	HENRARD	87	DM2 $e^+ e^- \rightarrow \Sigma^{*+}$
0.29±0.11±0.10	26	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*-}$
0.31±0.11±0.11	28	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6±0.5 OUR AVERAGE				
3.4±1.8±1.5	1.1k	¹ ABLIKIM 15H	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$
3.2±0.6±0.4		JOUSSET 90	DM2	$J/\psi \rightarrow \phi 2(\pi^+\pi^-)$
2.1±0.5±0.4	25	² JOUSSET 90	DM2	$J/\psi \rightarrow \phi\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.6±0.2±0.1	16	BECKER 87	MRK3	$J/\psi \rightarrow \phi K \bar{K} \pi$

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(15) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{107} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
9.36 ± 2.31 ± 1.54	78	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{108} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
2.08 ± 1.63 ± 1.47	9	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{109} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.78 ± 0.68	471	¹ ABLIKIM	19Q BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \eta \pi^+ \pi^-$

¹ From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow \eta \pi^+ \pi^-$ assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$.

$\Gamma(\eta \rho) / \Gamma_{\text{total}}$ Γ_{110} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193 ± 0.023 OUR AVERAGE				
0.194 ± 0.017 ± 0.029	299	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.193 ± 0.013 ± 0.029		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^+ \pi^- \eta$

$\Gamma(\omega \eta'(958)) / \Gamma_{\text{total}}$ Γ_{111} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89 ± 0.18 OUR AVERAGE				
2.08 ± 0.30 ± 0.14	137	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
2.26 ± 0.43	218	² ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta'$
1.8 ^{+1.0} _{-0.8} ± 0.3	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.66 ± 0.17 ± 0.19		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

² Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+ \pi^- \gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega f_0(980)) / \Gamma_{\text{total}}$ Γ_{112} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.27 ± 0.47	¹ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\rho \eta'(958)) / \Gamma_{\text{total}}$ Γ_{113} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.8 OUR AVERAGE				Error includes scale factor of 1.6.
7.90 ± 0.19 ± 0.49	3476	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
8.3 ± 3.0 ± 1.2	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
11.4 ± 1.4 ± 1.6		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

$\Gamma(a_2(1320)^\pm \pi^\mp) / \Gamma_{\text{total}}$ Γ_{114} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 43 × 10⁻⁴	90	BRAUNSCH...	76 DASP	$e^+ e^-$

$\Gamma(K\bar{K}_2^*(1430)+c.c.)/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<40 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+e^- \rightarrow K^0\bar{K}_2^{*0}$
$<66 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+e^- \rightarrow K^\pm\bar{K}_2^{*\mp}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	¹ BAI 99C	BES	e^+e^-

¹ Assuming $B(K_1(1270) \rightarrow K\rho)=0.42 \pm 0.06$

$\Gamma(K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$8.54^{+1.07+2.35}_{-1.20-2.13}$	ABLIKIM 18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(K_2^*(1430)^0\bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^- K^+K^-$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{120}/Γ

The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	¹ ABLIKIM 15K	BES3	$e^+e^- \rightarrow J/\psi \rightarrow K^+K^-\gamma\gamma$
$0.124 \pm 0.033 \pm 0.030$		35 ± 9	² ABLIKIM 15K	BES3	$e^+e^- \rightarrow J/\psi \rightarrow K^+K^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.4	90	³ ABLIKIM 05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$
<6.8	90	COFFMAN 88	MRK3	$e^+e^- \rightarrow K^+K^-\pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+K^-\pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+K^-\pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.01 \pm 0.58 \pm 0.82$	172		¹ ABLIKIM 15H	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17	90	² FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
-------	----	-------------------------	-----	-------------------------------------

¹ With 3.6σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

$\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.2 \times 10^{-4}$ 90 ¹ VANNUCCI 77 MRK1 $e^+e^- \rightarrow \pi^+\pi^-\pi^0 K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.8 \times 10^{-4}$ 90 ¹ FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

$\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<3.9 \times 10^{-6}$ 95 ABLIKIM 13P BES3 $J/\psi \rightarrow \gamma\pi^0 p\bar{p}$

$\Gamma(\omega X(1835), X \rightarrow \eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$<6.2 \times 10^{-5}$ ¹ ABLIKIM 19AC BES3 $J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.1 \times 10^{-7}$ 90 ¹ ABLIKIM 16K BES3 $J/\psi \rightarrow p\bar{p}K_S^0 K_L^0, p\bar{p}K^+K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

$\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.8 \times 10^{-4}$ 90 ABLIKIM 15H BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<6.13 \times 10^{-5}$ 90 ABLIKIM 15H BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.20 \pm 0.14 \pm 0.37$ 471 ABLIKIM 15H BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.52 \times 10^{-4}$ 90 ABLIKIM 10C BES2 $J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<0.82 \times 10^{-5}$ 90 ABLIKIM 13F BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.2 \times 10^{-3}$ 90 HENRARD 87 DM2 e^+e^-

$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------


$<0.1 \times 10^{-3}$ 90 HENRARD 87 DM2 e^+e^-

$\Gamma(\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{132}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.1 \times 10^{-6}$	90	ABLIKIM 12B	BES3	$J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$	
$\Gamma(\bar{\Lambda}(1520)\Lambda + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{133}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.80 \times 10^{-3}$	90	LU 19	BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$	
$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{134}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.1 \times 10^{-5}$	90	BAI 04G	BES2	$e^+ e^-$	
$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{135}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.1 \times 10^{-5}$	90	BAI 04G	BES2	$e^+ e^-$	
$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{136}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-5}$	90	BAI 04G	BES2	$e^+ e^-$	
$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{137}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.6 \times 10^{-5}$	90	BAI 04G	BES2	$e^+ e^-$	
$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{138}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.1 \times 10^{-5}$	90	BAI 04G	BES2	$e^+ e^-$	

———— STABLE HADRONS ————

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$					Γ_{139}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.73 ± 0.32	OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
4.73 ± 0.44	228k	¹ ABLIKIM 19Q	BES3	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$	
3.25 ± 0.49	46055	AUGUSTIN 89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$	
3.17 ± 0.42	147	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow \text{hadrons}$	
3.64 ± 0.52	1500	BURMESTER 77D	PLUT	$e^+ e^-$	
4 ± 1	675	JEAN-MARIE 76	MRK1	$e^+ e^-$	

¹From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$, assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.



$\Gamma(3(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}$ **Γ_{140} / Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.029±0.006 OUR AVERAGE				
0.028±0.009	11	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow \text{hadrons}$
0.029±0.007	181	JEAN-MARIE 76	MRK1	$e^+ e^-$

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ **Γ_{141} / Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
21.0 ±0.8 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
21.37±0.04 ^{+0.64} _{-0.62}	1.8M	^{1,2} ABLIKIM 12H	BES3	$e^+ e^- \rightarrow J/\psi$
23.0 ±2.0 ±0.4	256	³ AUBERT 07AU	BABR	10.6 $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
21.84±0.05±2.01	220k	^{1,4} BAI 04H	BES	$e^+ e^-$
20.91±0.21±1.16		^{4,5} BAI 04H	BES	$e^+ e^-$
15 ±2	168	FRANKLIN 83	MRK2	$e^+ e^-$


¹ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}} = 0.808 \pm 0.013$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁵ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}$ **Γ_{145}/Γ**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.3	309	VANNUCCI 77	MRK1	$e^+ e^-$

$\Gamma(4(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$ **Γ_{146}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
90±30	13	JEAN-MARIE 76	MRK1	$e^+ e^-$

$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ **Γ_{147}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.2±2.3	205	VANNUCCI 77	MRK1	$e^+ e^-$

$\Gamma(K \bar{K} \pi)/\Gamma_{\text{total}}$ **Γ_{155}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ±10 OUR AVERAGE				
55.2±12.0	25	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow K^+ K^- \pi^0$
78.0±21.0	126	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ **Γ_{156}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.01±0.12	183k	ABLIKIM 19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{156}/\Gamma_{141}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0±0.3±0.9	23k	LEES	17C	BABR $J/\psi \rightarrow h^+ h^- \pi^0$

$\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{157}/\Gamma_{141}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
26.5±0.5±2.1	24k	LEES	17C	BABR $J/\psi \rightarrow h^0 h^+ h^-$

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{total}$ Γ_{162}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.57±0.30 OUR AVERAGE				

3.53±0.12±0.29	1107	¹ ABLIKIM	05H	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+ \pi^-, J/\psi \rightarrow 2(\pi^+ \pi^-)$
----------------	------	----------------------	-----	---

4.0 ±1.0 76 JEAN-MARIE 76 MRK1 $e^+ e^-$
¹ Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{total}$ Γ_{163}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

40±20	32	JEAN-MARIE	76	MRK1 $e^+ e^-$
-------	----	------------	----	----------------

$\Gamma(2(\pi^+ \pi^-)\eta)/\Gamma_{total}$ Γ_{165}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.26±0.08±0.27	4.8k	ABLIKIM	05c	BES2 $e^+ e^- \rightarrow 2(\pi^+ \pi^-)\eta$
-----------------------	------	---------	-----	---

$\Gamma(3(\pi^+ \pi^-)\eta)/\Gamma_{total}$ Γ_{166}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.24±0.96±1.11	616	ABLIKIM	05c	BES2 $e^+ e^- \rightarrow 3(\pi^+ \pi^-)\eta$
-----------------------	-----	---------	-----	---

$\Gamma(\rho\bar{\rho})/\Gamma_{total}$ Γ_{169}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.120±0.029 OUR AVERAGE

2.112±0.004±0.031	314k	ABLIKIM	12c	BES3 $e^+ e^-$
2.17 ±0.16 ±0.04	317	¹ WU	06	BELL $B^+ \rightarrow \rho\bar{\rho}K^+$
2.26 ±0.01 ±0.14	63316	BAI	04E	BES2 $e^+ e^- \rightarrow J/\psi$
1.97 ±0.22	99	BALDINI	98	FENI $e^+ e^-$
1.91 ±0.04 ±0.30		PALLIN	87	DM2 $e^+ e^-$
2.16 ±0.07 ±0.15	1420	EATON	84	MRK2 $e^+ e^-$
2.5 ±0.4	133	BRANDELIK	79c	DASP $e^+ e^-$
2.0 ±0.5		BESCH	78	BONA $e^+ e^-$
2.2 ±0.2	331	² PERUZZI	78	MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ±0.3	48	ANTONELLI	93	SPEC $e^+ e^-$
----------	----	-----------	----	----------------

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \rho\bar{\rho})/\Gamma_{total}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming angular distribution $(1+\cos^2\theta)$.


$\Gamma(\rho\bar{p}\pi^0)/\Gamma_{\text{total}}$ **Γ_{170}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.1.		
1.33 ± 0.02 ± 0.11	11k	ABLIKIM	09B	BES2 e^+e^-
1.13 ± 0.09 ± 0.09	685	EATON	84	MRK2 e^+e^-
1.4 ± 0.4		BRANDELIK	79C	DASP e^+e^-
1.00 ± 0.15	109	PERUZZI	78	MRK1 e^+e^-

$\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{171}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
6.46 ± 0.17 ± 0.43	1435	EATON	84	MRK2 e^+e^-
3.8 ± 1.6	48	BESCH	81	BONA e^+e^-
5.5 ± 0.6	533	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(\rho\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{172}/Γ**

Including $\rho\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
3.36 ± 0.65 ± 0.28	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$ **Γ_{173}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				
1.91 ± 0.02 ± 0.17	13k	¹ ABLIKIM	09	BES2 e^+e^-

$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2	e^+e^-
2.5 ± 1.2		BRANDELIK	79C	DASP	e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1	e^+e^-

¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.


$\Gamma(p\bar{p}\rho)/\Gamma_{total}$ **Γ_{174}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.31 \times 10^{-3}$	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{total}$ **Γ_{175}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.10 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
$0.90 \pm 0.02 \pm 0.09$	2670	ABLIKIM	13P	BES3 e^+e^-
$0.98 \pm 0.03 \pm 0.14$	2449	ABLIKIM	08	BES2 e^+e^-
$1.10 \pm 0.17 \pm 0.18$	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
 0.98 ± 0.10 (Error scaled by 1.3)



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{total}$ **Γ_{176}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.129 ± 0.014 OUR AVERAGE		Error includes scale factor of 2.0.		
$0.126 \pm 0.002 \pm 0.007$	16k	¹ ABLIKIM	19N	BES3 e^+e^-
$0.200 \pm 0.023 \pm 0.028$	265 ± 31	² ABLIKIM	09	BES2 e^+e^-
$0.68 \pm 0.23 \pm 0.17$	19	EATON	84	MRK2 e^+e^-
1.8 ± 0.6	19	PERUZZI	78	MRK1 e^+e^-

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

² From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels.

$\Gamma(\rho\bar{p}a_0(980) \rightarrow \rho\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{177}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.8 \pm 1.2 \pm 1.3$	ABLIKIM	14N	BES3 $e^+e^- \rightarrow J/\psi$

$\Gamma(\rho\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{178}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.519 ± 0.033 OUR AVERAGE				
$0.523 \pm 0.006 \pm 0.033$	14k	ABLIKIM	16K	BES3 $J/\psi \rightarrow \rho\bar{p}K_S^0 K_L^0,$ $\rho\bar{p}K^+ K^-$
$0.45 \pm 0.13 \pm 0.07$		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{179}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.09 ± 0.16 OUR AVERAGE				
$2.07 \pm 0.01 \pm 0.17$	36k	ABLIKIM	12C	BES3 e^+e^-
2.31 ± 0.49	79	BALDINI	98	FENI e^+e^-
1.8 ± 0.9		BESCH	78	BONA e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90 ± 0.55	40	ANTONELLI	93	SPEC e^+e^-

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{180}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8 ± 3.6	5	BESCH	81	BONA e^+e^-

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{181}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.50 \pm 0.10 \pm 0.22$	399	ABLIKIM	08O	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{182}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.172 ± 0.032 OUR AVERAGE		Error includes scale factor of 1.4.		
$1.164 \pm 0.004 \pm 0.023$	111k	ABLIKIM	17L	BES3 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
$1.33 \pm 0.04 \pm 0.11$	1.7k	ABLIKIM	06	BES2 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
$1.06 \pm 0.04 \pm 0.23$	884	PALLIN	87	DM2 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
$1.58 \pm 0.16 \pm 0.25$	90	EATON	84	MRK2 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78	MRK1 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.4 ± 2.6	3	BESCH	81	BONA $e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

$\Gamma(2(\pi^+\pi^-)K^+K^-)/\Gamma_{\text{total}}$ Γ_{183}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
31 ± 13	30	VANNUCCI	77	MRK1 e^+e^-

$\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{184}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.12 ± 0.09 OUR AVERAGE				
$2.36 \pm 0.02 \pm 0.21$	59k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \rho\pi^-\bar{n}$
$2.47 \pm 0.02 \pm 0.24$	55k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \bar{p}\pi^+n$


$2.02 \pm 0.07 \pm 0.16$	1288	EATON	84	MRK2	$e^+e^- \rightarrow \rho\pi^-$
$1.93 \pm 0.07 \pm 0.16$	1191	EATON	84	MRK2	$e^+e^- \rightarrow \bar{\rho}\pi^+$
1.7 ± 0.7	32	BESCH	81	BONA	$e^+e^- \rightarrow \rho\pi^-$
1.6 ± 1.2	5	BESCH	81	BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$
2.16 ± 0.29	194	PERUZZI	78	MRK1	$e^+e^- \rightarrow \rho\pi^-$
2.04 ± 0.27	204	PERUZZI	78	MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$

$\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$

Γ_{188}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
$1.040 \pm 0.006 \pm 0.074$	43k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Xi^- \Xi^+$
$0.90 \pm 0.03 \pm 0.18$	961	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^- \Xi^+$
$0.70 \pm 0.06 \pm 0.12$	132	HENRARD	87 DM2	$e^+e^- \rightarrow \Xi^- \Xi^+$
$1.14 \pm 0.08 \pm 0.20$	194	EATON	84 MRK2	$e^+e^- \rightarrow \Xi^- \Xi^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Xi^- \Xi^+$

WEIGHTED AVERAGE
 0.97 ± 0.08 (Error scaled by 1.4)




$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

Γ_{189}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89 ± 0.09 OUR AVERAGE		Error includes scale factor of 2.8. See the ideogram below.		
$1.943 \pm 0.003 \pm 0.033$	441k	ABLIKIM	17L BES3	e^+e^-
$2.03 \pm 0.03 \pm 0.15$	8887	ABLIKIM	06 BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
$1.96 \begin{smallmatrix} +0.47 \\ -0.44 \end{smallmatrix} \pm 0.04$	46	¹ WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
$1.08 \pm 0.06 \pm 0.24$	631	BAI	98G BES	e^+e^-

1.38 ±0.05 ±0.20	1847	PALLIN	87	DM2	e^+e^-
1.58 ±0.08 ±0.19	365	EATON	84	MRK2	e^+e^-
2.6 ±1.6	5	BESCH	81	BONA	e^+e^-
1.1 ±0.2	196	PERUZZI	78	MRK1	e^+e^-

¹WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\Lambda\bar{\Sigma}^- \pi^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ **Γ_{190}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ±0.07 OUR AVERAGE		Error includes scale factor of 1.2.		
0.770±0.051±0.083	335	¹ ABLIKIM	07H BES2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
0.747±0.056±0.076	254	¹ ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
0.90 ±0.06 ±0.16	225 ± 15	HENRARD	87 DM2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
1.11 ±0.06 ±0.20	342 ± 18	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
1.53 ±0.17 ±0.38	135	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
1.38 ±0.21 ±0.35	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$

¹Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

$\Gamma(pK^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{191}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.11 OUR AVERAGE				
0.84 ^{+0.17} _{-0.15} ±0.02	45	¹ LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
0.89±0.07±0.14	307	EATON	84 MRK2	e^+e^-

¹LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	¹ HUANG 03	BELL	$B^+ \rightarrow 2(K^+K^-)K^+$
0.7 ± 0.3		VANNUCCI 77	MRK1	e^+e^-

¹Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

$\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.29 \pm 0.06 \pm 0.05$	90	EATON	84	MRK2 e^+e^-

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{194}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.86 \pm 0.09 \pm 0.19$	1k	¹ METREVELI 12		$\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.39 \pm 0.24 \pm 0.22$	107	² BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ± 0.9	6	² BRANDELIK 79C	DASP	e^+e^-

¹Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.


²Interference with non-resonant K^+K^- production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.95 ± 0.11 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.		
$1.93 \pm 0.01 \pm 0.05$	110k	ABLIKIM 17AH	BES3	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$
$2.62 \pm 0.15 \pm 0.14$	0.3k	¹ METREVELI 12		$\psi(2S) \rightarrow \pi^+\pi^-K_S^0 K_L^0$
$1.82 \pm 0.04 \pm 0.13$	2.1k	² BAI 04A	BES2	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.18 \pm 0.12 \pm 0.18$		JOUSSET 90	DM2	$J/\psi \rightarrow$ hadrons
$1.01 \pm 0.16 \pm 0.09$	74	BALTRUSAIT..85D	MRK3	e^+e^-

¹Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

²Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$.



$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{196}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30±0.13±0.99	2.4k	ABLIKIM	12P BES2	J/ψ

$\Gamma(\Lambda \bar{\Lambda} \eta)/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				
15.7±0.80±1.54	454	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$
26.2±6.0 ±4.4	44	² ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma \gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma \gamma) = 39.4\%$.

$\Gamma(\Lambda \bar{\Lambda} \pi^0)/\Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 6.4	90		² ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$
23 ±7 ±8		11	BAI	98G BES	$e^+ e^-$
22 ±5 ±5		19	HENRARD	87 DM2	$e^+ e^-$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma \gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\bar{\Lambda} n K_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46±0.20±1.07	1058	¹ ABLIKIM	08C BES2	$e^+ e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{200}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				
1.47±0.13±0.13	140	¹ METREVELI 12		$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-
1.0 ±0.5	5	BRANDELIK 78B	DASP	e^+e^-
1.6 ±1.6	1	VANNUCCI 77	MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{201}/Γ**

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83±0.23 OUR AVERAGE					
2.74±0.24±0.22	234 ± 21	¹ ABLIKIM	12B	BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92±0.22±0.24	308 ± 24	² ABLIKIM	12B	BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
••• We do not use the following data for averages, fits, limits, etc. •••					
<18		² HENRARD	87	DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15	90	PERUZZI	78	MRK1	$e^+e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ **Γ_{202}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁸	95	¹ ABLIKIM 17AH	BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<1 × 10 ⁻⁶	95	¹ BAI 04D	BES	e^+e^-
<5.2 × 10 ⁻⁶	90	¹ BALTRUSAIT..85C	MRK3	e^+e^-

¹ Forbidden by CP.

————— RADIATIVE DECAYS —————

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ **Γ_{203}/Γ**

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.6±2.2 OUR AVERAGE					
11.3±1.8±2.0	113 ± 18	ABLIKIM	13I	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ±3 ±2	24.2 ^{+7.2} _{-6.0}	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

••• We do not use the following data for averages, fits, limits, etc. •••

<55	90	PARTRIDGE 80	CBAL	e^+e^-
-----	----	--------------	------	----------

$\Gamma(4\gamma)/\Gamma_{\text{total}}$ **Γ_{204}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9 × 10⁻⁶	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$ **Γ_{205}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<15 × 10⁻⁶	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$				Γ_{206}/Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT
1.15 ± 0.05		¹ ABLIKIM	15AE BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ The uncertainty is systematic as statistical is negligible.

$\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$			Γ_{207}/Γ	
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
21.4 ± 1.8 ± 2.5	596	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$				Γ_{208}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5 × 10⁻⁶	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$				Γ_{209}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6.6 × 10⁻⁶	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{210}/Γ
VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
8.1 ± 0.4		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$				Γ_{211}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.5.		
2.00 ± 0.31 ± 0.02		¹ MITCHELL	09 CLEO	$e^+e^- \rightarrow \gamma X$
1.27 ± 0.36		GAISER	86 CBAL	$J/\psi \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen		ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
0.79 ± 0.20	273 ± 43	² AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
seen	16	BALTRUSAITIS	84 MRK3	$J/\psi \rightarrow 2\phi\gamma$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by the authors using an average of $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$ from BALTRUSAITIS 86, BISELLO 91, BAI 04 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$				Γ_{212}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8^{+1.3}_{-1.0} OUR AVERAGE		Error includes scale factor of 1.1.		
4.5 ± 1.2 ± 0.6	33 ± 9	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
1.2 ^{+2.7} _{-1.1} ± 0.3	1.2 ^{+2.8} _{-1.1}	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta')/\Gamma_{\text{total}}$ Γ_{213}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.86 ± 0.62 ± 0.45	137	ABLIKIM	21C	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$

$\Gamma(\gamma\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{214}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.3 ± 0.2 ± 3.1	¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

¹ 4π mass less than 2.0 GeV.

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{215}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 ± 1.0 OUR AVERAGE			
5.85 ± 0.3 ± 1.05	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$
7.8 ± 1.2 ± 2.4	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$

¹ Broad enhancement at 1700 MeV.

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{216}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.2 ± 2.2 ± 0.9	BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{217}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
1.66 ± 0.1 ± 0.58	^{1,2} BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
3.8 ± 0.3 ± 0.6	³ AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$
4.0 ± 0.7 ± 1.0	³ EDWARDS	82E	CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{3,4} SCHARRE	80	MRK2 e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.78 ± 0.21 ± 0.33	^{3,5,6} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$
0.83 ± 0.13 ± 0.18	^{3,7,8} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$
0.66 ^{+0.17 +0.24} _{-0.16 -0.15}	^{3,6,9} BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1.03 ^{+0.21 +0.26} _{-0.18 -0.19}	^{3,8,10} BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K\bar{K}\pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

³ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.

⁵ From fit to the $a_0(980)\pi 0^-+$ partial wave.


⁶ $a_0(980)\pi$ mode.

⁷ From fit to the $K^*(892)K 0^-+$ partial wave.

⁸ K^*K mode.

⁹ From $a_0(980)\pi$ final state.

¹⁰ From $K^*(890)K$ final state.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$ **Γ_{218}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
$1.07 \pm 0.17 \pm 0.11$	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$0.64 \pm 0.12 \pm 0.07$	¹ COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ **Γ_{219}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.5 OUR AVERAGE				
$2.6 \pm 0.7 \pm 0.4$		BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$3.38 \pm 0.33 \pm 0.64$		¹ BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$7.0 \pm 0.6 \pm 1.1$	261	² AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Via $a_0(980)\pi$.
² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi) / \Gamma_{\text{total}}$ **Γ_{220}/Γ**

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<82	95		BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$7.03 \pm 0.92 \pm 0.91$	1.3k		¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$10.36 \pm 1.51 \pm 1.54$	1.9k		² ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

$\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{221}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.63 \times 10^{-6}$	90	ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{222}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.86 \times 10^{-6}$	90	ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$					Γ_{223}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
4.5 ± 0.8 OUR AVERAGE					
$4.7 \pm 0.3 \pm 0.9$		¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$	
$3.75 \pm 1.05 \pm 1.20$		² BURKE	82	MRK2	$J/\psi \rightarrow 4\pi\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.09	90	³ BISELLO	89B		$J/\psi \rightarrow 4\pi\gamma$
¹ 4π mass less than 2.0 GeV.					
² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .					
³ 4π mass in the range 2.0–25 GeV.					


$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$					Γ_{224}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.4 \times 10^{-4}$	90	ABLIKIM	08A	BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$					Γ_{225}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8.8 \times 10^{-5}$	90	ABLIKIM	08A	BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$					Γ_{226}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.25 ± 0.07 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
$5.27 \pm 0.03 \pm 0.05$	36k	ABLIKIM	19T	BES	$J/\psi \rightarrow \gamma\eta'$
$5.43 \pm 0.23 \pm 0.09$	5.0k	¹ ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
$4.77 \pm 0.22 \pm 0.06$		² ABLIKIM	11	BES3	$J/\psi \rightarrow \eta'\gamma$
$5.24 \pm 0.12 \pm 0.11$		PEDLAR	09	CLE3	$J/\psi \rightarrow \eta'\gamma$
5.55 ± 0.44	35k	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta'\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$4.50 \pm 0.14 \pm 0.53$		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
$4.30 \pm 0.31 \pm 0.71$		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
$4.04 \pm 0.16 \pm 0.85$	622	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$4.39 \pm 0.09 \pm 0.66$	2420	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$4.1 \pm 0.3 \pm 0.6$		BLOOM	83	CBAL	$e^+e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ± 1.1	6	BRANDELIK	79C	DASP	$e^+e^- \rightarrow 3\gamma$
2.4 ± 0.7	57	BARTEL	76	CNTR	$e^+e^- \rightarrow 2\gamma\rho$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.




$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ **Γ_{227}/Γ**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
4.32 ± 0.14 ± 0.73	¹ BISELLO 89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
2.08 ± 0.13 ± 0.35	² BISELLO 89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
3.05 ± 0.08 ± 0.45	² BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 ± 0.45 ± 1.20	³ BURKE 82	MRK2	e^+e^-

¹ 4π mass less than 3.0 GeV.
² 4π mass less than 2.0 GeV.
³ 4π mass less than 2.5 GeV.

WEIGHTED AVERAGE
 2.8 ± 0.5 (Error scaled by 1.9)



$\Gamma(\gamma f_2(1270) f_2(1270)) / \Gamma_{\text{total}}$ Γ_{228} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 0.7 \pm 1.6$	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant})) / \Gamma_{\text{total}}$ Γ_{229} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$8.2 \pm 0.8 \pm 1.7$	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{230} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.1 \pm 0.1 \pm 0.6$	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{231} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.7 \pm 0.5 \pm 0.5$	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi\pi / \text{total} = 0.167$.

$\Gamma(\gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{232} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61 ± 0.33 OUR AVERAGE				
$6.0 \pm 4.8 \pm 1.8$		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma \omega \pi^+ \pi^-$
$1.41 \pm 0.2 \pm 0.42$	120 ± 17	BISELLO	87 SPEC	$e^+ e^-$, hadrons γ
$1.76 \pm 0.09 \pm 0.45$		BALTRUSAIT..85c	MRK3	$e^+ e^- \rightarrow \text{hadrons } \gamma$


$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ **Γ_{233}/Γ**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ± 0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36 ± 0.38	1,2 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.
² Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ **Γ_{234}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.64 ± 0.12 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
2.07 ± 0.16 ^{+0.02} _{-0.07}	2.4k	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
1.63 ± 0.26 ^{+0.02} _{-0.06}		3 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.42 ± 0.21 ^{+0.01} _{-0.05}		4 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
1.33 ± 0.05 ± 0.20		5 AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.36 ± 0.09 ± 0.23		5 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.48 ± 0.25 ± 0.30	178	EDWARDS	82B CBAL	$e^+e^- \rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER	78 PLUT	e^+e^-
1.2 ± 0.6	30	6 BRANDELIK	78B DASP	$e^+e^- \rightarrow \pi^+\pi^-\gamma$



¹ Using CLEO-c data but not authored by the CLEO Collaboration.
² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi)$

$= (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.

$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{235}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{236}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$4.19 \pm 0.73 \pm 1.34$	478	¹ DOBBS 15	$J/\psi \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{237}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.07^{+0.08+0.36}_{-0.07-0.34}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{238}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.59 \pm 0.16^{+0.18}_{-0.56}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{239}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

$9.5^{+1.0}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.00^{+0.12+1.24}_{-0.08-0.40}$			¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$11.76 \pm 0.54 \pm 0.94$	1.2k		² DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
$9.62 \pm 0.29^{+3.51}_{-1.86}$			³ BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
$5.0 \pm 0.8^{+1.8}_{-0.4}$			^{1,4} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$9.2 \pm 1.4 \pm 1.4$			¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$10.4 \pm 1.2 \pm 1.6$			¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$9.6 \pm 1.2 \pm 1.8$			¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6 ± 0.2	$^{+0.6}_{-0.2}$		1,5 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8		90	6 BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
1.6 ± 0.4	± 0.3		7 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
3.8 ± 1.6			8 EDWARDS	82D CBAL	$e^+e^- \rightarrow \eta\eta\gamma$

¹ Includes unknown branching fraction to K^+K^- or $K_S^0K_S^0$. We have multiplied K^+K^- measurement by 2, and $K_S^0K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to K^+K^- or $K_S^0K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.


⁵ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁶ Includes unknown branching fraction to $\rho^0\rho^0$.

⁷ Includes unknown branching fraction to $\pi^+\pi^-$.

⁸ Includes unknown branching fraction to $\eta\eta$.

WEIGHTED AVERAGE
9.5+1.0-0.5 (Error scaled by 1.5)



$\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi) / \Gamma_{\text{total}}$

Γ_{240} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5	OUR AVERAGE			
$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.5 ± 1.6	± 0.8	BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$
---------------	-----------	-----	---------	---------------------------------------

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{241} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.06 ± 0.08	180	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{242} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35^{+0.13+1.24}_{-0.11-0.74}	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma \eta) / \Gamma_{\text{total}}$ Γ_{243} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.108 ± 0.027 OUR AVERAGE				
1.12 ± 0.05 ± 0.01	18.6k	¹ ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
1.101 ± 0.029 ± 0.022		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.11		BLOOM	83	CBAL $e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C	DASP $e^+ e^-$
1.3 ± 0.4	21	BARTEL	77	CNTR $e^+ e^-$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{244} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79 ± 0.13 OUR AVERAGE			
0.68 ± 0.04 ± 0.24	BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.76 ± 0.15 ± 0.21	^{1,2} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
0.87 ± 0.14 ^{+0.14} _{-0.11}	¹ BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

² From fit to the $K^*(892) K 1^{++}$ partial wave.

$\Gamma(\gamma f_1(1285)) / \Gamma_{\text{total}}$ Γ_{245} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.61 ± 0.08 OUR AVERAGE			
0.69 ± 0.16 ± 0.20	¹ BAI	04J	BES2 $J/\psi \rightarrow \gamma \gamma \rho^0$
0.61 ± 0.04 ± 0.21	² BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 ± 0.09 ± 0.17	³ BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.625 ± 0.063 ± 0.103	⁴ BOLTON	92	MRK3 $J/\psi \rightarrow \gamma f_1(1285)$
0.70 ± 0.08 ± 0.16	⁵ BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.

² Assuming $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} = 0.090 \pm 0.004$.

³ Assuming $\Gamma(f_1(1285) \rightarrow \eta\pi\pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.

⁴ Obtained summing the sequential decay channels

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi\pi\pi\pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow \eta\pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma\rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}.$$

⁵ Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

$\Gamma(\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{246}/Γ


VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.7$	BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}}$ Γ_{247}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$5.7^{+0.8}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.1 $\pm 0.9 \pm 0.2$	750	1,2	DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
3.85 $\pm 0.17^{+1.91}_{-0.73}$		3	BAI	03G	BES $J/\psi \rightarrow \gamma K\bar{K}$
3.6 $\pm 0.4^{+1.4}_{-0.4}$		3	BAI	96C	BES $J/\psi \rightarrow \gamma K^+K^-$
5.6 $\pm 1.4 \pm 0.9$		3	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+K^-$
4.5 $\pm 0.4 \pm 0.9$		3	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.8 $\pm 1.6 \pm 1.4$		3	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<3.4	90	4	BRANDELIK	79C	DASP $e^+e^- \rightarrow \pi^+\pi^-\gamma$
<2.3	90	3	ALEXANDER	78	PLUT $e^+e^- \rightarrow K^+K^-\gamma$



¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{248}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{249}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$ Γ_{250}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$ Γ_{251}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{252}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \phi)/\Gamma_{\text{total}}$ Γ_{253}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.5 ± 0.6 OUR AVERAGE				
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma \omega \phi$
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma \omega \phi$

$\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{254}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{255}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.7 \pm 0.1 \pm 0.2$	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{256}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.0 \pm 0.3 \pm 1.3$	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

¹ Summed over all charges.


$\Gamma(\gamma \phi \phi)/\Gamma_{\text{total}}$ Γ_{257}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.0 ± 1.2 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
$3.4 \pm 0.8 \pm 0.6$	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.1 \pm 0.7 \pm 0.4$		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ $\phi \phi$ mass less than 2.9 GeV, η_c excluded.



$\Gamma(\gamma \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{258}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$0.38 \pm 0.07 \pm 0.07$ 49 EATON 84 MRK2 $e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.11 90 PERUZZI 78 MRK1 $e^+ e^-$

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$ Γ_{259}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.14^{+0.50}_{-0.19}$				OUR AVERAGE
$2.40 \pm 0.10^{+2.47}_{-0.18}$		1,2 ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$4.4 \pm 0.4 \pm 0.8$	196	2 ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.3 \pm 0.8 \pm 0.5$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$2.7 \pm 0.6 \pm 0.6$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$2.4^{+1.5}_{-1.0}$		3,4 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for $\eta(2100)$, 0^{-+} phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^{-+} state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

² Includes unknown branching fraction to $\phi\phi$.

³ Estimated by us from various fits.

⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{260}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.09	1,2 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{261}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{262}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.80 \times 10^{-6}$	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{263}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.7^{+0.6}_{-0.8}$				OUR AVERAGE Error includes scale factor of 1.6.

$3.93 \pm 0.38^{+0.31}_{-0.84}$ ¹ ABLIKIM 16J BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

$2.2 \pm 0.4 \pm 0.4$ 264 ABLIKIM 05R BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.87 \pm 0.09^{+0.49}_{-0.52}$ 4265 ² ABLIKIM 11C BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma X(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{264}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

$0.77^{+0.15}_{-0.09}$ OUR AVERAGE

$0.90^{+0.04+0.27}_{-0.11-0.55}$		¹ ABLIKIM	12D	BES3	$J/\psi \rightarrow \gamma p\bar{p}$
$1.14^{+0.43+0.42}_{-0.30-0.26}$	231	² ALEXANDER	10	CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
$0.70 \pm 0.04^{+0.19}_{-0.08}$		BAI	03F	BES2	$J/\psi \rightarrow \gamma p\bar{p}$

¹ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A.

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

$\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{265}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$3.31^{+0.33+1.96}_{-0.30-1.29}$ ABLIKIM 15T BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\phi(1020))/\Gamma_{\text{total}}$ Γ_{266}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.77 \pm 0.35 \pm 0.25$	305	¹ ABLIKIM	18i	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$8.09 \pm 1.99 \pm 1.36$	1.3k	² ABLIKIM	18i	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{267}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

$< 3.56 \times 10^{-6}$ 90 ABLIKIM 180 BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma X(1840) \rightarrow \gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{271}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

$2.44 \pm 0.36^{+0.60}_{-0.74}$ 0.6k ABLIKIM 13U BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

$\Gamma(\gamma X(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$ Γ_{268}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$1.79 \pm 0.23 \pm 0.65$ ABLIKIM 20Q BES3 $J/\psi \rightarrow \gamma K^+ K^- \eta'$

$\Gamma(\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$ Γ_{269}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$1.18 \pm 0.32 \pm 0.39$ ABLIKIM 20Q BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$

$\Gamma(\gamma X(2370) \rightarrow \gamma \eta \eta \eta') / \Gamma_{\text{total}}$ Γ_{270} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9.2	90	ABLIKIM	21C	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta \eta'$

$\Gamma(\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}]) / \Gamma_{\text{total}}$ Γ_{272} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
0.58 ± 0.03 ± 0.20	¹ BAI	00D	BES $J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$
2.1 ± 0.1 ± 0.7	² BAI	00D	BES $J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$

¹ For a broad structure around 1800 MeV.
² For a broad structure around 2040 MeV.

$\Gamma(\gamma \pi^0) / \Gamma_{\text{total}}$ Γ_{273} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.56 ± 0.17 OUR AVERAGE				
3.59 ± 0.20 ± 0.03	1.6k	¹ ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
3.63 ± 0.36 ± 0.13		PEDLAR	09	CLE3 $J/\psi \rightarrow \pi^0 \gamma$
3.13 ^{+0.65} _{-0.47}	586	ABLIKIM	06E	BES2 $J/\psi \rightarrow \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.6 ± 1.1 ± 0.7		BLOOM	83	CBAL $e^+ e^-$
7.3 ± 4.7	10	BRANDELIK	79C	DASP $e^+ e^-$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma \pi^0) / \Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \pi^0) / \Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma \rho \bar{\rho} \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{274} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.79 × 10⁻³	90	EATON	84	MRK2 $e^+ e^-$

$\Gamma(\gamma \Lambda \bar{\Lambda}) / \Gamma_{\text{total}}$ Γ_{275} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.13 × 10⁻³	90	HENRARD	87	DM2 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.16 × 10 ⁻³	90	BAI	98G	BES $e^+ e^-$
--------------------------	----	-----	-----	---------------

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{276} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.13^{+0.09+0.64}_{-0.10-0.28}	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{277} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200)) / \Gamma_{\text{total}}$ Γ_{278} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-----------------------	----	-----	---

¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{279} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.86 \pm 0.49 \pm 1.20$	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{280} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.72^{+0.08+0.17}_{-0.06-0.47}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_J(2220)) / \Gamma_{\text{total}}$ Γ_{281} / Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300			¹ BAI	96B	BES	$e^+ e^- \rightarrow \gamma \bar{p} p, K \bar{K}$
>250	99.9		² HASAN	96	SPEC	$\bar{p} p \rightarrow \pi^+ \pi^-$
< 2.3	95		³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95		³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	³ BALTRUSAIT..86D	MRK3		$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	³ BALTRUSAIT..86D	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{282} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.9	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 $\pm 8 \pm 4$	BAI	98H	BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
$8.4 \pm 2.6 \pm 3.0$	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{283}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6		³ DEL-AMO-SA...100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
< 2.9		³ DEL-AMO-SA...100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.6 \pm 2.9 \pm 2.4$		BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$		BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{284}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.6 \pm 0.5$	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma \rho \bar{\rho}$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{285}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.95 \pm 0.21^{+0.66}_{-0.72}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{286}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{287}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$5.54^{+0.34+3.82}_{-0.40-1.49}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{288}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.24 OUR AVERAGE				
$1.21 \pm 0.29 \pm 0.24$	174	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
$1.00 \pm 0.03 \pm 0.45$		² ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1.02 \pm 0.09 \pm 0.45$		² ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.7 ± 0.8		^{3,4} BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
---------------	--	---------------------	----	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S-wave dipions.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{289} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma A \rightarrow \gamma \text{invisible}) / \Gamma_{\text{total}}$ Γ_{290} / Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-6}$	90	88M	¹ ABLIKIM	20K BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$<6.3 \times 10^{-6}$	90	3.7M	² INSLER	10 CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ For a narrow state A with mass $m_A < 1.2$ GeV. The limit varies with m_A , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_A = 0$.

² The limit varies with mass m_A of a narrow state A and is 4.3×10^{-6} for $m_A = 0$, reaches its largest value of 6.3×10^{-6} at $m_A = 500$ MeV, and is 3.6×10^{-6} at $m_A = 960$ MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-) / \Gamma_{\text{total}}$ Γ_{291} / Γ
 (narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.5 \times 10^{-5}$	90	¹ ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$<2.1 \times 10^{-5}$	90	² ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

————— DALITZ DECAYS —————

$\Gamma(\pi^0 e^+ e^-) / \Gamma_{\text{total}}$ Γ_{292} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

$\Gamma(\eta e^+ e^-) / \Gamma_{\text{total}}$ Γ_{293} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.43 \pm 0.04 \pm 0.06$	2.47k	^{1,2} ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$
$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM	14I BES3	$J/\psi \rightarrow \eta e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.84 \pm 0.11 \pm 0.08$ GeV. Supersedes ABLIKIM 14I.

$\Gamma(\eta'(958) e^+ e^-) / \Gamma_{\text{total}}$ Γ_{294} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.59 \pm 0.07 \pm 0.17$	8.9k	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.81 \pm 0.16 \pm 0.31$ 1.4k ^{1,2} ABLIKIM 14I BES3 $J/\psi \rightarrow \eta'(958) e^+ e^-$

¹ Using both $\eta' \rightarrow \gamma \pi^+ \pi^-$ and $\eta' \rightarrow \pi^+ \pi^- \eta$ decays.

² Superseded by ABLIKIM 19H.

$\Gamma(\eta U \rightarrow \eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{295}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.11 \times 10^{-7}$	90	¹ ABLIKIM 19A	BES3	$J/\psi \rightarrow \eta e^+ e^-$

¹ For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

$\Gamma(\eta'(958) U \rightarrow \eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{296}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-7}$	90	¹ ABLIKIM 19H	BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

¹ For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

$\Gamma(\phi e^+ e^-)/\Gamma_{\text{total}}$ Γ_{297}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ ABLIKIM 19AB	BES3	$J/\psi \rightarrow \phi e^+ e^-$

¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) = (34.49 \pm 0.30)\%$.

WEAK DECAYS

$\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{298}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-5}$	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{299}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-8}$	90	¹ ABLIKIM 17AF	BES3	$e^+ e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-5}$ 90 ABLIKIM 06M BES2 $e^+ e^- \rightarrow J/\psi$

¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{300}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.6 \times 10^{-5}$ 90 ¹ ABLIKIM 06M BES2 $e^+ e^- \rightarrow J/\psi$

¹ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5 \%$.

$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{301}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{302}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.5 \times 10^{-5}$	90	ABLIKIM	08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{303}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-4}$	90	ABLIKIM	08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{304}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	90	ABLIKIM	14K	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{305}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-4}$	90	ABLIKIM	08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{306}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	ABLIKIM	14K	BES3	$e^+ e^- \rightarrow J/\psi$

————— CHARGE CONJUGATION (C), PARITY (P), —————
 ————— LEPTON FAMILY NUMBER (LF) VIOLATING MODES —————

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{307}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.7 \times 10^{-7}$	90	ABLIKIM	14Q	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.5 \times 10^{-5}$	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$<1.6 \times 10^{-4}$	90	¹ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
$<2.2 \times 10^{-5}$	90	ABLIKIM	07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$<50 \times 10^{-5}$	90	BARTEL	77	CNTR	$e^+ e^-$

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.020 \times 10^{-3}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$					Γ_{308}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-6}$	90	ABLIKIM	14Q	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					Γ_{309}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-7}$	90	ABLIKIM	13L	BES3	$e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.1 \times 10^{-6}$	90	BAI	03D	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$					Γ_{310}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8.3 \times 10^{-6}$	90	ABLIKIM	04	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$			Γ_{312}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.9 \times 10^{-8}$	90	ABLIKIM	19AF BES3	$e^+ e^- \rightarrow J/\psi \rightarrow p K^- \pi^+ e^-$ (+ c.c.)	

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$			Γ_{311}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	ABLIKIM	04 BES	$e^+ e^- \rightarrow J/\psi$	

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$			Γ_{313}/Γ_5		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-2}$	90	LEES	13I BABR	$B \rightarrow K^{(*)} J/\psi$	

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$			Γ_{313}/Γ_7		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

$J/\psi(1S)$ REFERENCES

ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	20	JHEP 2007 112	V. V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)

AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13L	PR D87 112007	Ablikim M. <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	10O	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)

AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
BALTRUSAIT...	84	PRL 52 2126	Translated from YAF 41 733.	(CIT, UCSC+)
EATON	84	PR D29 804	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BLOOM	83	ARNS 33 143	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
EDWARDS	83B	PRL 51 859	E.D. Bloom, C. Peck	(SLAC, CIT)
FRANKLIN	83	PRL 51 963	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BURKE	82	PRL 49 632	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82E	PRL 49 259	E.D. Bloom, C. Peck	(SLAC, CIT)
LEMOIGNE	82	PL 113B 509	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BESCH	81	ZPHY C8 1	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
GIDAL	81	PL 107B 153	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
PARTRIDGE	80	PRL 44 712	G. Gidal <i>et al.</i>	(SLAC, LBL)
SCHARRE	80	PL 97B 329	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
ZHOLENTZ	80	PL 96B 214	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	79C	ZPHY C1 233	Translated from YAF 34 1471.	(NOVO)
ALEXANDER	78	PL 72B 493	R. Brandelik <i>et al.</i>	(DASP Collab.)
BESCH	78	PL 78B 347	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BRANDELIK	78B	PL 74B 292	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
PERUZZI	78	PR D17 2901	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	77	PL 66B 489	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BURMESTER	77D	PL 72B 135	W. Bartel <i>et al.</i>	(DESY, HEIDP)
FELDMAN	77	PRPL 33C 285	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
VANNUCCI	77	PR D15 1814	G.J. Feldman, M.L. Perl	(LBL, SLAC)
BARTEL	76	PL 64B 483	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BRAUNSCH...	76	PL 63B 487	W. Bartel <i>et al.</i>	(DESY, HEIDP)
JEAN-MARIE	76	PRL 36 291	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BALDINI-...	75	PL 58B 471	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BOYARSKI	75	PRL 34 1357	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
DASP	75	PL 56B 491	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
ESPOSITO	75B	LNC 14 73	W. Braunschweig <i>et al.</i>	(DASP Collab.)
FORD	75	PRL 34 604	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
			R.L. Ford <i>et al.</i>	(SLAC, PENN)