

**$a_2(1700)$** 

$$I^G(J^{PC}) = 1^-(2^{++})$$

 **$a_2(1700)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1698 ± 44</b>		<sup>1</sup> AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1638.9 ± 2.3 <sup>+57.4</sup> <sub>-0.1</sub>		<sup>2</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
1722 ± 15 ± 67		<sup>3</sup> RODAS 19	JPAC	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
1681 <sup>+22</sup> <sub>-35</sub>	46M	<sup>4,5</sup> AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow$ $\pi^- \pi^+ \pi^- p$
1720 ± 10 ± 60		<sup>6</sup> JACKURA 18	JPAC	$\pi^- p \rightarrow \eta \pi^- p$
1726 ± 12 ± 25		<sup>5</sup> ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
1675 ± 25		ANISOVICH 09	RVUE	0.0 $\bar{p}p$ , $\pi N$
1722 ± 9 ± 15	18k	<sup>7</sup> SCHEGELSKY 06	RVUE	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$
1702 ± 7	80k	<sup>8</sup> UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
1721 ± 13 ± 44	145k	LU 05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1737 ± 5 ± 7		ABE 04	BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
1767 ± 14	221	<sup>9</sup> ACCIARRI 01H	L3	$\gamma \gamma \rightarrow K_S^0 K_S^0$ , $E_{cm}^{ee} =$ 91, 183–209 GeV
1660 ± 40		<sup>5</sup> ABELE 99B	CBAR	1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$
~ 1775		<sup>10</sup> GRYGOREV 99	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1752 ± 21 ± 4		ACCIARRI 97T	L3	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> T-matrix pole.<sup>2</sup> T-matrix pole, 2 poles, 2 channels ( $\pi \eta$ ,  $K \bar{K}$ ).<sup>3</sup> The coupled-channel analysis of both the  $\eta \pi$  and  $\eta' \pi$  systems using ADOLPH 15 data. The mass is extracted from the T-matrix pole.<sup>4</sup> Statistical error negligible.<sup>5</sup> Breit-Wigner mass.<sup>6</sup> Superseded by RODAS 19.<sup>7</sup> From analysis of L3 data at 183–209 GeV.<sup>8</sup> Statistical error only.<sup>9</sup> Spin 2 dominant, isospin not determined, could also be  $I=1$ .<sup>10</sup> Possibly two  $J^P = 2^+$  resonances with isospins 0 and 1. **$a_2(1700)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>265 ± 55</b>		<sup>1</sup> AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
224.0 ± 2.5 <sup>+1.8</sup> <sub>-48.3</sub>		<sup>2</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
247 ± 17 ± 63		<sup>3</sup> RODAS 19	JPAC	191 $\pi^- p \rightarrow \eta' \pi^- p$
436 <sup>+20</sup> <sub>-16</sub>	46M	<sup>4,5</sup> AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

280 ± 10 ± 70		<sup>6</sup> JACKURA	18	JPAC	$\pi^- p \rightarrow \eta \pi^- p$
190 ± 18 ± 30		<sup>5</sup> ABLIKIM	17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
270 + 50 - 20		ANISOVICH	09	RVUE	0.0 $\bar{p} p, \pi N$
336 ± 20 ± 20	18k	<sup>7</sup> SCHEGELSKY	06	RVUE	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$
417 ± 19	80k	<sup>8</sup> UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
279 ± 49 ± 66	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
151 ± 22 ± 24		ABE	04	BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
187 ± 60	221	<sup>9</sup> ACCIARRI	01H	L3	$\gamma \gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} =$ 91, 183–209 GeV
280 ± 70		<sup>5</sup> ABELE	99B	CBAR	1.94 $\bar{p} p \rightarrow \pi^0 \eta \eta$
150 ± 110 ± 34		ACCIARRI	97T	L3	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> T-matrix pole.

<sup>2</sup> T-matrix pole, 2 poles, 2 channels ( $\pi \eta, K \bar{K}$ ).

<sup>3</sup> The coupled-channel analysis of both the  $\eta \pi$  and  $\eta' \pi$  systems using ADOLPH 15 data. The width is extracted from the T-matrix pole.

<sup>4</sup> Statistical error negligible.

<sup>5</sup> Breit-Wigner width.

<sup>6</sup> Superseded by RODAS 19.

<sup>7</sup> From analysis of L3 data at 183–209 GeV.

<sup>8</sup> Statistical error only.

<sup>9</sup> Spin 2 dominant, isospin not determined, could also be  $I=1$ .

## $a_2(1700)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta \pi$	(3.6 ± 1.1) %
$\Gamma_2$ $\gamma \gamma$	(1.13 ± 0.30) × 10 <sup>-6</sup>
$\Gamma_3$ $\rho \pi$	seen
$\Gamma_4$ $f_2(1270) \pi$	seen
$\Gamma_5$ $K \bar{K}$	(1.9 ± 1.2) %
$\Gamma_6$ $\omega \pi^- \pi^0$	seen
$\Gamma_7$ $\omega \rho$	seen

## $a_2(1700)$ PARTIAL WIDTHS

$\Gamma(\eta \pi)$					$\Gamma_1$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>9.5 ± 2.0</b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

$\Gamma(\gamma \gamma)$					$\Gamma_2$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.30 ± 0.05</b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

$\Gamma(K\bar{K})$   $\Gamma_5$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$5.0 \pm 3.0$	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

$a_2(1700) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $(\Gamma_3 + \Gamma_4)\Gamma_2/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.29 \pm 0.04 \pm 0.02$		ACCIARRI 97T	L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37^{+0.12}_{-0.08} \pm 0.10$	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
---------------------------------	-----	----------------------------	------	--

<sup>1</sup> From analysis of L3 data at 183–209 GeV.

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_5\Gamma_2/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$20.6 \pm 4.2 \pm 4.6$	<sup>1</sup> ABE 04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
$49 \pm 11 \pm 13$	<sup>2</sup> ACCIARRI 01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{ee} = 91, 183\text{--}209 \text{ GeV}$

<sup>1</sup> Assuming spin 2.

<sup>2</sup> Spin 2 dominant, isospin not determined, could also be  $I=1$ .

$a_2(1700)$  BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$   $\Gamma_3/\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.4 \pm 0.4 \pm 0.1$	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
-----------------------	-----	----------------------------	------	--

<sup>1</sup> From analysis of L3 data at 183–209 GeV.

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$   $\Gamma_5/\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.134 \pm 0.106^{+4.909}_{-2.988}$	<sup>1</sup> ALBRECHT 20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
-------------------------------------	--------------------------	------	---

<sup>1</sup> Residues from T-matrix pole, 2 poles, 2 channels ( $\pi\eta, K\bar{K}$ ).

## $a_2(1700)$ REFERENCES

ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>	
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)
		Translated from YAF 62 513.		

---