

$f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Scalar Mesons below 2 GeV" and a note on "Non- $q\bar{q}$ Candidates" in PDG 06, Journal of Physics **G33** 1 (2006).

$f_0(1370)$ T-MATRIX POLE POSITION

Note that $\Gamma \approx 2 \text{Im}(\sqrt{s_{\text{pole}}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1200–1500)–i(150–250) OUR ESTIMATE			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$(1280.6 \pm 1.6 \pm 47.4)$ – $i(205.2 \pm 1.7 \pm 20.7)$	¹ ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
(1290 ± 50) – $i(170^{+20}_{-40})$	² ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
(1373 ± 15) – $i(137 \pm 10)$	³ BARGIOTTI	03	OBLX $\bar{p}p$
(1302 ± 17) – $i(166 \pm 18)$	⁴ BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_S$
$(1312 \pm 25 \pm 10)$ – $i(109 \pm 22 \pm 15)$	BARBERIS	99D	OMEG $450 pp \rightarrow K^+ K^-,$ $\pi^+ \pi^-$
(1406 ± 19) – $i(80 \pm 6)$	⁵ KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
(1300 ± 20) – $i(120 \pm 20)$	ANISOVICH	98B	RVUE Compilation
(1290 ± 15) – $i(145 \pm 15)$	BARBERIS	97B	OMEG $450 pp \rightarrow$ $pp2(\pi^+ \pi^-)$
(1548 ± 40) – $i(560 \pm 40)$	BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
(1380 ± 40) – $i(180 \pm 25)$	ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
(1300 ± 15) – $i(115 \pm 8)$	BUGG	96	RVUE
(1330 ± 50) – $i(150 \pm 40)$	⁶ AMSLER	95B	CBAR $\bar{p}p \rightarrow 3\pi^0$
(1360 ± 35) – $i(150-300)$	⁶ AMSLER	95C	CBAR $\bar{p}p \rightarrow \pi^0 \eta \eta$
(1390 ± 30) – $i(190 \pm 40)$	⁷ AMSLER	95D	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta,$ $\pi^0 \pi^0 \eta$
1346 – i 249	^{8,9} JANSSEN	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1214 – i 168	^{9,10} TORNVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
1364 – i 139	AMSLER	94D	CBAR $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
(1365^{+20}_{-55}) – $i(134 \pm 35)$	ANISOVICH	94	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$
(1340 ± 40) – $i(127^{+30}_{-20})$	¹¹ BUGG	94	RVUE $\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0,$ $\eta \pi^0 \pi^0$
(1430 ± 5) – $i(73 \pm 13)$	¹² KAMINSKI	94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1420 – i 220	¹³ AU	87	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

¹ T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$), and BINON 84C ($\eta\eta'$).

² Another pole is found at $(1510 \pm 130) - i(800^{+100}_{-150})$ MeV.

³ Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0,$ and $K^\pm K_S^0 \pi^\mp$.

⁴ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

⁵ T-matrix pole on sheet – – –.

⁶ Supersedes ANISOVICH 94.

⁷ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta,$ and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁸ Analysis of data from FALVARD 88.

⁹ The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

¹⁰ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

¹¹ Reanalysis of ANISOVICH 94 data.

¹² T-matrix pole on sheet III.

¹³ Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

$f_0(1370)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>			
1200 to 1500 OUR ESTIMATE					
$\pi\pi$ MODE					
<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1400 ± 40		¹ AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$	
1470 ⁺ ₋₇ ⁶⁺ ₋₂₅₅		² UEHARA	08A BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	
1259 ± 55	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^-\pi^+\pi^+$	
1309 ± 1 ± 15		³ BUGG	07A RVUE	0.0 $p\bar{p} \rightarrow 3\pi^0$	
1449 ± 13	4.3k	⁴ GARMASH	06 BELL	$B^+ \rightarrow K^+\pi^+\pi^-$	
1350 ± 50		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$	
1265 ± 30 ⁺ ₋₃₅		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$	
1434 ± 18 ± 9	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$	
1308 ± 10		BARBERIS	99B OMEG	450 $pp \rightarrow p_s p_f \pi^+\pi^-$	
1315 ± 50		BELLAZZINI	99 GAM4	450 $pp \rightarrow pp\pi^0\pi^0$	
1315 ± 30		ALDE	98 GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0n$	
1280 ± 55		BERTIN	98 OBLX	0.05–0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$	
1186		^{5,6} TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$	
1472 ± 12		ARMSTRONG	91 OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$	
1275 ± 20		BREAKSTONE	90 SFM	62 $pp \rightarrow pp\pi^+\pi^-$	
1420 ± 20		AKESSON	86 SPEC	63 $pp \rightarrow pp\pi^+\pi^-$	
1256		FROGGATT	77 RVUE	$\pi^+\pi^-$ channel	

¹ Breit-Wigner mass.

² Breit-Wigner mass. May also be the $f_0(1500)$.

³ Reanalysis of ABELE 96C data.

⁴ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0\pi^+\pi^-$ decays. Supersedes GARMASH 05.

⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁶ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1422 \pm 15 \pm 28$		¹ AAIJ	19H LHCb	$p\bar{p} \rightarrow D^\pm X$
$1360 \pm 31 \pm 28$	430	^{2,3} DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
$1350 \pm 48 \pm 15$	168	^{2,3} DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
1440 ± 6		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1391 ± 10		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440 ± 50		BOLONKIN	88 SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1463 ± 9		ETKIN	82B MPS	$23 \pi^- p \rightarrow n 2K_S^0$
1425 ± 15		WICKLUND	80 SPEC	$6 \pi N \rightarrow K^+ K^- N$
~ 1300		POLYCHRO...	79 STRC	$7 \pi^- p \rightarrow n 2K_S^0$

¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the isobar model A.² Using CLEO-c data but not authored by the CLEO Collaboration.³ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 346$ MeV. **4π MODE $2(\pi\pi)_S + \rho\rho$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1395 ± 40		ABELE	01 CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1374 ± 38		AMSLER	94 CBAR	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
1345 ± 12		ADAMO	93 OBLX	$\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
1386 ± 30		GASPERO	93 DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 1410	5751	¹ BETTINI	66 DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant. **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1262^{+51+82}_{-78-103}$	¹ UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1430	AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
1220 ± 40	ALDE	86D GAM4	$100 \pi^- p \rightarrow n 2\eta$

¹ Breit-Wigner mass. May also be the $f_0(1500)$.**COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1330.2^{+5.9}_{-6.5} \pm 5.1$	¹ AAIJ	19H LHCb	$p\bar{p} \rightarrow D^\pm X$
1306 ± 20	² ANISOVICH	03 RVUE	

¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUEDE 18.² K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

$f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)

DOCUMENT ID

200 to 500 OUR ESTIMATE **$\pi\pi$ MODE**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
300 ± 80		¹ AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
$90^+_{-1} \quad 2^+_{-1} \quad 50_{-22}$		² UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
298 ± 21	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
126 ± 25	4286	³ GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
265 ± 40		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$350 \pm 100^+_{-60} \quad 105_{-60}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
$173 \pm 32 \pm 6$	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
222 ± 20		BARBERIS	99B OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
255 ± 60		BELLAZZINI	99 GAM4	$450 pp \rightarrow pp \pi^0 \pi^0$
190 ± 50		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
323 ± 13		BERTIN	98 OBLX	$0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
350		^{4,5} TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195 ± 33		ARMSTRONG	91 OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
285 ± 60		BREAKSTONE	90 SFM	$62 pp \rightarrow pp\pi^+ \pi^-$
460 ± 50		AKESSON	86 SPEC	$63 pp \rightarrow pp\pi^+ \pi^-$
~ 400		⁶ FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

¹ The systematic errors are not reported.² Breit-Wigner width. May also be the $f_0(1500)$.³ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.⁴ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.⁵ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays⁶ Width defined as distance between 45 and 135° phase shift. **$K\bar{K}$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$324 \pm 38 \pm 42$	¹ AAIJ	19H LHCb	$pp \rightarrow D^\pm X$
121 ± 15	VLADIMIRSK..06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
55 ± 26	TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
250 ± 80	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$118^+_{-16} \quad 138_{-16}$	ETKIN	82B MPS	$23 \pi^- p \rightarrow n 2K_S^0$
160 ± 30	WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
~ 150	POLYCHRO...	79 STRC	$7 \pi^- p \rightarrow n 2K_S^0$

¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the isobar model A.

4π MODE 2(ππ)_S+ρρ

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • •				We do not use the following data for averages, fits, limits, etc. • • •
275 ± 55		ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 ± 61		AMSLER	94	CBAR 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
398 ± 26		ADAMO	93	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 ± 50		GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 90	5751	¹ BETTINI	66	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$
¹ ρρ dominant.				

ηη MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • •			We do not use the following data for averages, fits, limits, etc. • • •
484 ⁺²⁴⁶⁺²⁴⁶ ₋₁₇₀₋₂₆₃	¹ UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
250	AMSLER	92	CBAR 0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
320 ± 40	ALDE	86D GAM4	100 $\pi^-p \rightarrow n2\eta$
¹ Breit-Wigner width. May also be the $f_0(1500)$.			

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN	
• • •			We do not use the following data for averages, fits, limits, etc. • • •
147 ⁺³⁰ ₋₅₀	¹ ANISOVICH	03 RVUE	
¹ K-matrix pole from combined analysis of $\pi^-p \rightarrow \pi^0\pi^0n$, $\pi^-p \rightarrow K\bar{K}n$, $\pi^+\pi^- \rightarrow \pi^+\pi^-$, $\bar{p}p \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$, $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, $K_S^0K_S^0\pi^0$, $K^+K_S^0\pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+$, $K_S^0K^-\pi^0$, $K_S^0K_S^0\pi^-$ at rest.			

f₀(1370) DECAY MODES

Mode	Fraction (Γ _{<i>i</i>} /Γ)
Γ ₁ ππ	seen
Γ ₂ 4π	seen
Γ ₃ 4π ⁰	seen
Γ ₄ 2π ⁺ 2π ⁻	seen
Γ ₅ π ⁺ π ⁻ 2π ⁰	seen
Γ ₆ ρρ	seen
Γ ₇ 2(ππ) _S -wave	seen
Γ ₈ π(1300)π	seen
Γ ₉ a ₁ (1260)π	seen
Γ ₁₀ ηη	seen
Γ ₁₁ K \bar{K}	seen
Γ ₁₂ K \bar{K} nπ	not seen
Γ ₁₃ 6π	not seen
Γ ₁₄ ωω	not seen

Γ_{15}	$\gamma\gamma$	seen
Γ_{16}	e^+e^-	not seen

$f_0(1370)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$ Γ_{15}
 See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90.

$\Gamma(e^+e^-)$ Γ_{16}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^0\pi^0$

$f_0(1370)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{10}\Gamma_{15}/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$121^{+133}_{-53} + 169_{-106}$	¹ UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

¹ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

$f_0(1370)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.10	95	OCHS	13	RVUE
0.26 ± 0.09		BUGG	96	RVUE
<0.15		¹ AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
<0.06		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$

¹ Using AMSLER 95B ($3\pi^0$).

$\Gamma(4\pi)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
>0.72	GASPERO 93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$

$\Gamma(4\pi^0)/\Gamma(4\pi)$ Γ_3/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	ABELE 96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
0.068 ± 0.005	¹ GASPERO 93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$

¹ Model-dependent evaluation.

$\Gamma(2\pi^+2\pi^-)/\Gamma(4\pi)$ $\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3+\Gamma_4+\Gamma_5)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.420±0.014	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

¹ Model-dependent evaluation.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(4\pi)$ $\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3+\Gamma_4+\Gamma_5)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.512±0.019	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

¹ Model-dependent evaluation.

$\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_6/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.26±0.07	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(\pi\pi)$ Γ_7/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.6±2.6	¹ ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^-4\pi^0 p$

¹ From the combined data of ABELE 96 and ABELE 96C.

$\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$ Γ_7/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.51±0.09	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$ Γ_6/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
large	BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
1.6 ±0.2	AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
~ 0.65	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_8/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.17±0.06	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_9/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.06±0.02	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(\eta\eta)/\Gamma(4\pi)$ **$\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3+\Gamma_4+\Gamma_5)$**

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(28 \pm 11) \times 10^{-3}$	¹ ANISOVICH	02D	SPEC	Combined fit
$(4.7 \pm 2.0) \times 10^{-3}$	BARBERIS	00E		450 $p\bar{p} \rightarrow p_f \eta \eta p_S$

¹From a combined K-matrix analysis of Crystal Barrel ($0. \rho\bar{\rho} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

VALUE DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.35 ± 0.13	BUGG	96	RVUE
-----------------	------	----	------

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ **Γ_{11}/Γ_1**

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08 ± 0.08	ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-, \phi K^+ K^-$
0.91 ± 0.20	¹ BARGIOTTI	03	OBLX	$\bar{p}p$
0.12 ± 0.06	² ANISOVICH	02D	SPEC	Combined fit
$0.46 \pm 0.15 \pm 0.11$	BARBERIS	99D	OMEG	450 $p\bar{p} \rightarrow K^+ K^-, \pi^+ \pi^-$

¹Coupled channel analysis of $\pi^+\pi^-\pi^0, K^+K^-\pi^0$, and $K^\pm K_S^0 \pi^\mp$.

²From a combined K-matrix analysis of Crystal Barrel ($0. \rho\bar{\rho} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

$\Gamma(K\bar{K}n\pi)/\Gamma_{\text{total}}$ **Γ_{12}/Γ**

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.03	GASPERO	93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$
---------	---------	----	-----	---

$\Gamma(6\pi)/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.22	GASPERO	93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$
---------	---------	----	-----	---

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13	GASPERO	93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$
---------	---------	----	-----	---

$f_0(1370)$ REFERENCES

- ALBRECHT 20 EPJ C80 453 M. Albrecht *et al.* (Crystal Barrel Collab.)
AAIJ 19H JHEP 1904 063 R. Aaij *et al.* (LHCb Collab.)
AOUDE 18 PR D98 056021 R.T. Aoude *et al.*
DOBBS 15 PR D91 052006 S. Dobbs *et al.* (NWES)
OCHS 13 JP G40 043001 W. Ochs
UEHARA 10A PR D82 114031 S. Uehara *et al.* (BELLE Collab.)
ANISOVICH 09 IJMP A24 2481 V.V. Anisovich, A.V. Sarantsev
AUBERT 09L PR D79 072006 B. Aubert *et al.* (BABAR Collab.)
PDG 08 PL B667 1 C. Amsler *et al.* (PDG Collab.)
UEHARA 08A PR D78 052004 S. Uehara *et al.* (BELLE Collab.)
BONVICINI 07 PR D76 012001 G. Bonvicini *et al.* (CLEO Collab.)
BUGG 07A JP G34 151 D.V. Bugg *et al.*
GARMASH 07 PR D75 012006 A. Garmash *et al.* (BELLE Collab.)
GARMASH 06 PRL 96 251803 A. Garmash *et al.* (BELLE Collab.)
PDG 06 JP G33 1 W.-M. Yao *et al.* (PDG Collab.)
VLADIMIRSK... 06 PAN 69 493 V.V. Vladimirovsky *et al.* (ITEP, Moscow)
Translated from YAF 69 515.
ABLIKIM 05 PL B607 243 M. Ablikim *et al.* (BES Collab.)
ABLIKIM 05Q PR D72 092002 M. Ablikim *et al.* (BES Collab.)
GARMASH 05 PR D71 092003 A. Garmash *et al.* (BELLE Collab.)
ANISOVICH 03 EPJ A16 229 V.V. Anisovich *et al.*
BARGIOTTI 03 EPJ C26 371 M. Bargiotti *et al.* (OBELIX Collab.)
TIKHOMIROV 03 PAN 66 828 G.D. Tikhomirov *et al.*
Translated from YAF 66 860.
ANISOVICH 02D PAN 65 1545 V.V. Anisovich *et al.*
Translated from YAF 65 1583.
ABELE 01 EPJ C19 667 A. Abele *et al.* (Crystal Barrel Collab.)
ABELE 01B EPJ C21 261 A. Abele *et al.* (Crystal Barrel Collab.)
AITALA 01A PRL 86 765 E.M. Aitala *et al.* (FNAL E791 Collab.)
ASNER 00 PR D61 012002 D.M. Asner *et al.* (CLEO Collab.)
BARBERIS 00C PL B471 440 D. Barberis *et al.* (WA 102 Collab.)
BARBERIS 00E PL B479 59 D. Barberis *et al.* (WA 102 Collab.)
BARBERIS 99B PL B453 316 D. Barberis *et al.* (Omega Expt.)
BARBERIS 99D PL B462 462 D. Barberis *et al.* (Omega Expt.)
BELLAZZINI 99 PL B467 296 R. Bellazzini *et al.*
KAMINSKI 99 EPJ C9 141 R. Kaminski, L. Lesniak, B. Loiseau (CRAC, PARIN)
ALDE 98 EPJ A3 361 D. Alde *et al.* (GAM4 Collab.)
Also PAN 62 405 D. Alde *et al.* (GAMS Collab.)
Translated from YAF 62 446.
ANISOVICH 98B SPU 41 419 V.V. Anisovich *et al.*
Translated from UFN 168 481.
BERTIN 98 PR D57 55 A. Bertin *et al.* (OBELIX Collab.)
BARBERIS 97B PL B413 217 D. Barberis *et al.* (WA 102 Collab.)
BERTIN 97C PL B408 476 A. Bertin *et al.* (OBELIX Collab.)
ABELE 96 PL B380 453 A. Abele *et al.* (Crystal Barrel Collab.)
ABELE 96B PL B385 425 A. Abele *et al.* (Crystal Barrel Collab.)
ABELE 96C NP A609 562 A. Abele *et al.* (Crystal Barrel Collab.)
BUGG 96 NP B471 59 D.V. Bugg, A.V. Sarantsev, B.S. Zou (LOQM, PNPI)
AMSLER 95B PL B342 433 C. Amsler *et al.* (Crystal Barrel Collab.)
AMSLER 95C PL B353 571 C. Amsler *et al.* (Crystal Barrel Collab.)
AMSLER 95D PL B355 425 C. Amsler *et al.* (Crystal Barrel Collab.)
JANSSEN 95 PR D52 2690 G. Janssen *et al.* (STON, ADLD, JULI)
TORNQVIST 95 ZPHY C68 647 N.A. Tornqvist (HELs)
AMSLER 94 PL B322 431 C. Amsler *et al.* (Crystal Barrel Collab.) JPC
AMSLER 94D PL B333 277 C. Amsler *et al.* (Crystal Barrel Collab.)
ANISOVICH 94 PL B323 233 V.V. Anisovich *et al.* (Crystal Barrel Collab.) JPC
BUGG 94 PR D50 4412 D.V. Bugg *et al.* (LOQM)
KAMINSKI 94 PR D50 3145 R. Kaminski, L. Lesniak, J.P. Maillet (CRAC+)
ADAMO 93 NP A558 13C A. Adamo *et al.* (OBELIX Collab.) JPC
GASPERO 93 NP A562 407 M. Gaspero (ROMAI) JPC
AMSLER 92 PL B291 347 C. Amsler *et al.* (Crystal Barrel Collab.)
ARMSTRONG 91 ZPHY C51 351 T.A. Armstrong *et al.* (ATHU, BARI, BIRM+)
ARMSTRONG 91B ZPHY C52 389 T.A. Armstrong *et al.* (ATHU, BARI, BIRM+)
BREAKSTONE 90 ZPHY C48 569 A.M. Breakstone *et al.* (ISU, BGNA, CERN+)
MORGAN 90 ZPHY C48 623 D. Morgan, M.R. Pennington (RAL, DURH)
ASTON 88 NP B296 493 D. Aston *et al.* (SLAC, NAGO, CINC, INUS)
BOLONKIN 88 NP B309 426 B.V. Bolonkin *et al.* (ITEP, SERP)
FALVARD 88 PR D38 2706 A. Falvard *et al.* (CLER, FRAS, LALO+)
VOROBYEV 88 SJNP 48 273 P.V. Vorobiev *et al.* (NOVO)
Translated from YAF 48 436.

AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)
