

$\Lambda(1670) \ 1/2^-$  $I(J^P) = 0(\frac{1}{2}^-)$  Status: \*\*\*\*

The measurements of the mass, width, and elasticity published before 1974 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** 1 (1982).

 **$\Lambda(1670)$  POLE POSITIONS****REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1670 to 1678 (<math>\approx 1674</math>) OUR ESTIMATE</b>			
1676 $\pm 2$	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
1669 $\begin{smallmatrix} +3 \\ -8 \end{smallmatrix}$	<sup>1</sup> KAMANO	15	DPWA $\bar{K}N$ multichannel
1677.5 $\pm 0.8$	GARCIA-REC...03		DPWA $\bar{K}N$ multichannel
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1667	ZHANG	13A	DPWA $\bar{K}N$ multichannel

<sup>1</sup>From the preferred solution A in KAMANO 15.

**-2×IMAGINARY PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>28 to 36 (<math>\approx 32</math>) OUR ESTIMATE</b>			
33 $\pm 4$	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
19 $\begin{smallmatrix} +18 \\ -2 \end{smallmatrix}$	<sup>1</sup> KAMANO	15	DPWA $\bar{K}N$ multichannel
29.2 $\pm 1.4$	GARCIA-REC...03		DPWA $\bar{K}N$ multichannel
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
26	ZHANG	13A	DPWA $\bar{K}N$ multichannel

<sup>1</sup>From the preferred solution A in KAMANO 15.

 **$\Lambda(1670)$  POLE RESIDUES**

The normalized residue is the residue divided by  $\Gamma_{pole}/2$ .

**Normalized residue in  $\bar{K}N \rightarrow \Lambda(1670) \rightarrow \bar{K}N$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.30 <math>\pm 0.06</math></b>	<b>-145 <math>\pm 11</math></b>	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.351	164	<sup>1</sup> KAMANO	15	DPWA $\bar{K}N$ multichannel

<sup>1</sup>From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma\pi$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.19 <math>\pm 0.06</math></b>	<b>145 <math>\pm 14</math></b>	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.327	125	<sup>1</sup> KAMANO	15	DPWA $\bar{K}N$ multichannel

<sup>1</sup>From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda\eta$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.26 <math>\pm</math> 0.09</b>	<b>104 <math>\pm</math> 14</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.474	59	<sup>1</sup> KAMANO 15	DPWA	Multichannel

<sup>1</sup>From the preferred solution A in KAMANO 15.**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Xi K$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.02 <math>\pm</math> 0.02</b>	<b>100 <math>\pm</math> 25</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda\omega, S=1/2, S\text{-wave}$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.09 <math>\pm</math> 0.04</b>	<b>-60 <math>\pm</math> 35</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda\omega, S=3/2, D\text{-wave}$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.05 <math>\pm</math> 0.04</b>		SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow N\bar{K}^*(892), S=1/2, S\text{-wave}$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.31 <math>\pm</math> 0.14</b>	<b>100 <math>\pm</math> 45</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow N\bar{K}^*(892), S=3/2, D\text{-wave}$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.06 <math>\pm</math> 0.03</b>	<b>-85 <math>\pm</math> 40</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda\sigma$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.25 <math>\pm</math> 0.08</b>	<b>160 <math>\pm</math> 15</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma(1385)\pi$** 

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.13 <math>\pm</math> 0.06</b>	<b>110 <math>\pm</math> 12</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0988	-104	<sup>1</sup> KAMANO 15	DPWA	Multichannel

<sup>1</sup>From the preferred solution A in KAMANO 15. **$\Lambda(1670)$  MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1670 to 1678 (<math>\approx</math> 1674) OUR ESTIMATE</b>			
1677 $\pm$ 2	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
1672 $\pm$ 3	ZHANG 13A	DPWA	Multichannel
1670.8 $\pm$ 1.7	KOISO 85	DPWA	$K^- p \rightarrow \Sigma \pi$
1667 $\pm$ 5	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1671 $\pm$ 3	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1675 $\pm$ 2	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma \pi$
1679 $\pm$ 1	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$
1665 $\pm$ 5	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1673 ± 2	MANLEY	02	DPWA	$\bar{K}N$ multichannel
1668.9 ± 2.0	ABAEV	96	DPWA	$K^- p \rightarrow \Lambda \eta$
1670 ± 5	GOPAL	77	DPWA	$\bar{K}N$ multichannel
1664	<sup>1</sup> MARTIN	77	DPWA	$\bar{K}N$ multichannel

<sup>1</sup>MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.

### $\Lambda(1670)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>25 to 35 (≈ 30) OUR ESTIMATE</b>			
33 ± 4	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
29 ± 5	ZHANG	13A	DPWA $\bar{K}N$ multichannel
34.1 ± 3.7	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
29 ± 5	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
29 ± 5	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
46 ± 5	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
40 ± 3	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
19 ± 5	PREVOST	74	DPWA $K^- N \rightarrow \Sigma(1385)\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

23 ± 6	MANLEY	02	DPWA	$\bar{K}N$ multichannel
21.1 ± 3.6	ABAEV	96	DPWA	$K^- p \rightarrow \Lambda \eta$
45 ± 10	GOPAL	77	DPWA	$\bar{K}N$ multichannel
12	<sup>1</sup> MARTIN	77	DPWA	$\bar{K}N$ multichannel

<sup>1</sup>MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.

### $\Lambda(1670)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $N\bar{K}$	20–30 %
$\Gamma_2$ $\Sigma \pi$	25–55 %
$\Gamma_3$ $\Lambda \eta$	10–25 %
$\Gamma_4$ $\Sigma(1385)\pi$ , <i>D</i> -wave	( 6.0 ± 2.0 ) %
$\Gamma_5$ $N\bar{K}^*(892)$ , <i>S</i> =1/2, <i>S</i> -wave	
$\Gamma_6$ $N\bar{K}^*(892)$ , <i>S</i> =3/2, <i>D</i> -wave	( 5 ± 4 ) %
$\Gamma_7$ $\Lambda \sigma$	( 20 ± 8 ) %

### $\Lambda(1670)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
<b>0.20 to 0.30 OUR ESTIMATE</b>				
0.33 ± 0.07	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
0.26 ± 0.25	ZHANG	13A	DPWA	$\bar{K}N$ multichannel

0.18 ±0.03	GOPAL	80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
0.17 ±0.03	ALSTON-...	78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.318	<sup>1</sup> KAMANO	15	DPWA	$\bar{K}N$ multichannel
0.37 ±0.07	MANLEY	02	DPWA	$\bar{K}N$ multichannel
0.20 ±0.03	GOPAL	77	DPWA	See GOPAL 80
0.15	<sup>2</sup> MARTIN	77	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

<sup>2</sup> MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.

### $\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.12 ±0.03</b>	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.289	<sup>1</sup> KAMANO	15	DPWA Multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Lambda\eta)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.20 ±0.08</b>	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.373	KAMANO	15	DPWA Multichannel
0.30 ±0.08	ABAEV	96	DPWA $K^- p \rightarrow \Lambda\eta$

### $\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.06 ±0.02</b>	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.019	KAMANO	15	DPWA Multi-channel

### $\Gamma(\Lambda\sigma)/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.20 ±0.08</b>	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

### $\Gamma(N\bar{K}^*(892), S=1/2, S\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	<sup>1</sup> KAMANO	15	DPWA Multichannel

<sup>1</sup> Not seen in the preferred solution A in KAMANO 15.

### $\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.05 ±0.04</b>	ZHANG	13A	DPWA Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	<sup>1</sup> KAMANO	15	DPWA Multichannel

<sup>1</sup> Not seen in the preferred solution A in KAMANO 15.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma \pi$	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
$-0.29 \pm 0.06$	ZHANG	13A	DPWA Multichannel
$-0.26 \pm 0.02$	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
$-0.31 \pm 0.03$	GOPAL	77	DPWA $\bar{K} N$ multichannel
$-0.29 \pm 0.03$	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
$-0.23 \pm 0.03$	LONDON	75	HLBC $K^- p \rightarrow \Sigma^0 \pi^0$
$-0.27 \pm 0.02$	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$-0.38 \pm 0.03$	MANLEY	02	DPWA $\bar{K} N$ multichannel
$-0.13$	<sup>1</sup> MARTIN	77	DPWA $\bar{K} N$ multichannel
<sup>1</sup> MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.			

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda \eta$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
$-0.30 \pm 0.10$	ZHANG	13A	DPWA Multichannel
$+0.20 \pm 0.05$	BAXTER	73	DPWA $K^- p \rightarrow$ neutrals
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$+0.24 \pm 0.04$	MANLEY	02	DPWA $\bar{K} N$ multichannel
0.24	KIM	71	DPWA K-matrix analysis
0.26	ARMENTEROS69C	HBC	
0.20 or 0.23	BERLEY	65	HBC

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma(1385)\pi$ , D-wave	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
$-0.17 \pm 0.06$	MANLEY	02	DPWA $\bar{K} N$ multichannel
$-0.18 \pm 0.05$	PREVOST	74	DPWA $K^- N \rightarrow \Sigma(1385)\pi$

### $\Lambda(1670)$ REFERENCES

SARANTSEV 19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
KAMANO 15	PR C92 025205	H. Kamano <i>et al.</i>	(ANL, OSAK)
ZHANG 13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
GARCIA-REC... 03	PR D67 076009	C. Garcia-Recio <i>et al.</i>	(GRAN, VALE)
MANLEY 02	PRL 88 012002	D.M. Manley <i>et al.</i>	(BNL Crystal Ball Collab.)
ABAEV 96	PR C53 385	V.V. Abaev, B.M.K. Nefkens	(UCLA)
KOISO 85	NP A433 619	H. Koiso <i>et al.</i>	(TOKY, MASA)
PDG 82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL 80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
ALSTON-... 78	PR D18 182	M. Alston-Garnjost <i>et al.</i>	(LBL, MTHO+) IJP
Also	PRL 38 1007	M. Alston-Garnjost <i>et al.</i>	(LBL, MTHO+) IJP
GOPAL 77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
MARTIN 77	NP B127 349	B.R. Martin, M.K. Pidcock, R.G. Moorhouse	(LOUC+) IJP
Also	NP B126 266	B.R. Martin, M.K. Pidcock	(LOUC)
Also	NP B126 285	B.R. Martin, M.K. Pidcock	(LOUC) IJP
HEPP 76B	PL 65B 487	V. Hepp <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
LONDON 75	NP B85 289	G.W. London <i>et al.</i>	(BNL, CERN, EPOL+)
KANE 74	LBL-2452	D.F. Kane	(LBL) IJP
PREVOST 74	NP B69 246	J. Prevost <i>et al.</i>	(SACL, CERN, HEID)
BAXTER 73	NP B67 125	D.F. Baxter <i>et al.</i>	(OXF) IJP
KIM 71	PRL 27 356	J.K. Kim	(HARV) IJP
Also	Duke Conf. 161	J.K. Kim	(HARV) IJP
Hyperon Resonances, 1970			

ARMENTEROS 69C Lund Paper 229 R. Armenteros *et al.* (CERN, HEID, SACL) IJP  
Values are quoted in LEVI-SETTI 69.  
BERLEY 65 PRL 15 641 D. Berley *et al.* (BNL) IJP

---