

$\psi(4160)$

$$J^{PC} = 0^{-}(1^{-}-)$$

$\psi(4160)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4191 ± 5 OUR AVERAGE			
4191 $\begin{smallmatrix} +9 \\ -8 \end{smallmatrix}$	AAIJ	13BC	LHCB $B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 ± 6.5	¹ ABLIKIM	08D	BES2 $e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
4193 ± 7	² MO	10	RVUE $e^+ e^- \rightarrow$ hadrons
4151 ± 4	³ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
4155 ± 5	⁴ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C	DASP $e^+ e^-$

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4160)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70 ± 10 OUR AVERAGE			
65 $\begin{smallmatrix} +22 \\ -16 \end{smallmatrix}$	AAIJ	13BC	LHCB $B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8 ± 12.3	¹ ABLIKIM	08D	BES2 $e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
79 ± 14	² MO	10	RVUE $e^+ e^- \rightarrow$ hadrons
107 ± 10	³ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
107 ± 16	⁴ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
78 ± 20	BRANDELIK	78C	DASP $e^+ e^-$

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4160)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$	
Γ_2 $\mu^+ \mu^-$	seen	
Γ_3 $D \bar{D}$	seen	
Γ_4 $D^0 \bar{D}^0$	seen	
Γ_5 $D^+ D^-$	seen	
Γ_6 $D^* \bar{D} + \text{c.c.}$	seen	
Γ_7 $D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen	
Γ_8 $D^*(2010)^+ D^- + \text{c.c.}$	seen	
Γ_9 $D^* \bar{D}^*$	seen	
Γ_{10} $D^*(2007)^0 \bar{D}^*(2007)^0$	seen	
Γ_{11} $D^*(2010)^+ D^*(2010)^-$	seen	
Γ_{12} $D^0 D^- \pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0 \bar{D}^0 + \text{c.c.}$, $D^*(2010)^+ D^- + \text{c.c.}$)	not seen	
Γ_{13} $D \bar{D}^* \pi + \text{c.c.}$ (excl. $D^* \bar{D}^*$)	seen	
Γ_{14} $D^0 D^{*-} \pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+ D^*(2010)^-$)	not seen	
Γ_{15} $D_s^+ D_s^-$	not seen	
Γ_{16} $D_s^{*+} D_s^- + \text{c.c.}$	seen	
Γ_{17} $J/\psi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
Γ_{18} $J/\psi \pi^0 \pi^0$	$< 3 \times 10^{-3}$	90%
Γ_{19} $J/\psi K^+ K^-$	$< 2 \times 10^{-3}$	90%
Γ_{20} $J/\psi \eta$	$< 8 \times 10^{-3}$	90%
Γ_{21} $J/\psi \pi^0$	$< 1 \times 10^{-3}$	90%
Γ_{22} $J/\psi \eta'$	$< 5 \times 10^{-3}$	90%
Γ_{23} $J/\psi \pi^+ \pi^- \pi^0$	$< 1 \times 10^{-3}$	90%
Γ_{24} $\psi(2S) \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
Γ_{25} $\chi_{c1} \gamma$	$< 5 \times 10^{-3}$	90%
Γ_{26} $\chi_{c2} \gamma$	$< 1.3 \%$	90%
Γ_{27} $\chi_{c1} \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%
Γ_{28} $\chi_{c2} \pi^+ \pi^- \pi^0$	$< 8 \times 10^{-3}$	90%
Γ_{29} $h_c(1P) \pi^+ \pi^-$	$< 5 \times 10^{-3}$	90%
Γ_{30} $h_c(1P) \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%
Γ_{31} $h_c(1P) \eta$	$< 2 \times 10^{-3}$	90%
Γ_{32} $h_c(1P) \pi^0$	$< 4 \times 10^{-4}$	90%
Γ_{33} $\phi \pi^+ \pi^-$	$< 2 \times 10^{-3}$	90%

Γ_{34}	$\gamma\chi_{c1}(3872)$	< 1.8	$\times 10^{-3}$	90%
Γ_{35}	$\gamma X(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.36	$\times 10^{-4}$	90%
Γ_{36}	$\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.18	$\times 10^{-4}$	90%
Γ_{37}	$\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.47	$\times 10^{-4}$	90%
Γ_{38}	$\gamma X(3915) \rightarrow \gamma\gamma J/\psi$	< 1.26	$\times 10^{-4}$	90%
Γ_{39}	$\gamma X(3930) \rightarrow \gamma\gamma J/\psi$	< 8.8	$\times 10^{-5}$	90%
Γ_{40}	$\gamma X(3940) \rightarrow \gamma\gamma J/\psi$	< 1.79	$\times 10^{-4}$	90%
Γ_{41}	$K^+ K^-$			
Γ_{42}	$K_S^0 K^\pm \pi^\mp$			
Γ_{43}	$p\bar{p}p\bar{p}$	not seen		

$\psi(4160)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.48±0.22	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.4 to 1.1	² MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
0.83±0.08	³ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.84±0.13	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.77±0.23	BRANDELIK	78C DASP	e^+e^-

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\Gamma(\mu^+\mu^-)$

Γ_2

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.45±1.24±0.94	^{1,2} ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

¹ From a fit to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$.

² From solution 1 of 8 with equal fit quality. Other solutions range from $2.08 \pm 0.99 \pm 0.80$ to $2.45 \pm 1.24 \pm 0.94$ keV.

$\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi\eta')$ $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_{22}\Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17±0.04	86	^{1,2} ABLIKIM	20A BES3	$e^+e^- \rightarrow \eta' J/\psi$
1.07±0.09	86	^{1,3} ABLIKIM	20A BES3	$e^+e^- \rightarrow \eta' J/\psi$

¹ Based on a fit to $\sigma(e^+e^- \rightarrow \eta' J/\psi)$ from $\sqrt{s} = 4.18$ to 4.60 GeV assuming interfering $\psi(4160)$ and $\psi(4260)$ contributions. At $\sqrt{s} = 4.18$ GeV, $\sigma(e^+e^- \rightarrow \eta' J/\psi) = 2.4 \pm 0.3 \pm 0.2$ pb.

² Solution I of the fit, corresponding to a phase of -0.03 ± 0.44 rad.

³ Solution II of the fit, corresponding to a phase of 2.54 ± 0.04 rad.

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	¹ HAN	15 BELL	10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.1	90	¹ HAN	15 BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{42}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
2.71 ± 0.13 ± 0.12	¹ ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0095 ± 0.0088 ± 0.0004	² ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

¹ Solution I of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

² Solution II of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

$\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.9 ± 0.9	¹ WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$
12.8 ± 1.7 ± 2.0	² WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

¹ Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

² Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

$\psi(4160)$ BRANCHING RATIOS

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$

¹ AAIJ 13BC report $B(B^+ \rightarrow K^+ \psi(4160)) B(\psi(4160) \rightarrow \mu^+ \mu^-) = (3.5_{-0.8}^{+0.9}) \times 10^{-9}$.

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$				Γ_3/Γ_9
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.02 \pm 0.03 \pm 0.02$	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$				Γ_4/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^0\bar{D}^0$
seen	PAKHLOVA	08	BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$				Γ_5/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^+D^-$
seen	PAKHLOVA	08	BELL	$e^+e^- \rightarrow D^+D^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^+D^-\gamma$

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_7/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*0}\bar{D}^0$

$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_8/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	¹ ZHUKOVA	18	BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*+}D^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA	07	BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$
¹ Supersedes PAKHLOVA 07.				

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$				Γ_6/Γ_9
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.34 \pm 0.14 \pm 0.05$	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$				Γ_{10}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$

$\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$				Γ_{11}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	¹ ZHUKOVA	18	BELL	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*+}D^{*-}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA	07	BELL	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
¹ Supersedes PAKHLOVA 07.				

$$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.))} / \Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

$$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)) / \Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

$$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)) / \Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

$$\Gamma(D_s^+ D_s^-) / \Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

$$\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$

$$\Gamma(J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$$\Gamma(J/\psi \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$$\Gamma(J/\psi K^+ K^-) / \Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$$\Gamma(J/\psi \eta) / \Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	¹ ABLIKIM	15L	BES3	$e^+ e^- \rightarrow J/\psi \eta$
seen	WANG	13B	BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

¹ An enhancement around 4.2 GeV is observed.

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(J/\psi\eta')/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<13	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

¹At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	¹ PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

¹At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$ **Γ_{31}/Γ**

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2	90		¹ PEDLAR 11	CLEO	$e^+e^- \rightarrow h_c(1P)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 41 ² ABLIKIM 17R BES3 $e^+e^- \rightarrow h_c(1P)\eta$

¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

² An enhancement around 4.2 GeV is observed.

$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$ **Γ_{32}/Γ**

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	¹ PEDLAR 11	CLEO	$e^+e^- \rightarrow h_c(1P)\pi^0$

¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{33}/Γ**

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$ **Γ_{34}/Γ**

VALUE	CL%	DOCUMENT ID	COMMENT
<1.8 $\times 10^{-3}$	90	^{1,2} XIAO 13	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.012 90 ^{1,3} XIAO 13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

² XIAO 13 reports $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 0.68 \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.8 \times 10^{-2}$.

³ XIAO 13 reports $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 1.05 \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 8 \times 10^{-3}$.

$\Gamma(\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{35}/Γ**

VALUE	CL%	DOCUMENT ID	COMMENT
<1.36 $\times 10^{-4}$	90	¹ XIAO 13	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{36}/Γ**

VALUE	CL%	DOCUMENT ID	COMMENT
<1.18 $\times 10^{-4}$	90	¹ XIAO 13	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{37}/Γ**

VALUE	CL%	DOCUMENT ID	COMMENT
<1.47 $\times 10^{-4}$	90	¹ XIAO 13	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3915) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.26 \times 10^{-4}$	90	¹ XIAO 13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration. $\Gamma(\gamma X(3930) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<0.88 \times 10^{-4}$	90	¹ XIAO 13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration. $\Gamma(\gamma X(3940) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
$<1.79 \times 10^{-4}$	90	¹ XIAO 13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration. $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2 \times 10^{-5}$	90	¹ DRUZHININ 15	RVUE	$e^+ e^- \rightarrow \psi(3770)$
---------------------	----	---------------------------	------	----------------------------------

¹ DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+ e^- \rightarrow K^+ K^-$ and $e^+ e^- \rightarrow K_S^0 K_L^0$. $\Gamma(p\bar{p}p\bar{p})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 21D	BES3	4.0–4.6 $e^+ e^- \rightarrow p\bar{p}p\bar{p}$

 $\psi(4160)$ REFERENCES

ABLIKIM 21D	PR D103 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA 18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM 17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GAO 17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan	
PDG 16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM 15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DRUZHININ 15	PR D92 054024	V.P. Druzhinin	(NOVO)
HAN 15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
AAIJ 13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG 13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
XIAO 13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)
PAKHLOVA 11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR 11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA... 10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO 10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT 09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN... 09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA 09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM 08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA 08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN 06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH 05A	PR D72 017501	K.K. Seth	
BAI 02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)

BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
