

**$\rho(1570)$** 

$$I^G(J^{PC}) = 1^+(1^{--})$$

## OMITTED FROM SUMMARY TABLE

May be an OZI-violating decay mode of  $\rho(1700)$ . See our mini-review under the  $\rho(1700)$ .

 **$\rho(1570)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1570 \pm 36 \pm 62</math></b>	54	<sup>1</sup> AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1585 ± 15		<sup>2</sup> ACHASOV	20C SND	$1.3\text{--}2.0 e^+ e^- \rightarrow K^+ K^- \pi^0$
1480 ± 40		<sup>3</sup> BITYUKOV	87 SPEC	$32.5 \pi^- p \rightarrow \phi \pi^0 n$

<sup>1</sup> From the fit with two resonances.<sup>2</sup> From a fit using a two resonance model in which the mass and width of the other resonance are fixed at the  $\rho(1700)$  values from PDG 20.<sup>3</sup> Systematic errors not estimated. **$\rho(1570)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>144 \pm 75 \pm 43</math></b>	54	<sup>4</sup> AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
75 ± 30		<sup>5</sup> ACHASOV	20C SND	$1.3\text{--}2.0 e^+ e^- \rightarrow K^+ K^- \pi^0$
130 ± 60		<sup>6</sup> BITYUKOV	87 SPEC	$32.5 \pi^- p \rightarrow \phi \pi^0 n$

<sup>4</sup> From the fit with two resonances.<sup>5</sup> From a fit using a two resonance model in which the mass and width of the other resonance are fixed at the  $\rho(1700)$  values from PDG 20.<sup>6</sup> Systematic errors not estimated. **$\rho(1570)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	
$\Gamma_2$ $\phi \pi$	not seen
$\Gamma_3$ $\omega \pi$	

 **$\rho(1570)$   $\Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$** 

$\Gamma(\phi \pi) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_2 \Gamma_1/\Gamma$	
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.5 \pm 0.9 \pm 0.3</math></b>		54	<sup>7</sup> AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<70	90		<sup>8</sup> AULCHENKO	87B ND	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0$

<sup>7</sup> From the fit with two resonances.<sup>8</sup> Using mass and width of BITYUKOV 87.

## $\rho(1570)$ BRANCHING RATIOS

$\Gamma(\phi\pi)/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>not seen</b>	ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01 <sup>9</sup> DONNACHIE 91 RVUE

<sup>9</sup> Using data from BISELLO 91B, DOLINSKY 86, and ALBRECHT 87L.

$\Gamma(\phi\pi)/\Gamma(\omega\pi)$				$\Gamma_2/\Gamma_3$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>0.5	95	BITYUKOV 87	SPEC	$32.5 \pi^- p \rightarrow \phi \pi^0 n$

## $\rho(1570)$ REFERENCES

ACHASOV	20C	EPJ C80 1139	M.N. Achasov <i>et al.</i>	
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(SND Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(PDG Collab.)
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(BABAR Collab.)
BISELLO	91B	NPBPS B21 111	D. Bisello	(Crystal Barrel Collab.)
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(DM2 Collab.)
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(MCHS, LANC)
AULCHENKO	87B	JETPL 45 145	V.M. Aulchenko <i>et al.</i>	(ARGUS Collab.)
		Translated from ZETFP 45 118.		(NOVO)
BITYUKOV	87	PL B188 383	S.I. Bityukov <i>et al.</i>	(SERP)
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)