

$\Upsilon(11020)$ 

$$J^{PC} = 0^{-}(1^{- -})$$

 **$\Upsilon(11020)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>11000 ± 4 OUR AVERAGE</b>			
11000.0 <sup>+4.0</sup> <sub>-4.5</sub> ± 1.0 <sup>+1.0</sup> <sub>-1.3</sub>	<sup>1</sup> MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10999.0 <sup>+7.3</sup> <sub>-7.8</sub> ± 16.9 <sup>+16.9</sup> <sub>-1.0</sub>	<sup>2</sup> MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
11001 ± 1	<sup>3</sup> DONG	20A	$e^+e^- \rightarrow b\bar{b}$
11003.0 ± 1.1 <sup>+0.9</sup> <sub>-1.0</sub>	<sup>4,5</sup> SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons
10987.5 <sup>+6.4</sup> <sub>-2.5</sub> ± 9.1 <sup>+9.1</sup> <sub>-2.3</sub>	<sup>6,7</sup> SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10996 ± 2	<sup>8</sup> AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
11019 ± 5 ± 7	BESSION	85	CLEO $e^+e^- \rightarrow$ hadrons
11020 ± 30	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

<sup>2</sup> From a simultaneous fit to the  $h_b(nP)\pi^+\pi^-$ ,  $n = 1, 2$  cross sections at 22 energy points within  $\sqrt{s} = 10.77\text{--}11.02$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single relative phase, a single relative amplitude, and two overall normalization factors, one for each  $n$ ). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

<sup>3</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

<sup>4</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).

<sup>5</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions.

<sup>6</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.

<sup>7</sup> Superseded by MIZUK 19.

<sup>8</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

**$\Upsilon(11020)$  WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>24 <math>\pm \frac{8}{6}</math> OUR AVERAGE</b>			
23.8 $^{+8.0+0.7}_{-6.8-1.8}$	<sup>1</sup> MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
27 $^{+27+5}_{-11-12}$	<sup>2</sup> MIZUK	16 BELL	$e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
35.1 $\pm 1.2$	<sup>3</sup> DONG	20A	$e^+e^- \rightarrow b\bar{b}$
39.3 $^{+1.7+1.3}_{-1.6-2.4}$	<sup>4,5</sup> SANTEL	16 BELL	$e^+e^- \rightarrow \text{hadrons}$
61 $^{+9+2}_{-19-20}$	<sup>6,7</sup> SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
37 $\pm 3$	<sup>8</sup> AUBERT	09E BABR	$e^+e^- \rightarrow \text{hadrons}$
61 $\pm 13 \pm 22$	BESSION	85 CLEO	$e^+e^- \rightarrow \text{hadrons}$
90 $\pm 20$	LOVELOCK	85 CUSB	$e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

<sup>2</sup> From a simultaneous fit to the  $h_b(nP)\pi^+\pi^-$ ,  $n = 1, 2$  cross sections at 22 energy points within  $\sqrt{s} = 10.77\text{--}11.02$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single relative phase, a single relative amplitude, and two overall normalization factors, one for each  $n$ ). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

<sup>3</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

<sup>4</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).

<sup>5</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions.

<sup>6</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n=1, 2, 3$ , cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.

<sup>7</sup> Superseded by MIZUK 19.

<sup>8</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

## $\Upsilon(11020)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	$(5.4^{+1.9}_{-2.1}) \times 10^{-6}$
$\Gamma_2$ $\Upsilon(1S)\pi^+\pi^-$	
$\Gamma_3$ $\Upsilon(2S)\pi^+\pi^-$	
$\Gamma_4$ $\Upsilon(3S)\pi^+\pi^-$	
$\Gamma_5$ $\chi_{bJ}(1P)\pi^+\pi^-\pi^0$	$(9^{+9}_{-8}) \times 10^{-3}$
$\Gamma_6$ $\chi_{b1}(1P)\pi^+\pi^-\pi^0$	seen
$\Gamma_7$ $\chi_{b2}(1P)\pi^+\pi^-\pi^0$	seen

## $\Upsilon(11020)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					$\Gamma_1$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
<b>0.130±0.030 OUR AVERAGE</b>					
0.095±0.03 ±0.035	BESSON	85	CLEO	$e^+e^- \rightarrow$ hadrons	
0.156±0.040	LOVELOCK	85	CUSB	$e^+e^- \rightarrow$ hadrons	

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_2/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.46±0.08	<sup>1,2</sup> MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$ . <sup>2</sup> Reported as the range 0.38–0.54 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.					

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_3/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.65±0.52	<sup>1,2</sup> MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$ . <sup>2</sup> Reported as the range 0.13–1.16 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.					

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_4/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.33±0.16	<sup>1,2</sup> MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$ . <sup>2</sup> Reported as the range 0.17–0.49 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.					

$\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_5/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.7 \pm 4.3^{+7.6}_{-6.6}$	YIN	18	BELL	$e^+e^- \rightarrow$ hadrons

  

$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_6/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>seen</b>	YIN	18	BELL	$e^+e^- \rightarrow$ hadrons

  

$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_7/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>seen</b>	YIN	18	BELL	$e^+e^- \rightarrow$ hadrons

  

$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)$				$\Gamma_7/\Gamma_6$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.4 \pm 0.2$	YIN	18	BELL	$e^+e^- \rightarrow$ hadrons

### $\Upsilon(11020)$ REFERENCES

DONG	20A	CP C44 083001	X.-K. Dong <i>et al.</i>	
MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)
YIN	18	PR D98 091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSION	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)