

94. Leptoquarks

Revised August 2021 by S. Rolli (DOE) and M. Tanabashi (Nagoya U.; KMI, Nagoya U.).

Leptoquarks are hypothetical particles carrying both baryon number (B) and lepton number (L). The possible quantum numbers of leptoquark states can be restricted by assuming that their direct interactions with the ordinary Standard Model (SM) fermions are dimensionless and invariant under the SM gauge group. Table 94.1 shows the list of all possible quantum numbers with this assumption [1]. The columns of $SU(3)_C$, $SU(2)_W$, and $U(1)_Y$ in Table 94.1 indicate the QCD representation, the weak isospin representation, and the weak hypercharge, respectively. The spin of a leptoquark state is taken to be 1 (vector leptoquark) or 0 (scalar leptoquark).

Table 94.1: Possible leptoquarks and their quantum numbers.

Spin	$3B + L$	$SU(3)_C$	$SU(2)_W$	$U(1)_Y$	Allowed coupling
0	-2	$\bar{3}$	1	1/3	$\bar{q}_L^c \ell_L$ or $\bar{u}_R^c e_R$
0	-2	$\bar{3}$	1	4/3	$\bar{d}_R^c e_R$
0	-2	$\bar{3}$	3	1/3	$\bar{q}_L^c \ell_L$
1	-2	$\bar{3}$	2	5/6	$\bar{q}_L^c \gamma^\mu e_R$ or $\bar{d}_R^c \gamma^\mu \ell_L$
1	-2	$\bar{3}$	2	-1/6	$\bar{u}_R^c \gamma^\mu \ell_L$
0	0	3	2	7/6	$\bar{q}_L e_R$ or $\bar{u}_R \ell_L$
0	0	3	2	1/6	$\bar{d}_R \ell_L$
1	0	3	1	2/3	$\bar{q}_L \gamma^\mu \ell_L$ or $\bar{d}_R \gamma^\mu e_R$
1	0	3	1	5/3	$\bar{u}_R \gamma^\mu e_R$
1	0	3	3	2/3	$\bar{q}_L \gamma^\mu \ell_L$

If we do not require leptoquark states to couple directly with SM fermions, different assignments of quantum numbers become possible [2, 3].

Leptoquark states are expected to exist in various extensions of the SM. The Pati-Salam model [4] is an example predicting the existence of a leptoquark state. Leptoquark states also exist in grand unification theories based on $SU(5)$ [5], $SO(10)$ [6], which includes Pati-Salam color $SU(4)$, and larger gauge groups. Scalar quarks in supersymmetric models with R -parity violation may also have leptoquark-type Yukawa couplings. The bounds on the leptoquark states can therefore be applied to constrain R -parity-violating supersymmetric models. Scalar leptoquarks are expected to exist at the TeV scale in extended technicolor models [7, 8] where leptoquark states appear as the bound states of techni-fermions. Compositeness of quarks and leptons also provides examples of models which may have light leptoquark states [9].

Bounds on leptoquark states are obtained both directly and indirectly. Direct limits are from their production cross sections at colliders, while indirect limits are calculated from bounds on leptoquark-induced four-fermion interactions, which are obtained from low-energy experiments, or from collider experiments below threshold. These four-fermion interactions often cause lepton-flavor non-universalities in heavy quark decays. Anomalies observed recently in the R_K and R_D ratios [10–12] in the semi-leptonic B decays may be explained in models with TeV scale leptoquarks.

If a leptoquark couples to quarks (leptons) belonging to more than a single generation in the mass eigenbasis, it can induce four-fermion interactions causing flavor-changing neutral currents (lepton-family-number violations). The quantum number assignment of Table 1 allows several leptoquark states to couple to both left- and right-handed quarks simultaneously. Such leptoquark states are called non-chiral and may cause four-fermion interactions affecting the $(\pi \rightarrow e\nu)/(\pi \rightarrow \mu\nu)$

ratio [13]. Non-chiral scalar leptoquarks also contribute to the muon anomalous magnetic moment [14, 15]. Since indirect limits provide more stringent constraints on these types of leptoquarks, it is often assumed that a leptoquark state couples only to a single generation of quarks and a single generation of leptons in a chiral interaction, for which indirect limits become much weaker. Additionally, this assumption gives strong constraints on models of leptoquarks.

Refs. [16–18] give extensive lists of the bounds on the leptoquark-induced four-fermion interactions. For the isoscalar scalar and vector leptoquarks S_0 and V_0 , for example, which couple with the first- (second-) generation left-handed quark, and the first-generation left-handed lepton, the bounds $\lambda^2 < 0.07 \times (M_{LQ}/1 \text{ TeV})^2$ for S_0 , and $\lambda^2 < 0.4 \times (M_{LQ}/1 \text{ TeV})^2$ for V_0 ($\lambda^2 < 0.7 \times (M_{LQ}/1 \text{ TeV})^2$ for S_0 , and $\lambda^2 < 0.5 \times (M_{LQ}/1 \text{ TeV})^2$ for V_0) with λ being the leptoquark coupling strength, can be derived from the limits listed in Ref. [19]. The e^+e^- experiments are sensitive to the indirect effects coming from t - and u -channel exchanges of leptoquarks in the $e^+e^- \rightarrow q\bar{q}$ process. The HERA experiments give bounds on the leptoquark-induced four-fermion interaction. For detailed bounds obtained in this way, see the Boson Particle Listings for “Indirect Limits for Leptoquarks” and its references.

Collider experiments provide direct limits on the leptoquark states through limits on the pair- and single-production cross sections. The leading-order cross sections of the parton processes

$$\begin{aligned} q + \bar{q} &\rightarrow LQ + \bar{LQ} \\ g + g &\rightarrow LQ + \bar{LQ} \\ e + q &\rightarrow LQ \end{aligned} \tag{94.1}$$

may be written as [20]

$$\begin{aligned} \hat{\sigma}_{LO}[q\bar{q} \rightarrow LQ + \bar{LQ}] &= \frac{2\alpha_s^2\pi}{27\hat{s}}\beta^3, \\ \hat{\sigma}_{LO}[gg \rightarrow LQ + \bar{LQ}] &= \frac{\alpha_s^2\pi}{96\hat{s}} \times \left[\beta(41 - 31\beta^2) + (18\beta^2 - \beta^4 - 17) \log \frac{1 + \beta}{1 - \beta} \right], \\ \hat{\sigma}_{LO}[eq \rightarrow LQ] &= \frac{\pi\lambda^2}{4}\delta(\hat{s} - M_{LQ}^2) \end{aligned} \tag{94.2}$$

for a scalar leptoquark. Here $\sqrt{\hat{s}}$ is the invariant energy of the parton subprocess, and $\beta \equiv \sqrt{1 - 4M_{LQ}^2/\hat{s}}$. The leptoquark Yukawa coupling is given by λ . Leptoquarks are also produced singly at hadron colliders through $g + q \rightarrow LQ + \ell$ [21], which allows extending to higher masses the collider reach in the leptoquark search [22], depending on the leptoquark Yukawa coupling. See also Ref. [23] for a comprehensive review on the leptoquark phenomenology in precision experiments and particle colliders.

Leptoquark states which couple only to left- or right-handed quarks are called chiral leptoquarks. Leptoquark states which couple only to the first (second, third) generation are referred as the first- (second-, third-) generation leptoquarks.

The LHC, Tevatron and LEP experiments have been searching for pair production of the leptoquark states, which arises from the leptoquark gauge interaction. Due to the typical decay of the leptoquark into charged and neutral leptons and quarks, the searches are carried on in signatures including high p_T charged leptons, high E_T jets and large missing transverse energy. Additionally, searches for pair produced LQs are often organized by the decay mode of the pair of LQs, via

the decay parameter β , which represents the branching fraction into a charge lepton vs a neutrino: $\beta = 1$ for both LQs decaying into a charged lepton, $\beta = 0.5$ for one LQ decaying into a charged lepton and one into a neutrino. The gauge couplings of a scalar leptoquark are determined uniquely according to its quantum numbers in Table 94.1. Since all of the leptoquark states belong to color-triplet representation, the scalar leptoquark pair-production cross section at the Tevatron and LHC can be determined solely as a function of the leptoquark mass without making further assumptions. This is in contrast to the indirect or single-production limits, which give constraints in the leptoquark mass-coupling plane.

Older results from the Tevatron run can be found here: [24–27].

Since the previous version of this review, both ATLAS and CMS have updated their results concerning searches for first, second, and third generation LQs and leptoquark states which couple only with the i -th generation quarks and the j -th generation leptons ($i \neq j$) without causing conflicts with severe indirect constraints. The datasets were almost all collected at center of mass energy of 13 TeV and corresponding to the latest integrated luminosity collected before the shutdown of the LHC occurring in 2019 and 2020.

It is worthy to note that organizing LQs by flavor quantum number first before organizing them by gauge quantum number is becoming more common and advantageous because it relates more closely to some of the experimental searches being performed. The traditional nomenclature for 1st, 2nd, and 3rd generation LQ encourages only looking for the diagonal elements in a flavor matrix of possibilities, which has been the traditional experimental search strategy.

Current results extend previous mass limits for scalar leptoquarks to > 1435 GeV (first generation, CMS, $\beta = 1$, $\sqrt{s} = 13$ TeV) and > 1270 GeV (first generation, CMS, $\beta = 0.5$, $\sqrt{s} = 13$ TeV) [28]; > 1800 GeV (first generation, ATLAS, $\beta = 1$, $\sqrt{s} = 13$ TeV) [29] and > 1290 GeV (first generation, ATLAS, $\beta = 0.5$, $\sqrt{s} = 13$ TeV) [30]; > 1530 GeV (second generation, CMS, $\beta = 1$, $\sqrt{s} = 13$ TeV) and > 1285 GeV (second generation, CMS, $\beta = 0.5$, $\sqrt{s} = 13$ TeV) [31]; and > 1700 GeV (second generation, ATLAS, $\beta = 1$, $\sqrt{s} = 13$ TeV) [29] and > 1230 GeV (second generation, ATLAS, $\beta = 0.5$, $\sqrt{s} = 13$ TeV) [30]. All limits are presented at 95% C.L.

As for third generation leptoquarks, CMS results are the following: 1) assuming that all leptoquarks decay to a top quark and a τ lepton, the existence of pair produced, third-generation leptoquark up to a mass of 900 GeV ($\beta = 1$, 13 TeV) is excluded at 95% confidence level [32]; 2) assuming that all leptoquarks decay to a bottom quark and a τ lepton, the existence of pair produced, third-generation leptoquark up to a mass of 1020 GeV ($\beta = 1$, 13 TeV) is excluded at 95% confidence level [33]; 3) assuming that all leptoquarks decay to a bottom quark and a τ neutrino, the existence of pair produced, third-generation leptoquark up to a mass of 450 GeV ($\beta = 0$, 7 TeV) is excluded at 95% confidence level [34]. In a recent paper [35] signatures of top quark τ lepton ν bottom and top $\tau \nu$ - not previously explored in dedicated searches, were analyzed in the context of searches for scalar leptoquark of charge $-1/3e$ coupling to a top quark plus a τ lepton ($t\tau$) or a bottom quark plus a neutrino ($b\nu$), or a vector particle of charge $+2/3e$, coupling to $t\nu$ or $b\tau$. These choices are motivated by models that can explain a series of anomalies observed in the measurement of B meson decays. The data are found to be in agreement with the standard model prediction. Lower limits at 95% confidence level are set on the LQ mass in the range 0.98–1.73 TeV, depending on the LQ spin and its coupling λ to a lepton and a quark, and assuming equal couplings for the two LQ decay modes considered. These are the most stringent constraints to date on the existence of leptoquarks in this scenario.

In [36], the ATLAS collaboration has limits on pair production of third generation scalar leptoquarks where all possible decays of the leptoquark into a quark (t , b) and a lepton (τ , ν) of the third generation are considered. The limits are presented as a function of the leptoquark mass and the branching ratio into charged leptons for leptoquark of up-type ($LQ_3^{up} \rightarrow \tau\nu/b\tau$) and down-type

($LQ_3^d \rightarrow b\nu/\tau\tau$); many results are re-interpretation of previously published ATLAS searches. The collaboration finds that masses below 800 GeV are excluded for both LQ_3^u and LQ_3^d independently of the branching ratio, with masses below about 1 TeV being excluded for the limiting cases of branching ratios equal to zero or unity. In a more recent paper [37] ATLAS present the result of searches for pair production of third-generation scalar leptoquarks decaying into a top quark and a τ -lepton, using 139 fb^{-1} of data collected at 13 TeV. Scalar leptoquarks decaying exclusively into $\tau\tau$ are excluded up to masses of 1.43 TeV while, for a branching fraction of 50% into $t\tau$, the lower mass limit is 1.22 TeV.

It is also possible to consider leptoquark states which couple only with the i -th generation quarks and the j -th generation leptons ($i \neq j$) without causing conflicts with severe indirect constraints. Such couplings have received renewed attention because they may provide an explanation to anomalies in rare $B - meson$ decays and the anomalous magnetic moment of the muon. See Ref. [38,39] and [40] and references therein for collider search strategies and limits on the pair production cross sections of this class of leptoquark states. In this framework, a novel CMS result [41] presents a non-traditional search for pair production of LQs coupled to a top quark and a muon. As no deviation from the Standard Model prediction was observed, scalar LQs decaying exclusively into $top - \mu$ are excluded up to masses of 1420 GeV. In a recent ATLAS paper [42] a search for pair production of scalar leptoquarks, each decaying into either an electron or a muon and a top quark, is presented. This is the first leptoquark search using ATLAS data to investigate top-philic cross-generational couplings that could provide explanations for recently observed anomalies in B meson decays. In the absence of any significant deviation from the background expectation, lower limits on the leptoquark masses are set at 1480 GeV and 1470 GeV for the electron and muon channel, respectively.

The magnetic-dipole-type and the electric-quadrupole-type interactions of a vector leptoquark are not determined even if we fix its gauge quantum numbers as listed in the Table 94.1 [43]. The production of vector leptoquarks depends in general on additional assumptions, where the leptoquark couplings and their pair production cross sections are enhanced relative to the scalar leptoquark contributions. The most stringent limits on vector LQ production are now from CMS [44] where previous searches for squarks and gluinos have been reinterpreted to constrain models of leptoquark production. LQ masses below 1530 GeV are excluded assuming the Yang-Mills case with coupling $\kappa = 1$, or 1115 GeV in the minimal coupling case where $\kappa = 0$, placing the most stringent constraint to date from pair production of vector LQs.

The leptoquark pair-production cross sections in e^+e^- collisions depend on the leptoquark $SU(2) \times U(1)$ quantum numbers and Yukawa coupling with electron [45].

Searches for first generation leptoquark singly produced were performed by the HERA experiments. Since the leptoquark single-production cross section depends on its Yukawa coupling, the leptoquark mass limits from HERA are usually displayed in the mass-coupling plane. For leptoquark Yukawa coupling $\lambda = 0.1$, early ZEUS Collaboration bounds on the first-generation leptoquarks range from 248 to 290 GeV, depending on the leptoquark species [46]. The ZEUS Collaboration has recently released a new paper [47] where data corresponding to a luminosity of around 1 fb^{-1} have been used in the framework of $eeqq$ contact interactions (CI) to set limits on possible high-energy contributions beyond the Standard Model to electron-quark scattering. The analysis of the ep data has been based on simultaneous fits of parton distribution functions including contributions of Contact Interaction (CI) couplings to ep scattering. Several general CI models and scenarios with heavy leptoquarks were considered. As unambiguous deviations from the SM cannot be established, limits for CI compositeness scales and LQ mass scales were set that are in the TeV range. The H1 Collaboration has a comprehensive summary of searches for first generation leptoquarks using the full data sample collected in ep collisions at HERA (446 pb^{-1}). No evidence of production of

leptoquarks was observed in final states with a large transverse momentum electron or large missing transverse momentum. For a coupling strength $\lambda = 0.3$, first generation leptoquarks with masses up to 800 GeV are excluded at 95% C.L. [48].

At the LHC, the CMS collaboration performed searches for single production of first and second generation leptoquarks [49], which is complementary to the HERA searches in the high λ region (for coupling strength $\lambda = 1.0$, first generation leptoquarks are excluded for masses up to 1.73 TeV and second generation leptoquark are excluded up to masses of 530 GeV). CMS also recently searched for third generation LQ decaying into τ and *bottom* in [50]. Assuming unit Yukawa coupling (λ), a third generation scalar leptoquark is excluded for masses below 740 GeV. Limits are also set on λ of the hypothesized leptoquark as a function of its mass. Above $\lambda = 1.4$, the results provide the best upper limit on the mass of a third-generation scalar leptoquark decaying to a τ lepton and a bottom quark.

Searches for LQ will continue with more LHC data, particularly in light of the renewed interest in this type of particle to explain violation of lepton flavor universality and other anomalies, which point to explanations laying outside the Standard Model.

References

- [1] W. Buchmuller, R. Ruckl and D. Wyler, *Phys. Lett. B* **191**, 442 (1987), [Erratum: *Phys.Lett.B* 448, 320–320 (1999)].
- [2] K. Babu, C. F. Kolda and J. March-Russell, *Phys. Lett. B* **408**, 261 (1997), [hep-ph/9705414].
- [3] J. L. Hewett and T. G. Rizzo, *Phys. Rev. D* **58**, 055005 (1998), [hep-ph/9708419].
- [4] J. C. Pati and A. Salam, *Phys. Rev. D* **10**, 275 (1974), [Erratum: *Phys.Rev.D* 11, 703–703 (1975)].
- [5] H. Georgi and S. Glashow, *Phys. Rev. Lett.* **32**, 438 (1974).
- [6] H. Georgi, *AIP Conf. Proc.* **23**, 575 (1975); H. Fritzsch and P. Minkowski, *Annals Phys.* **93**, 193 (1975).
- [7] For a review, see, E. Farhi and L. Susskind, *Phys. Rept.* **74**, 277 (1981).
- [8] K. D. Lane and M. Ramana, *Phys. Rev. D* **44**, 2678 (1991).
- [9] See, for example, B. Schrempp and F. Schrempp, *Phys. Lett.* **153B**, 101 (1985).
- [10] R. Aaij *et al.* (LHCb), *Phys. Rev. Lett.* **120**, 17, 171802 (2018), [arXiv:1708.08856]; R. Aaij *et al.* (LHCb), *Phys. Rev. D* **97**, 7, 072013 (2018), [arXiv:1711.02505].
- [11] R. Aaij *et al.* (LHCb), *JHEP* **08**, 055 (2017), [arXiv:1705.05802]; R. Aaij *et al.* (LHCb), *Phys. Rev. Lett.* **122**, 19, 191801 (2019), [arXiv:1903.09252]; R. Aaij *et al.* (LHCb), *Phys. Rev. Lett.* **125**, 1, 011802 (2020), [arXiv:2003.04831]; R. Aaij *et al.* (LHCb) (2021), [arXiv:2103.11769].
- [12] Y. S. Amhis *et al.* (HFLAV), *Eur. Phys. J. C* **81**, 3, 226 (2021), updated results and plots available at <https://hflav.web.cern.ch/>, [arXiv:1909.12524].
- [13] O. U. Shanker, *Nucl. Phys. B* **204**, 375 (1982).
- [14] U. Mahanta, *Eur. Phys. J. C* **21**, 171 (2001), [hep-ph/0102176].
- [15] K.-M. Cheung, *Phys. Rev. D* **64**, 033001 (2001), [hep-ph/0102238].
- [16] S. Davidson, D. C. Bailey and B. A. Campbell, *Z. Phys. C* **61**, 613 (1994), [hep-ph/9309310].
- [17] M. Leurer, *Phys. Rev. D* **49**, 333 (1994), [hep-ph/9309266]; M. Leurer, *Phys. Rev. D* **50**, 536 (1994), [hep-ph/9312341].
- [18] M. Carpentier and S. Davidson, *Eur. Phys. J. C* **70**, 1071 (2010), [arXiv:1008.0280].
- [19] S. Davidson and A. Saporta, *Phys. Rev. D* **99**, 1, 015032 (2019), [arXiv:1807.10288].

- [20] T. Plehn *et al.*, *Z. Phys. C* **74**, 611 (1997), [[hep-ph/9703433](#)]; M. Kramer *et al.*, *Phys. Rev. Lett.* **79**, 341 (1997), [[hep-ph/9704322](#)].
- [21] J. Hewett and S. Pakvasa, *Phys. Rev. D* **37**, 3165 (1988); O. J. Eboli and A. V. Olinto, *Phys. Rev. D* **38**, 3461 (1988); A. Dobado, M. J. Herrero and C. Munoz, *Phys. Lett. B* **207**, 97 (1988); V. D. Barger *et al.*, *Phys. Lett. B* **220**, 464 (1989); M. De Montigny and L. Marleau, *Phys. Rev. D* **40**, 2869 (1989), [Erratum: *Phys.Rev.D* 56, 3156 (1997)].
- [22] A. Belyaev *et al.*, *JHEP* **09**, 005 (2005), [[hep-ph/0502067](#)].
- [23] I. Doršner *et al.*, *Phys. Rept.* **641**, 1 (2016), [[arXiv:1603.04993](#)].
- [24] V. Abazov *et al.* (D0), *Phys. Lett. B* **681**, 224 (2009), [[arXiv:0907.1048](#)].
- [25] A. Abulencia *et al.* (CDF), *Phys. Rev. D* **73**, 051102 (2006), [[hep-ex/0512055](#)].
- [26] V. Abazov *et al.* (D0), *Phys. Lett. B* **671**, 224 (2009), [[arXiv:0808.4023](#)].
- [27] V. M. Abazov *et al.* (D0), *Phys. Lett. B* **693**, 95 (2010), [[arXiv:1005.2222](#)].
- [28] A. M. Sirunyan *et al.* (CMS), *Phys. Rev. D* **99**, 5, 052002 (2019), [[arXiv:1811.01197](#)].
- [29] G. Aad *et al.* (ATLAS), *JHEP* **10**, 112 (2020), [[arXiv:2006.05872](#)].
- [30] M. Aaboud *et al.* (ATLAS), *Eur. Phys. J. C* **79**, 9, 733 (2019), [[arXiv:1902.00377](#)].
- [31] A. M. Sirunyan *et al.* (CMS), *Phys. Rev. D* **99**, 3, 032014 (2019), [[arXiv:1808.05082](#)].
- [32] A. M. Sirunyan *et al.* (CMS), *Eur. Phys. J. C* **78**, 707 (2018), [[arXiv:1803.02864](#)].
- [33] A. M. Sirunyan *et al.* (CMS), *JHEP* **03**, 170 (2019), [[arXiv:1811.00806](#)].
- [34] S. Chatrchyan *et al.* (CMS), *JHEP* **12**, 055 (2012), [[arXiv:1210.5627](#)].
- [35] A. M. Sirunyan *et al.* (CMS), *Phys. Lett. B* **819**, 136446 (2021), [[arXiv:2012.04178](#)].
- [36] M. Aaboud *et al.* (ATLAS), *JHEP* **06**, 144 (2019), [[arXiv:1902.08103](#)].
- [37] G. Aad *et al.* (ATLAS), *JHEP* **06**, 179 (2021), [[arXiv:2101.11582](#)].
- [38] B. Diaz, M. Schmaltz and Y.-M. Zhong, *JHEP* **10**, 097 (2017), [[arXiv:1706.05033](#)].
- [39] M. Schmaltz and Y.-M. Zhong, *JHEP* **01**, 132 (2019), [[arXiv:1810.10017](#)].
- [40] D. Müller, *EPJ Web Conf.* **179**, 01015 (2018), [[arXiv:1801.03380](#)].
- [41] A. M. Sirunyan *et al.* (CMS), *Phys. Rev. Lett.* **121**, 24, 241802 (2018), [[arXiv:1809.05558](#)].
- [42] G. Aad *et al.* (ATLAS), *Eur. Phys. J. C* **81**, 4, 313 (2021), [[arXiv:2010.02098](#)].
- [43] J. Blumlein, E. Boos and A. Kryukov, *Z. Phys. C* **76**, 137 (1997), [[hep-ph/9610408](#)].
- [44] A. M. Sirunyan *et al.* (CMS), *Phys. Rev. D* **98**, 3, 032005 (2018), [[arXiv:1805.10228](#)].
- [45] J. Blumlein and R. Ruckl, *Phys. Lett. B* **304**, 337 (1993).
- [46] S. Chekanov *et al.* (ZEUS), *Phys. Rev. D* **68**, 052004 (2003), [[hep-ex/0304008](#)].
- [47] H. Abramowicz *et al.* (ZEUS), *Phys. Rev. D* **99**, 9, 092006 (2019), [[arXiv:1902.03048](#)].
- [48] F. Aaron *et al.* (H1), *Phys. Lett. B* **704**, 388 (2011), [[arXiv:1107.3716](#)].
- [49] V. Khachatryan *et al.* (CMS), *Phys. Rev. D* **93**, 3, 032005 (2016), [Erratum: *Phys.Rev.D* 95, 039906 (2017)], [[arXiv:1509.03750](#)].
- [50] A. Sirunyan *et al.* (CMS), *JHEP* **07**, 115 (2018), [[arXiv:1806.03472](#)].