

## 54. Mass and Width of the $W$ Boson

Revised August 2021 by M. Grünewald (University Coll. Dublin) and A. Gurtu (CERN; TIFR Mumbai).

Precision determination of the  $W$ -mass is of great importance in testing the internal consistency of the Standard Model. From the time of its discovery in 1983, the  $W$ -boson has been studied and its mass determined in  $p\bar{p}$  and  $e^+e^-$  interactions; it is currently studied in  $pp$  interactions at the LHC. The  $W$  mass and width definition used here corresponds to a Breit-Wigner with mass-dependent width.

Production of on-shell  $W$  bosons at hadron colliders is tagged by the high  $p_T$  charged lepton from its decay. Owing to the unknown parton-parton effective energy and missing energy in the longitudinal direction, the collider experiments reconstruct the transverse mass of the  $W$ , and derive the  $W$  mass from comparing the transverse mass distribution with Monte Carlo predictions as a function of  $M_W$ . These analyses use the electron and muon decay modes of the  $W$  boson.


In the  $e^+e^-$  collider (LEP) a precise knowledge of the beam energy enables one to determine the  $e^+e^- \rightarrow W^+W^-$  cross section as a function of center of mass energy, as well as to reconstruct the  $W$  mass precisely from its decay products, even if one of them decays leptonically. Close to the  $W^+W^-$  threshold (161 GeV), the dependence of the  $W$ -pair production cross section on  $M_W$  is large, and this was used to determine  $M_W$ . At higher energies (172 to 209 GeV) this dependence is much weaker and  $W$ -bosons were directly reconstructed and the mass determined as the invariant mass of its decay products, improving the resolution with a kinematic fit.

In order to compute the LEP average  $W$  mass, each experiment provided its measured  $W$  mass for the  $q\bar{q}q\bar{q}$  and  $q\bar{q}\ell\nu_\ell$ ,  $\ell = e, \mu, \tau$  channels at each center-of-mass energy, along with a detailed break-up of errors: statistical, uncorrelated, partially correlated and fully correlated systematics [1]. These have been combined to obtain a LEP  $W$  mass of  $M_W = 80.376 \pm 0.033$  GeV. Errors due to uncertainties in LEP energy (9 MeV), and possible effect of color reconnection (CR) and Bose-Einstein correlations (BEC) between quarks from different  $W$ 's (8 MeV) are included. The mass difference between  $q\bar{q}q\bar{q}$  and  $q\bar{q}\ell\nu_\ell$  final states (due to possible CR and BEC effects) is  $-12 \pm 45$  MeV. In a similar manner, the width results obtained at LEP have been combined, resulting in  $\Gamma_W = 2.195 \pm 0.083$  GeV [1].

The two Tevatron experiments have also identified common systematic errors. Between the two experiments, uncertainties due to the parton distribution functions, radiative corrections, and choice of mass (width) in the width (mass) measurements are treated as correlated. An average  $W$  width of  $\Gamma_W = 2.046 \pm 0.049$  GeV [2] is obtained. Errors of 20 MeV and 7 MeV accounting for PDF and radiative correction uncertainties in this width combination dominate the correlated uncertainties. The CDF and D0 values of  $80.387 \pm 0.019$  GeV [3] and  $80.375 \pm 0.023$  GeV [4], respectively, are the two most precise determinations of the  $W$  boson mass. Combining all Tevatron results from Run-I and Run-II using an improved treatment of correlations, a new average of  $80.387 \pm 0.016$  GeV is obtained [5], with common uncertainties of 10 MeV (PDF) and 4 MeV (radiative corrections).


Good agreement between the LEP and Tevatron results is observed. Combining these results, assuming no common systematic uncertainties between the LEP and the Tevatron measurements, yields an average  $W$  mass of  $M_W = 80.385 \pm 0.015$  GeV and a  $W$  width of  $\Gamma_W = 2.085 \pm 0.042$  GeV.

Based on  $pp$  collisions at  $\sqrt{s} = 7$  TeV, the ATLAS collaboration has determined the  $W$  boson mass as  $M_W = 80.370 \pm 0.019$  GeV [6], which is compatible with the above world average and of similar precision to the best measurements of CDF and D0. Assuming a Tevatron/LHC common PDF uncertainty of 7 MeV [7], this results in a new world average of  $M_W = 80.379 \pm 0.012$  GeV.



**Figure 54.1:** Measurements of the  $W$  boson mass by the LEP, Tevatron and LHC experiments.

The LEP, Tevatron and LHC results on mass and width, which are based on all results available, are compared in Fig. 54.1 and Fig. 54.2. The Standard Model prediction from the electroweak fit, including Z-pole data and  $m_{\text{top}}$  and  $M_H$  measurements, gives a W-boson mass of  $M_W = 80.354 \pm 0.007$  GeV and a W-boson width of  $\Gamma_W = 2.091 \pm 0.001$  GeV [8].



**Figure 54.2:** Measurements of the  $W$ -boson width by the LEP and Tevatron experiments.

### References

- [1] S. Schael *et al.* (ALEPH, DELPHI, L3, OPAL, LEP Electroweak), *Phys. Rept.* **532**, 119 (2013), [arXiv:1302.3415].
- [2] The Tevatron Electroweak Working Group, for the CDF and D0 Collaborations, March 2010, [arXiv:1003.2826].
- [3] T. Aaltonen *et al.* (CDF), *Phys. Rev. Lett.* **108**, 151803 (2012), [arXiv:1203.0275].
- [4] V. M. Abazov *et al.* (D0), *Phys. Rev. Lett.* **108**, 151804 (2012), [arXiv:1203.0293].
- [5] T. A. Aaltonen *et al.* (CDF, D0), *Phys. Rev. D* **88**, 5, 052018 (2013), [arXiv:1307.7627].
- [6] M. Aaboud *et al.* (ATLAS), *Eur. Phys. J. C* **78**, 2, 110 (2018), [Erratum: *Eur. Phys. J. C* **78**, 898 (2018)], [arXiv:1701.07240].
- [7] J. Erler, in “52nd Rencontres de Moriond on EW Interactions and Unified Theories,” 85–94 (2017), [arXiv:1704.08330].
- [8] J. Haller *et al.*, *Eur. Phys. J. C* **78**, 8, 675 (2018), [arXiv:1803.01853].