

CHARMED BARYONS ($C = +1$)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc, \\ \Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

 Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Mass } m = 2286.46 \pm 0.14 \text{ MeV}$$

$$\text{Mean life } \tau = (202.4 \pm 3.1) \times 10^{-15} \text{ s} \quad (S = 1.7)$$

$$c\tau = 60.7 \text{ } \mu\text{m}$$

Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.84 \pm 0.09$$

$$\Sigma^+\pi^0 \quad \alpha = -0.55 \pm 0.11$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^0\pi^+ = -0.73 \pm 0.18$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.86 \pm 0.04$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow pK_S^0 = 0.2 \pm 0.5$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^- = -0.07 \pm 0.31$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+\nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}e^-\bar{\nu}_e = 0.00 \pm 0.04$$

$$A_{CP}(\Lambda X) \text{ in } \Lambda_c \rightarrow \Lambda X, \bar{\Lambda}_c \rightarrow \bar{\Lambda} X = (2 \pm 7)\%$$

$$\Delta A_{CP} = A_{CP}(\Lambda_c^+ \rightarrow pK^+K^-) - A_{CP}(\Lambda_c^+ \rightarrow p\pi^+\pi^-) = (0.3 \pm 1.1)\%$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Lambda_c^+ \rightarrow p\bar{K}^*(892)^0$ seen in $\Lambda_c^+ \rightarrow pK^-\pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0\pi^0$ decays.

Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Hadronic modes with a p or n : $S = -1$ final states			
$p K_S^0$	(1.59 $\pm$ 0.08) %	S=1.1	873
$p K^- \pi^+$	(6.28 $\pm$ 0.32) %	S=1.4	823
$p \bar{K}^*(892)^0$	[a] (1.96 $\pm$ 0.27) %		685
$\Delta(1232)^{++} K^-$	(1.08 $\pm$ 0.25) %		710
$\Lambda(1520) \pi^+$	[a] (2.2 $\pm$ 0.5) %		628
$p K^- \pi^+$ nonresonant	(3.5 $\pm$ 0.4) %		823
$p K_S^0 \pi^0$	(1.97 $\pm$ 0.13) %	S=1.1	823
$n K_S^0 \pi^+$	(1.82 $\pm$ 0.25) %		821

$p\bar{K}^0\eta$	(1.6 $\pm$ 0.4) %		568
$pK_S^0\pi^+\pi^-$	(1.60 $\pm$ 0.12) %	S=1.1	754
$pK^-\pi^+\pi^0$	(4.46 $\pm$ 0.30) %	S=1.5	759
$pK^*(892)^-\pi^+$	[a] (1.4 $\pm$ 0.5) %		580
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(4.6 $\pm$ 0.8) %		759
$\Delta(1232)\bar{K}^*(892)$	seen		419
$pK^-2\pi^+\pi^-$	(1.4 $\pm$ 0.9) $\times 10^{-3}$		671
$pK^-\pi^+2\pi^0$	(1.0 $\pm$ 0.5) %		678

Hadronic modes with a p : $S = 0$ final states

$p\pi^0$	< 2.7 $\times 10^{-4}$	CL=90%	945
$p\eta$	(1.24 $\pm$ 0.30) $\times 10^{-3}$		856
$p\omega(782)^0$	(9 $\pm$ 4) $\times 10^{-4}$		751
$p\pi^+\pi^-$	(4.61 $\pm$ 0.28) $\times 10^{-3}$		927
$pf_0(980)$	[a] (3.5 $\pm$ 2.3) $\times 10^{-3}$		614
$p2\pi^+2\pi^-$	(2.3 $\pm$ 1.4) $\times 10^{-3}$		852
pK^+K^-	(1.06 $\pm$ 0.06) $\times 10^{-3}$		616
$p\phi$	[a] (1.06 $\pm$ 0.14) $\times 10^{-3}$		590
$pK^+K^- \text{ non-}\phi$	(5.3 $\pm$ 1.2) $\times 10^{-4}$		616
$p\phi\pi^0$	(10 $\pm$ 4) $\times 10^{-5}$		460
$pK^+K^-\pi^0 \text{ nonresonant}$	< 6.3 $\times 10^{-5}$	CL=90%	494

Hadronic modes with a hyperon: $S = -1$ final states

$\Lambda\pi^+$	(1.30 $\pm$ 0.07) %	S=1.1	864
$\Lambda\pi^+\pi^0$	(7.1 $\pm$ 0.4) %	S=1.1	844
$\Lambda\rho^+$	< 6 %	CL=95%	636
$\Lambda\pi^-2\pi^+$	(3.64 $\pm$ 0.29) %	S=1.4	807
$\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+$	(1.0 $\pm$ 0.5) %		688
$\Sigma(1385)^-2\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-$	(7.6 $\pm$ 1.4) $\times 10^{-3}$		688
$\Lambda\pi^+\rho^0$	(1.5 $\pm$ 0.6) %		524
$\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$	(5 $\pm$ 4) $\times 10^{-3}$		363
$\Lambda\pi^-2\pi^+ \text{ nonresonant}$	< 1.1 %	CL=90%	807
$\Lambda\pi^-\pi^02\pi^+ \text{ total}$	(2.3 $\pm$ 0.8) %		757
$\Lambda\pi^+\eta$	[a] (1.84 $\pm$ 0.26) %		691
$\Sigma(1385)^+\eta$	[a] (9.1 $\pm$ 2.0) $\times 10^{-3}$		570
$\Lambda\pi^+\omega$	[a] (1.5 $\pm$ 0.5) %		517
$\Lambda\pi^-\pi^02\pi^+, \text{ no } \eta \text{ or } \omega$	< 8 $\times 10^{-3}$	CL=90%	757
$\Lambda K^+\bar{K}^0$	(5.7 $\pm$ 1.1) $\times 10^{-3}$	S=1.9	443
$\Xi(1690)^0K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$	(1.6 $\pm$ 0.5) $\times 10^{-3}$		286
$\Sigma^0\pi^+$	(1.29 $\pm$ 0.07) %	S=1.1	825
$\Sigma^+\pi^0$	(1.25 $\pm$ 0.10) %		827
$\Sigma^+\eta$	(4.4 $\pm$ 2.0) $\times 10^{-3}$		713
$\Sigma^+\eta'$	(1.5 $\pm$ 0.6) %		391

$\Sigma^+ \pi^+ \pi^-$	(4.50 ± 0.25) %	S=1.3	804
$\Sigma^+ \rho^0$	< 1.7 %	CL=95%	575
$\Sigma^- 2\pi^+$	(1.87 ± 0.18) %		799
$\Sigma^0 \pi^+ \pi^0$	(3.5 ± 0.4) %		803
$\Sigma^+ \pi^0 \pi^0$	(1.55 ± 0.15) %		806
$\Sigma^0 \pi^- 2\pi^+$	(1.11 ± 0.30) %		763
$\Sigma^+ \pi^+ \pi^- \pi^0$	—		767
$\Sigma^+ \omega$	[a] (1.70 ± 0.21) %		569
$\Sigma^- \pi^0 2\pi^+$	(2.1 ± 0.4) %		762
$\Sigma^+ K^+ K^-$	(3.5 ± 0.4) × 10 ⁻³	S=1.1	349
$\Sigma^+ \phi$	[a] (3.9 ± 0.6) × 10 ⁻³	S=1.1	295
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$	(1.02 ± 0.25) × 10 ⁻³		286
$\Sigma^+ K^-$			
$\Sigma^+ K^+ K^-$ nonresonant	< 8 × 10 ⁻⁴	CL=90%	349
$\Xi^0 K^+$	(5.5 ± 0.7) × 10 ⁻³		653
$\Xi^- K^+ \pi^+$	(6.2 ± 0.6) × 10 ⁻³	S=1.1	565
$\Xi(1530)^0 K^+$	(4.3 ± 0.9) × 10 ⁻³	S=1.1	473

Hadronic modes with a hyperon: S = 0 final states

ΛK^+	(6.1 ± 1.2) × 10 ⁻⁴		781
$\Lambda K^+ \pi^+ \pi^-$	< 5 × 10 ⁻⁴	CL=90%	637
$\Sigma^0 K^+$	(5.2 ± 0.8) × 10 ⁻⁴		735
$\Sigma^0 K^+ \pi^+ \pi^-$	< 2.6 × 10 ⁻⁴	CL=90%	574
$\Sigma^+ K^+ \pi^-$	(2.1 ± 0.6) × 10 ⁻³		670
$\Sigma^+ K^*(892)^0$	[a] (3.5 ± 1.0) × 10 ⁻³		470
$\Sigma^- K^+ \pi^+$	< 1.2 × 10 ⁻³	CL=90%	664

Doubly Cabibbo-suppressed modes

$p K^+ \pi^-$	(1.11 ± 0.18) × 10 ⁻⁴		823
---------------	------------------------------------	--	-----

Semileptonic modes

$\Lambda e^+ \nu_e$	(3.6 ± 0.4) %		871
$\Lambda \mu^+ \nu_\mu$	(3.5 ± 0.5) %		867

Inclusive modes

e^+ anything	(3.95 ± 0.35) %		—
p anything	(50 ± 16) %		—
n anything	(50 ± 16) %		—
Λ anything	(38.2 $^{+2.9}_{-2.4}$) %		—
K_S^0 anything	(9.9 ± 0.7) %		—
3prongs	(24 ± 8) %		—

**$\Delta C = 1$ weak neutral current ($C1$) modes, or
Lepton Family number (LF), or Lepton number (L), or
Baryon number (B) violating modes**

$p e^+ e^-$	$C1$	< 5.5	$\times 10^{-6}$	CL=90%	951
$p \mu^+ \mu^-$ non-resonant	$C1$	< 7.7	$\times 10^{-8}$	CL=90%	937
$p e^+ \mu^-$	LF	< 9.9	$\times 10^{-6}$	CL=90%	947
$p e^- \mu^+$	LF	< 1.9	$\times 10^{-5}$	CL=90%	947
$\bar{p} 2 e^+$	L, B	< 2.7	$\times 10^{-6}$	CL=90%	951
$\bar{p} 2 \mu^+$	L, B	< 9.4	$\times 10^{-6}$	CL=90%	937
$\bar{p} e^+ \mu^+$	L, B	< 1.6	$\times 10^{-5}$	CL=90%	947
$\Sigma^- \mu^+ \mu^+$	L	< 7.0	$\times 10^{-4}$	CL=90%	812

$\Lambda_c(2595)^+$

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that $\Sigma_c(2455)\pi$ decays, with little available phase space, are dominant. This assumes that $J^P = 1/2^+$ for the $\Sigma_c(2455)$.

Mass $m = 2592.25 \pm 0.28$ MeV

$m - m_{\Lambda_c^+} = 305.79 \pm 0.24$ MeV

Full width $\Gamma = 2.6 \pm 0.6$ MeV

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma_c(2455)\pi$ — the latter just barely — are the only strong decays allowed to an excited Λ_c^+ having this mass; and the submode seems to dominate.

$\Lambda_c(2595)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[b] —	117
$\Sigma_c(2455)^{++} \pi^-$	24 ± 7 %	†
$\Sigma_c(2455)^0 \pi^+$	24 ± 7 %	†
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	18 ± 10 %	117
$\Lambda_c^+ \pi^0$	[c] not seen	258
$\Lambda_c^+ \gamma$	not seen	288

$\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

Mass $m = 2628.11 \pm 0.19$ MeV ($S = 1.1$)

$m - m_{\Lambda_c^+} = 341.65 \pm 0.13$ MeV ($S = 1.1$)

Full width $\Gamma < 0.97$ MeV, CL = 90%

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma(2455) \pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

$\Lambda_c(2625)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	$\approx 67\%$		184
$\Sigma_c(2455)^{++} \pi^-$	<5	90%	102
$\Sigma_c(2455)^0 \pi^+$	<5	90%	102
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	[c] not seen		293
$\Lambda_c^+ \gamma$	not seen		319

$\Lambda_c(2860)^+$

$$I(J^P) = 0(\frac{3}{2}^+)$$

$$\text{Mass } m = 2856.1^{+2.3}_{-6.0} \text{ MeV}$$

$$\text{Full width } \Gamma = 68^{+12}_{-22} \text{ MeV}$$

$\Lambda_c(2860)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 p$	seen	259

$\Lambda_c(2880)^+$

$$I(J^P) = 0(\frac{5}{2}^+)$$

$$\text{Mass } m = 2881.63 \pm 0.24 \text{ MeV}$$

$$m - m_{\Lambda_c^+} = 595.17 \pm 0.28 \text{ MeV}$$

$$\text{Full width } \Gamma = 5.6^{+0.8}_{-0.6} \text{ MeV}$$

$\Lambda_c(2880)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	seen	471
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	376
$\Sigma_c(2520)^{0,++} \pi^\pm$	seen	317
$p D^0$	seen	316

$\Lambda_c(2940)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

$J^P = 3/2^-$ is favored, but is not certain

$$\text{Mass } m = 2939.6^{+1.3}_{-1.5} \text{ MeV}$$

$$\text{Full width } \Gamma = 20^{+6}_{-5} \text{ MeV}$$

$\Lambda_c(2940)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$p D^0$	seen	420
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	—

 $\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+)$$

$$\begin{aligned} \Sigma_c(2455)^{++} \text{ mass } m &= 2453.97 \pm 0.14 \text{ MeV} \\ \Sigma_c(2455)^+ \text{ mass } m &= 2452.9 \pm 0.4 \text{ MeV} \\ \Sigma_c(2455)^0 \text{ mass } m &= 2453.75 \pm 0.14 \text{ MeV} \\ m_{\Sigma_c^{++}} - m_{\Lambda_c^+} &= 167.510 \pm 0.017 \text{ MeV} \\ m_{\Sigma_c^+} - m_{\Lambda_c^+} &= 166.4 \pm 0.4 \text{ MeV} \\ m_{\Sigma_c^0} - m_{\Lambda_c^+} &= 167.290 \pm 0.017 \text{ MeV} \\ m_{\Sigma_c^{++}} - m_{\Sigma_c^0} &= 0.220 \pm 0.013 \text{ MeV} \\ m_{\Sigma_c^+} - m_{\Sigma_c^0} &= -0.9 \pm 0.4 \text{ MeV} \\ \Sigma_c(2455)^{++} \text{ full width } \Gamma &= 1.89^{+0.09}_{-0.18} \text{ MeV} \quad (S = 1.1) \\ \Sigma_c(2455)^+ \text{ full width } \Gamma &< 4.6 \text{ MeV, CL} = 90\% \\ \Sigma_c(2455)^0 \text{ full width } \Gamma &= 1.83^{+0.11}_{-0.19} \text{ MeV} \quad (S = 1.2) \end{aligned}$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2455)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	94

 $\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\begin{aligned} \Sigma_c(2520)^{++} \text{ mass } m &= 2518.41^{+0.21}_{-0.19} \text{ MeV} \quad (S = 1.1) \\ \Sigma_c(2520)^+ \text{ mass } m &= 2517.5 \pm 2.3 \text{ MeV} \\ \Sigma_c(2520)^0 \text{ mass } m &= 2518.48 \pm 0.20 \text{ MeV} \quad (S = 1.1) \\ m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} &= 231.95^{+0.17}_{-0.12} \text{ MeV} \quad (S = 1.3) \\ m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} &= 231.0 \pm 2.3 \text{ MeV} \\ m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} &= 232.02^{+0.15}_{-0.14} \text{ MeV} \quad (S = 1.3) \\ m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} &= 0.01 \pm 0.15 \text{ MeV} \\ \Sigma_c(2520)^{++} \text{ full width } \Gamma &= 14.78^{+0.30}_{-0.40} \text{ MeV} \\ \Sigma_c(2520)^+ \text{ full width } \Gamma &< 17 \text{ MeV, CL} = 90\% \\ \Sigma_c(2520)^0 \text{ full width } \Gamma &= 15.3^{+0.4}_{-0.5} \text{ MeV} \end{aligned}$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2520)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	179

$\Sigma_c(2800)$

$$I(J^P) = 1(?^?)$$

$$\begin{aligned} \Sigma_c(2800)^{++} \text{ mass } m &= 2801_{-6}^{+4} \text{ MeV} \\ \Sigma_c(2800)^+ \text{ mass } m &= 2792_{-5}^{+14} \text{ MeV} \\ \Sigma_c(2800)^0 \text{ mass } m &= 2806_{-7}^{+5} \text{ MeV} \quad (S = 1.3) \\ m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} &= 514_{-6}^{+4} \text{ MeV} \\ m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} &= 505_{-5}^{+14} \text{ MeV} \\ m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} &= 519_{-7}^{+5} \text{ MeV} \quad (S = 1.3) \\ \Sigma_c(2800)^{++} \text{ full width } \Gamma &= 75_{-17}^{+22} \text{ MeV} \\ \Sigma_c(2800)^+ \text{ full width } \Gamma &= 62_{-40}^{+60} \text{ MeV} \\ \Sigma_c(2800)^0 \text{ full width } \Gamma &= 72_{-15}^{+22} \text{ MeV} \end{aligned}$$

$\Sigma_c(2800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	seen	443

Ξ_c^+

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\begin{aligned} \text{Mass } m &= 2467.71 \pm 0.23 \text{ MeV} \quad (S = 1.3) \\ \text{Mean life } \tau &= (456 \pm 5) \times 10^{-15} \text{ s} \\ c\tau &= 136.6 \text{ } \mu\text{m} \end{aligned}$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Xi_c^+ \rightarrow \Sigma^+ \bar{K}^*(892)^0$ seen in $\Xi_c^+ \rightarrow \Sigma^+ K^- \pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$ decays.

Ξ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
-----------------------	--------------------------------	-----------------------------------	----------------

Cabibbo-favored ($S = -2$) decays

$p 2K_S^0$	$(2.5 \pm 1.3) \times 10^{-3}$		766
$\Lambda \bar{K}^0 \pi^+$	—		852
$\Sigma(1385)^+ \bar{K}^0$	[a] $(2.9 \pm 2.0)\%$		746

$\Lambda K^- 2\pi^+$	$(9 \pm 4) \times 10^{-3}$		787
$\Lambda \bar{K}^*(892)^0 \pi^+$	$[a] < 5 \times 10^{-3}$	CL=90%	608
$\Sigma(1385)^+ K^- \pi^+$	$[a] < 6 \times 10^{-3}$	CL=90%	678
$\Sigma^+ K^- \pi^+$	$(2.7 \pm 1.2) \%$		810
$\Sigma^+ \bar{K}^*(892)^0$	$[a] (2.3 \pm 1.1) \%$		658
$\Sigma^0 K^- 2\pi^+$	$(8 \pm 5) \times 10^{-3}$		735
$\Xi^0 \pi^+$	$(1.6 \pm 0.8) \%$		876
$\Xi^- 2\pi^+$	$(2.9 \pm 1.3) \%$		851
$\Xi(1530)^0 \pi^+$	$[a] < 2.9 \times 10^{-3}$	CL=90%	749
$\Xi(1620)^0 \pi^+$	seen		—
$\Xi(1690)^0 \pi^+$	seen		644
$\Xi^0 \pi^+ \pi^0$	$(6.7 \pm 3.5) \%$		856
$\Xi^0 \pi^- 2\pi^+$	$(5.0 \pm 2.6) \%$		818
$\Xi^0 e^+ \nu_e$	$(7 \pm 4) \%$		884
$\Omega^- K^+ \pi^+$	$(2.0 \pm 1.5) \times 10^{-3}$		399

Cabibbo-suppressed decays

$p K^- \pi^+$	$(6.2 \pm 3.0) \times 10^{-3}$	S=1.5	944
$p \bar{K}^*(892)^0$	$[a] (3.3 \pm 1.7) \times 10^{-3}$		828
$\Sigma^+ \pi^+ \pi^-$	$(1.4 \pm 0.8) \%$		922
$\Sigma^- 2\pi^+$	$(5.1 \pm 3.4) \times 10^{-3}$		918
$\Sigma^+ K^+ K^-$	$(4.3 \pm 2.5) \times 10^{-3}$		579
$\Sigma^+ \phi$	$[a] < 3.2 \times 10^{-3}$	CL=90%	549
$\Xi(1690)^0 K^+, \Xi^0 \rightarrow \Sigma^+ K^-$	$< 1.3 \times 10^{-3}$	CL=90%	501
$p \phi(1020)$	$(1.2 \pm 0.6) \times 10^{-4}$		751



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2470.44 \pm 0.28 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 2.72 \pm 0.23 \text{ MeV} \quad (S = 1.1)$$

$$\text{Mean life } \tau = (153 \pm 6) \times 10^{-15} \text{ s} \quad (S = 2.4)$$

$$c\tau = 45.8 \text{ } \mu\text{m}$$

Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

Ξ_c^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor $\frac{P}{\text{MeV}/c}$
-----------------------	--------------------------------	---------------------------------------


Cabibbo-favored decays

$p K^- K^- \pi^+$	$(4.8 \pm 1.2) \times 10^{-3}$	1.1	676
$p K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(2.0 \pm 0.6) \times 10^{-3}$		413

$p K^- K^- \pi^+ (\text{no } \bar{K}^{*0})$	$(3.0 \pm 0.9) \times 10^{-3}$		676
ΛK_S^0	$(3.0 \pm 0.8) \times 10^{-3}$		906
$\Lambda K^- \pi^+$	$(1.45 \pm 0.33) \%$	1.1	856
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen		786
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen		703
$\Xi^- \pi^+$	$(1.43 \pm 0.32) \%$	1.1	875
$\Xi^- \pi^+ \pi^+ \pi^-$	$(4.8 \pm 2.3) \%$		816
$\Omega^- K^+$	$(4.2 \pm 1.0) \times 10^{-3}$		522
$\Xi^- e^+ \nu_e$	$(1.8 \pm 1.2) \%$		882

Cabibbo-suppressed decays

$\Lambda_c^+ \pi^-$	$(5.5 \pm 1.8) \times 10^{-3}$		115
$\Xi^- K^+$	$(3.9 \pm 1.2) \times 10^{-4}$		789
$\Lambda K^+ K^- (\text{no } \phi)$	$(4.1 \pm 1.4) \times 10^{-4}$		648
$\Lambda \phi$	$(4.9 \pm 1.5) \times 10^{-4}$		621



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.


$$\text{Mass } m = 2578.2 \pm 0.5 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c^{'+}} - m_{\Xi_c^+} = 110.5 \pm 0.4 \text{ MeV}$$

$$m_{\Xi_c^{'+}} - m_{\Xi_c^{\prime 0}} = -0.5 \pm 0.6 \text{ MeV}$$

The $\Xi_c^{'+} - \Xi_c^+$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'+}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \gamma$	seen	108



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2578.7 \pm 0.5 \text{ MeV}$$

$$m_{\Xi_c^{'0}} - m_{\Xi_c^0} = 108.3 \pm 0.4 \text{ MeV}$$

The $\Xi_c^{'0} - \Xi_c^0$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'0}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \gamma$	seen	106

$\Xi_c(2645)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\Xi_c(2645)^+ \text{ mass } m = 2645.10 \pm 0.30 \text{ MeV} \quad (S = 1.2)$$

$$\Xi_c(2645)^0 \text{ mass } m = 2646.16 \pm 0.25 \text{ MeV} \quad (S = 1.3)$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 174.67 \pm 0.09 \text{ MeV}$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.45 \pm 0.10 \text{ MeV}$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c(2645)^0} = -1.06 \pm 0.27 \text{ MeV} \quad (S = 1.1)$$

$$\Xi_c(2645)^+ \text{ full width } \Gamma = 2.14 \pm 0.19 \text{ MeV} \quad (S = 1.1)$$

$$\Xi_c(2645)^0 \text{ full width } \Gamma = 2.35 \pm 0.22 \text{ MeV}$$

$\Xi_c \pi$ is the only strong decay allowed to a Ξ_c resonance having this mass.

$\Xi_c(2645)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \pi^+$	seen	102
$\Xi_c^+ \pi^-$	seen	106

 $\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

J^P has not been measured; $\frac{1}{2}^-$ is the quark-model prediction.

$$\Xi_c(2790)^+ \text{ mass } = 2791.9 \pm 0.5 \text{ MeV}$$

$$\Xi_c(2790)^0 \text{ mass } = 2793.9 \pm 0.5 \text{ MeV}$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c'^0} = 213.20 \pm 0.22 \text{ MeV}$$

$$m_{\Xi_c(2790)^0} - m_{\Xi_c'^+} = 215.70 \pm 0.22 \text{ MeV}$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c(2790)^0} = -2.0 \pm 0.7 \text{ MeV}$$

$$\Xi_c(2790)^+ \text{ width } = 8.9 \pm 1.0 \text{ MeV}$$

$$\Xi_c(2790)^0 \text{ width } = 10.0 \pm 1.1 \text{ MeV}$$

$\Xi_c(2790)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c'^0 \pi$	seen	159

 $\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\Xi_c(2815)^+ \text{ mass } m = 2816.51 \pm 0.25 \text{ MeV} \quad (S = 1.2)$$

$$\Xi_c(2815)^0 \text{ mass } m = 2819.79 \pm 0.30 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.80 \pm 0.10 \text{ MeV}$$

$$\begin{aligned}
m_{\Xi_c(2815)^0} - m_{\Xi_c^0} &= 349.35 \pm 0.11 \text{ MeV} \\
m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0} &= -3.27 \pm 0.27 \text{ MeV} \\
\Xi_c(2815)^+ \text{ full width } \Gamma &= 2.43 \pm 0.26 \text{ MeV} \\
\Xi_c(2815)^0 \text{ full width } \Gamma &= 2.54 \pm 0.25 \text{ MeV}
\end{aligned}$$

The $\Xi_c \pi \pi$ modes are consistent with being entirely via $\Xi_c(2645) \pi$.

$\Xi_c(2815)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c' \pi$	seen	188
$\Xi_c(2645) \pi$	seen	102
$\Xi_c^0 \gamma$	seen	325

$\Xi_c(2970)$

$$I(J^P) = \frac{1}{2}(??)$$

was $\Xi_c(2980)$

$$\begin{aligned}
\Xi_c(2970)^+ \text{ } m &= 2964.3 \pm 1.5 \text{ MeV} \quad (S = 3.9) \\
\Xi_c(2970)^0 \text{ } m &= 2967.1 \pm 1.7 \text{ MeV} \quad (S = 6.7) \\
m_{\Xi_c(2970)^+} - m_{\Xi_c^+} &= 496.6 \pm 1.5 \text{ MeV} \quad (S = 3.7) \\
m_{\Xi_c(2970)^0} - m_{\Xi_c^0} &= 496.7 \pm 1.8 \text{ MeV} \quad (S = 5.3) \\
m_{\Xi_c(2970)^+} - m_{\Xi_c(2970)^0} &= -2.8 \pm 1.9 \text{ MeV} \quad (S = 4.8) \\
\Xi_c(2970)^+ \text{ width } \Gamma &= 20.9_{-3.5}^{+2.4} \text{ MeV} \quad (S = 1.2)
\end{aligned}$$

$\Xi_c(2970)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	223
$\Sigma_c(2455) \bar{K}$	seen	121
$\Lambda_c^+ \bar{K}$	not seen	410
$\Lambda_c^+ K^-$	seen	410
$\Xi_c 2\pi$	seen	381
$\Xi_c' \pi$	seen	—
$\Xi_c(2645) \pi$	seen	274

$\Xi_c(3055)$

$$I(J^P) = ?(??)$$

$$\begin{aligned}
\text{Mass } m &= 3055.9 \pm 0.4 \text{ MeV} \\
\text{Full width } \Gamma &= 7.8 \pm 1.9 \text{ MeV}
\end{aligned}$$

$\Xi_c(3055)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Sigma^{++} K^-$	seen	—
ΛD^+	seen	316

$\Xi_c(3080)$

$$I(J^P) = \frac{1}{2}(??)$$

$$\begin{aligned} \Xi_c(3080)^+ \quad m &= 3077.2 \pm 0.4 \text{ MeV} \\ \Xi_c(3080)^0 \quad m &= 3079.9 \pm 1.4 \text{ MeV} \quad (S = 1.3) \\ \Xi_c(3080)^+ \quad \text{width } \Gamma &= 3.6 \pm 1.1 \text{ MeV} \quad (S = 1.5) \\ \Xi_c(3080)^0 \quad \text{width } \Gamma &= 5.6 \pm 2.2 \text{ MeV} \end{aligned}$$

$\Xi_c(3080)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	415
$\Sigma_c(2455) \bar{K}$	seen	342
$\Sigma_c(2455)^{++} K^-$	seen	342
$\Sigma_c(2520)^{++} K^-$	seen	239
$\Sigma_c(2455) \bar{K} + \Sigma_c(2520) \bar{K}$	seen	—
$\Lambda_c^+ \bar{K}$	not seen	536
$\Lambda_c^+ \bar{K} \pi^+ \pi^-$	not seen	144
ΛD^+	seen	362

 Ω_c^0

$$I(J^P) = 0(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2695.2 \pm 1.7 \text{ MeV} \quad (S = 1.3)$$

$$\text{Mean life } \tau = (268 \pm 26) \times 10^{-15} \text{ s}$$

$$c\tau = 80 \text{ } \mu\text{m}$$

Ω_c^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
--	--------------------------------	------------------	-------------

No absolute branching fractions have been measured.
The following are branching *ratios* relative to $\Omega^- \pi^+$.

Cabibbo-favored ($S = -3$) decays — relative to $\Omega^- \pi^+$

$\Omega^- \pi^+$	DEFINED AS 1	821
$\Omega^- \pi^+ \pi^0$	1.80 ± 0.33	797
$\Omega^- \rho^+$	>1.3	90% 532
$\Omega^- \pi^- 2\pi^+$	0.31 ± 0.05	753
$\Omega^- e^+ \nu_e$	2.4 ± 1.2	829
$\Xi^0 \bar{K}^0$	1.64 ± 0.29	950
$\Xi^0 K^- \pi^+$	1.20 ± 0.18	901
$\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$	0.68 ± 0.16	764
$\Xi^- \bar{K}^0 \pi^+$	2.12 ± 0.28	895
$\Xi^- K^- 2\pi^+$	0.63 ± 0.09	830

$\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow$	0.21 ± 0.06	757
$\Xi^- \pi^+$		
$\Xi^- \bar{K}^{*0} \pi^+$	0.34 ± 0.11	653
$p K^- K^- \pi^+$	seen	864
$\Sigma^+ K^- K^- \pi^+$	< 0.32	90% 689
$\Lambda \bar{K}^0 \bar{K}^0$	1.72 ± 0.35	837

 $\Omega_c(2770)^0$

$$I(J^P) = 0(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2765.9 \pm 2.0 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Omega_c(2770)^0} - m_{\Omega_c^0} = 70.7^{+0.8}_{-0.9} \text{ MeV}$$

The $\Omega_c(2770)^0 - \Omega_c^0$ mass difference is too small for any strong decay to occur.

$\Omega_c(2770)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Omega_c^0 \gamma$	presumably 100%	70

 $\Omega_c(3000)^0$

$$I(J^P) = ?(?^?)$$

$$\text{Mass } m = 3000.41 \pm 0.22 \text{ MeV}$$

$$\text{Full width } \Gamma = 4.5 \pm 0.7 \text{ MeV}$$

$\Omega_c(3000)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	182

 $\Omega_c(3050)^0$

$$I(J^P) = ?(?^?)$$

$$\text{Mass } m = 3050.20 \pm 0.13 \text{ MeV}$$

$$\text{Full width } \Gamma < 1.2 \text{ MeV, CL} = 95\%$$

$\Omega_c(3050)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	279

 $\Omega_c(3065)^0$

$$I(J^P) = ?(?^?)$$

$$\text{Mass } m = 3065.46 \pm 0.28 \text{ MeV}$$

$$\text{Full width } \Gamma = 3.5 \pm 0.4 \text{ MeV}$$

$\Omega_c(3065)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	303

$\Omega_c(3090)^0$

$$I(J^P) = ?(?^?)$$

Mass $m = 3090.0 \pm 0.5$ MeVFull width $\Gamma = 8.7 \pm 1.3$ MeV

$\Omega_c(3090)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	339

$\Omega_c(3120)^0$

$$I(J^P) = ?(?^?)$$

Mass $m = 3119.1 \pm 1.0$ MeVFull width $\Gamma < 2.6$ MeV, CL = 95%

$\Omega_c(3120)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	379

NOTES

- [a] This branching fraction includes all the decay modes of the final-state resonance.
- [b] See AALTONEN 11H, Fig. 8, for the calculated ratio of $\Lambda_c^+ \pi^0 \pi^0$ and $\Lambda_c^+ \pi^+ \pi^-$ partial widths as a function of the $\Lambda_c(2595)^+ - \Lambda_c^+$ mass difference. At our value of the mass difference, the ratio is about 4.
- [c] A test that the isospin is indeed 0, so that the particle is indeed a Λ_c^+ .