

CHARMED MESONS

($C = \pm 1$)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \text{ similarly for } D^{*'}s$$

$D^\pm$

$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1869.66 \pm 0.05 \text{ MeV}$$

$$\text{Mean life } \tau = (1040 \pm 7) \times 10^{-15} \text{ s}$$

$$c\tau = 311.8 \text{ } \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.096 \pm 0.004 \text{ } [a]$$

$$\Gamma(c \rightarrow D^{*(2010)^+} \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_L^0 e^\pm \nu) = (-0.6 \pm 1.6)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.41 \pm 0.09)\%$$

$$A_{CP}(K_L^0 K^\pm) \text{ in } D^\pm \rightarrow K_L^0 K^\pm = (-4.2 \pm 3.4) \times 10^{-2}$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.18 \pm 0.16)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (-0.3 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (-0.1 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \eta) \text{ in } D^\pm \rightarrow K_S^0 \pi^\pm \eta = (-0.9 \pm 3.1) \times 10^{-2}$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.0 \pm 1.2)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^- \pi^0) \text{ in } D^\pm \rightarrow K^\pm \pi^+ \pi^- \pi^0 = -0.04 \pm 0.06$$

$$A_{CP}(\pi^\pm \pi^0) = (2.4 \pm 1.2)\%$$

$$A_{CP}(\pi^\pm \eta) = (1.0 \pm 1.5)\% \quad (S = 1.4)$$

$$A_{CP}(\pi^\pm \eta'(958)) = (-0.6 \pm 0.7)\%$$

$$A_{CP}(\bar{K}^0 / K^0 K^\pm) = (0.11 \pm 0.17)\%$$

$$A_{CP}(K_S^0 K^\pm) = (-0.01 \pm 0.07)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_S^0 K^\pm \pi^0 = (1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K_L^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_L^0 K^\pm \pi^0 = (-1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.37 \pm 0.29)\%$$

$$A_{CP}(K^\pm K^{*0}) = (-0.3 \pm 0.4)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.01 \pm 0.09)\% \quad (S = 1.8)$$

$$A_{CP}(K^\pm K_0^{*}(1430)^0) = (8_{-6}^{+7})\%$$

$$A_{CP}(K^\pm K_2^{*}(1430)^0) = (43_{-26}^{+20})\%$$

$$A_{CP}(K^\pm K_0^{*}(700)) = (-12_{-13}^{+18})\%$$

$$A_{CP}(a_0(1450)^0 \pi^\pm) = (-19_{-16}^{+14})\%$$

$$A_{CP}(\phi(1680) \pi^\pm) = (-9 \pm 26)\%$$

$$\begin{aligned}
 A_{CP}(\pi^+\pi^-\pi^\pm) &= (-2 \pm 4)\% \\
 A_{CP}(\pi^+\pi^-\pi^\pm\eta) \text{ in } D^\pm \rightarrow \pi^+\pi^-\pi^\pm\eta &= (3 \pm 5) \times 10^{-2} \\
 A_{CP}(K_S^0 K^\pm \pi^+\pi^-) &= (-4 \pm 7)\% \\
 A_{CP}(K^\pm \pi^0) &= (-4 \pm 11)\%
 \end{aligned}$$

χ^2 tests of CP-violation (CPV)

$$\begin{aligned}
 \text{Local CPV in } D^\pm \rightarrow \pi^+\pi^-\pi^\pm &= 78.1\% \\
 \text{Local CPV in } D^\pm \rightarrow K^+K^-\pi^\pm &= 31\%
 \end{aligned}$$

CP violating asymmetries of P-odd (T-odd) moments

$$A_T(K_S^0 K^\pm \pi^+\pi^-) = (-12 \pm 11) \times 10^{-3} [b]$$

D^+ form factors

$$\begin{aligned}
 f_+(0)|V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= 0.719 \pm 0.011 \quad (S = 1.6) \\
 r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -2.13 \pm 0.14 \\
 r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -3 \pm 12 \quad (S = 1.5) \\
 f_+(0)|V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell &= 0.1407 \pm 0.0025 \\
 r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -2.00 \pm 0.13 \\
 r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -4 \pm 5 \\
 f_+(0)|V_{cd}| \text{ in } D^+ \rightarrow \eta \ell^+ \nu_\ell \ (\ell = e \text{ or } \nu) &= (8.4 \pm 0.4) \times 10^{-2} \\
 r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e &= -5.3 \pm 2.7 \quad (S = 1.9) \\
 r_\nu \equiv V(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e &= 1.24 \pm 0.11 \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e &= 1.06 \pm 0.16 \\
 r_\nu \equiv V(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e &= 1.64 \pm 0.10 \quad (S = 1.2) \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e &= 0.84 \pm 0.06 \\
 r_\nu \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.49 \pm 0.05 \quad (S = 2.1) \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.802 \pm 0.021 \\
 r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.0 \pm 0.4 \\
 \Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.13 \pm 0.08 \\
 \Gamma_+/\Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.22 \pm 0.06 \quad (S = 1.6)
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as $\bar{K}^0$ modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07 $\pm$ 0.30) %		—
μ^+ anything	(17.6 $\pm$ 3.2) %		—
K^- anything	(25.7 $\pm$ 1.4) %		—
$\overline{K}^0$ anything + K^0 anything	(61 $\pm$ 5) %		—
K^+ anything	(5.9 $\pm$ 0.8) %		—

$K^*(892)^-$ anything	(6 ± 5) %	—
$\bar{K}^*(892)^0$ anything	(23 ± 5) %	—
$K^*(892)^0$ anything	< 6.6 %	CL=90% —
η anything	(6.3 ± 0.7) %	—
η' anything	(1.04 ± 0.18) %	—
ϕ anything	(1.12 ± 0.04) %	—

Leptonic and semileptonic modes

$e^+ \nu_e$	< 8.8 × 10 ⁻⁶ CL=90%	935
$\gamma e^+ \nu_e$	< 3.0 × 10 ⁻⁵ CL=90%	935
$\mu^+ \nu_\mu$	(3.74 ± 0.17) × 10 ⁻⁴	932
$\tau^+ \nu_\tau$	(1.20 ± 0.27) × 10 ⁻³	90
$\bar{K}^0 e^+ \nu_e$	(8.73 ± 0.10) %	869
$\bar{K}^0 \mu^+ \nu_\mu$	(8.76 ± 0.19) %	865
$K^- \pi^+ e^+ \nu_e$	(4.02 ± 0.18) % S=3.2	864
$\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.77 ± 0.17) %	722
$(K^- \pi^+) [0.8-1.0] \text{ GeV } e^+ \nu_e$	(3.39 ± 0.09) %	864
$(K^- \pi^+)_{S\text{-wave}} e^+ \nu_e$	(2.28 ± 0.11) × 10 ⁻³	—
$\bar{K}^*(1410)^0 e^+ \nu_e,$	< 6 × 10 ⁻³ CL=90%	—
$\bar{K}^*(1410)^0 \rightarrow K^- \pi^+$		
$\bar{K}_2^*(1430)^0 e^+ \nu_e,$	< 5 × 10 ⁻⁴ CL=90%	—
$\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$		
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 × 10 ⁻³ CL=90%	864
$\bar{K}^*(892)^0 e^+ \nu_e$	(5.40 ± 0.10) % S=1.1	722
$K^- \pi^+ \mu^+ \nu_\mu$	(3.65 ± 0.34) %	851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu,$	(3.52 ± 0.10) %	717
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(1.9 ± 0.5) × 10 ⁻³	851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(5.27 ± 0.15) %	717
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.5 × 10 ⁻³ CL=90%	825
$\bar{K}_1(1270)^0 e^+ \nu_e, \bar{K}_1^0 \rightarrow K^- \pi^+ \pi^0$	(1.06 ± 0.15) × 10 ⁻³	—
$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.3 × 10 ⁻⁴ CL=90%	380
$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.5 × 10 ⁻³ CL=90%	105
$\pi^0 e^+ \nu_e$	(3.72 ± 0.17) × 10 ⁻³ S=2.0	930
$\pi^0 \mu^+ \nu_\mu$	(3.50 ± 0.15) × 10 ⁻³	927
$\eta e^+ \nu_e$	(1.11 ± 0.07) × 10 ⁻³	855
$\eta \mu^+ \nu_\mu$	(1.04 ± 0.11) × 10 ⁻³	851
$\pi^- \pi^+ e^+ \nu_e$	(2.45 ± 0.10) × 10 ⁻³	924
$f_0(500)^0 e^+ \nu_e, f_0(500)^0 \rightarrow \pi^+ \pi^-$	(6.3 ± 0.5) × 10 ⁻⁴	—
$\rho^0 e^+ \nu_e$	(2.18 ± 0.17 / 0.25) × 10 ⁻³	774

$\rho^0 \mu^+ \nu_\mu$	$(2.4 \pm 0.4) \times 10^{-3}$	770
$\omega e^+ \nu_e$	$(1.69 \pm 0.11) \times 10^{-3}$	771
$\omega \mu^+ \nu_\mu$	$(1.77 \pm 0.21) \times 10^{-3}$	767
$\eta'(958) e^+ \nu_e$	$(2.0 \pm 0.4) \times 10^{-4}$	690
$a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0$	$(1.7 \pm_{-0.7}^{+0.8}) \times 10^{-4}$	—
$b_1(1235)^0 e^+ \nu_e, b_1^0 \rightarrow \omega \pi^0$	$< 1.75 \times 10^{-4}$ CL=90%	—
$\phi e^+ \nu_e$	$< 1.3 \times 10^{-5}$ CL=90%	657
$D^0 e^+ \nu_e$	$< 1.0 \times 10^{-4}$ CL=90%	5

Hadronic modes with a $\bar{K}$ or $\bar{K}K\bar{K}$

$K_S^0 \pi^+$	$(1.562 \pm 0.031) \%$	S=1.7	863
$K_L^0 \pi^+$	$(1.46 \pm 0.05) \%$		863
$K^- 2\pi^+$	[c] $(9.38 \pm 0.16) \%$	S=1.6	846
$(K^- \pi^+)_{S\text{-wave}} \pi^+$	$(7.52 \pm 0.17) \%$		846
$\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[d] $(1.25 \pm 0.06) \%$		382
$\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.04 \pm 0.12) \%$		714
$\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+$	not seen		381
$\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[d] $(2.3 \pm 0.7) \times 10^{-4}$		371
$\bar{K}^*(1680)^0 \pi^+, \bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[d] $(2.2 \pm 1.1) \times 10^{-4}$		58
$K^-(2\pi^+)_{I=2}$	$(1.45 \pm 0.26) \%$		—
$K_S^0 \pi^+ \pi^0$	[c] $(7.36 \pm 0.21) \%$		845
$K_S^0 \rho^+$	$(6.14 \pm_{-0.35}^{+0.60}) \%$		677
$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.5 \pm_{-1.4}^{+1.2}) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	$(2.64 \pm 0.32) \times 10^{-3}$		714
$\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0$	$(2.7 \pm 0.9) \times 10^{-3}$		—
$\bar{K}_0^*(1680)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0$	$(10 \pm_{-10}^{+7}) \times 10^{-4}$		—
$\bar{K}^0 \pi^+, \bar{K}^0 \rightarrow K_S^0 \pi^0$	$(6 \pm_{-4}^{+5}) \times 10^{-3}$		—
$K_S^0 \pi^+ \pi^0$ nonresonant	$(3 \pm 4) \times 10^{-3}$		845
$K_S^0 \pi^+ \pi^0$ nonresonant and $\bar{K}^0 \pi^+$	$(1.37 \pm_{-0.40}^{+0.21}) \%$		—
$(K_S^0 \pi^0)_{S\text{-wave}} \pi^+$	$(1.27 \pm_{-0.33}^{+0.27}) \%$		845

$K_S^0 \pi^+ \eta$	(1.31 $\pm$ 0.05) %	722
$K_S^0 \pi^+ \eta'(958)$	(1.90 $\pm$ 0.21) $\times 10^{-3}$	481
$K^- 2\pi^+ \pi^0$	[e] (6.25 $\pm$ 0.18) %	817
$K_S^0 2\pi^+ \pi^-$	[e] (3.10 $\pm$ 0.09) %	814
$K^- 2\pi^+ \eta$	(1.35 $\pm$ 0.12) $\times 10^{-3}$	657
$K_S^0 \pi^+ \pi^0 \eta$	(1.22 $\pm$ 0.25) $\times 10^{-3}$	657
$K^- 3\pi^+ \pi^-$	[c] (5.7 $\pm$ 0.5) $\times 10^{-3}$ S=1.1	772
$\bar{K}^*(892)^0 2\pi^+ \pi^-$,	(1.2 $\pm$ 0.4) $\times 10^{-3}$	645
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$\bar{K}^*(892)^0 \rho^0 \pi^+$,	(2.3 $\pm$ 0.4) $\times 10^{-3}$	239
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$\bar{K}^*(892)^0 a_1(1260)^+$	[f] (9.3 $\pm$ 1.9) $\times 10^{-3}$	†
$K^- \rho^0 2\pi^+$	(1.72 $\pm$ 0.28) $\times 10^{-3}$	524
$K^- 3\pi^+ \pi^-$ nonresonant	(4.0 $\pm$ 2.9) $\times 10^{-4}$	772
$K^+ 2K_S^0$	(2.54 $\pm$ 0.13) $\times 10^{-3}$	545
$K^+ K^- K_S^0 \pi^+$	(2.4 $\pm$ 0.5) $\times 10^{-4}$	436

Pionic modes

$\pi^+ \pi^0$	(1.247 $\pm$ 0.033) $\times 10^{-3}$	925
$2\pi^+ \pi^-$	(3.27 $\pm$ 0.18) $\times 10^{-3}$	909
$\rho^0 \pi^+$	(8.3 $\pm$ 1.5) $\times 10^{-4}$	767
$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	(1.83 $\pm$ 0.16) $\times 10^{-3}$	909
$\sigma \pi^+$, $\sigma \rightarrow \pi^+ \pi^-$	(1.38 $\pm$ 0.12) $\times 10^{-3}$	—
$f_0(980) \pi^+$,	(1.56 $\pm$ 0.33) $\times 10^{-4}$	669
$f_0(980) \rightarrow \pi^+ \pi^-$		
$f_0(1370) \pi^+$,	(8 $\pm$ 4) $\times 10^{-5}$	—
$f_0(1370) \rightarrow \pi^+ \pi^-$		
$f_2(1270) \pi^+$,	(5.0 $\pm$ 0.9) $\times 10^{-4}$	485
$f_2(1270) \rightarrow \pi^+ \pi^-$		
$\rho(1450)^0 \pi^+$,	< 8 $\times 10^{-5}$ CL=95%	338
$\rho(1450)^0 \rightarrow \pi^+ \pi^-$		
$f_0(1500) \pi^+$,	(1.1 $\pm$ 0.4) $\times 10^{-4}$	—
$f_0(1500) \rightarrow \pi^+ \pi^-$		
$f_0(1710) \pi^+$,	< 5 $\times 10^{-5}$ CL=95%	—
$f_0(1710) \rightarrow \pi^+ \pi^-$		
$f_0(1790) \pi^+$,	< 7 $\times 10^{-5}$ CL=95%	—
$f_0(1790) \rightarrow \pi^+ \pi^-$		
$(\pi^+ \pi^+)_{S\text{-wave}} \pi^-$	< 1.2 $\times 10^{-4}$ CL=95%	909
$2\pi^+ \pi^-$ nonresonant	< 1.1 $\times 10^{-4}$ CL=95%	909
$\pi^+ 2\pi^0$	(4.7 $\pm$ 0.4) $\times 10^{-3}$	910
$2\pi^+ \pi^- \pi^0$	(1.16 $\pm$ 0.08) %	883
$3\pi^+ 2\pi^-$	(1.66 $\pm$ 0.16) $\times 10^{-3}$ S=1.1	845
$\eta \pi^+$	(3.77 $\pm$ 0.09) $\times 10^{-3}$	848

$\eta\pi^+\pi^0$	$(2.05 \pm 0.35) \times 10^{-3}$	S=2.2	831
$\eta 2\pi^+\pi^-$	$(3.41 \pm 0.20) \times 10^{-3}$		798
$\eta\pi^+2\pi^0$	$(3.20 \pm 0.33) \times 10^{-3}$		801
$\eta\eta\pi^+$	$(2.96 \pm 0.26) \times 10^{-3}$		700
$\omega\pi^+$	$(2.8 \pm 0.6) \times 10^{-4}$		764
$\omega\pi^+\pi^0$	$(3.9 \pm 0.9) \times 10^{-3}$		742
$\eta'(958)\pi^+$	$(4.97 \pm 0.19) \times 10^{-3}$		681
$\eta'(958)\pi^+\pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$		654

Hadronic modes with a $K\bar{K}$ pair

$K^+K_S^0$	$(3.04 \pm 0.09) \times 10^{-3}$	S=2.2	793
$K_L^0K^+$	$(3.21 \pm 0.16) \times 10^{-3}$		793
$K_S^0K^+\pi^0$	$(5.07 \pm 0.30) \times 10^{-3}$		744
$K_L^0K^+\pi^0$	$(5.24 \pm 0.31) \times 10^{-3}$		744
$K^+K^-\pi^+$	[c] $(9.68 \pm 0.18) \times 10^{-3}$		744
$K^+K^-\pi^+\pi^0$	$(6.62 \pm 0.32) \times 10^{-3}$		682
$\phi\pi^+$	$(5.70 \pm 0.14) \times 10^{-3}$		647
$\phi\pi^+, \phi \rightarrow K^+K^-$	$(2.69 \pm_{-0.08}^{+0.07}) \times 10^{-3}$		647
$K^+\bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(2.49 \pm_{-0.13}^{+0.08}) \times 10^{-3}$		613
$K^+\bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow K^-\pi^+$	$(1.82 \pm 0.35) \times 10^{-3}$		—
$K^+\bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^-\pi^+$	$(1.6 \pm_{-0.8}^{+1.2}) \times 10^{-4}$		—
$K^+\bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^-\pi^+$	$(6.8 \pm_{-2.1}^{+3.5}) \times 10^{-4}$		—
$a_0(1450)^0\pi^+, a_0^0 \rightarrow K^+K^-$	$(4.5 \pm_{-1.8}^{+7.0}) \times 10^{-4}$		—
$\phi(1680)\pi^+, \phi \rightarrow K^+K^-$	$(4.9 \pm_{-1.9}^{+4.0}) \times 10^{-5}$		—
$K_S^0K_S^0\pi^+$	$(2.70 \pm 0.13) \times 10^{-3}$		741
$K_S^0K_S^0\pi^+\pi^0$	$(1.34 \pm 0.21) \times 10^{-3}$		679
$K_S^0K^+\eta$	$(1.8 \pm 0.5) \times 10^{-4}$		516
$K^+K_S^0\pi^+\pi^-$	$(1.89 \pm 0.13) \times 10^{-3}$		678
$K_S^0K^+\pi^0\pi^0$	$(5.8 \pm 1.3) \times 10^{-4}$		683
$K_S^0K^-2\pi^+$	$(2.27 \pm 0.13) \times 10^{-3}$		678
$K^+K^-2\pi^+\pi^-$	$(2.3 \pm 1.2) \times 10^{-4}$		601

A few poorly measured branching fractions:

$\phi\pi^+\pi^0$	$(2.3 \pm 1.0) \%$		619
$\phi\rho^+$	$< 1.5 \%$	CL=90%	260
$K^+K^-\pi^+\pi^0$ non- ϕ	$(1.5 \pm_{-0.6}^{+0.7}) \%$		682
$K^*(892)^+K_S^0$	$(1.7 \pm 0.8) \%$		612

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(2.08 \pm 0.21) \times 10^{-4}$	$S=1.4$	864
$K^+ \eta$	$(1.25 \pm 0.16) \times 10^{-4}$	$S=1.1$	776
$K^+ \eta'(958)$	$(1.85 \pm 0.20) \times 10^{-4}$		571
$K^+ \pi^+ \pi^-$	$(4.91 \pm 0.09) \times 10^{-4}$		846
$K^+ \rho^0$	$(1.9 \pm 0.5) \times 10^{-4}$		679
$K^{*}(892)^0 \pi^+, K^{*}(892)^0 \rightarrow$	$(2.3 \pm 0.4) \times 10^{-4}$		714
$K^+ \pi^-$			
$K^+ f_0(980), f_0(980) \rightarrow$	$(4.4 \pm 2.6) \times 10^{-5}$		—
$\pi^+ \pi^-$			
$K_2^{*}(1430)^0 \pi^+, K_2^{*}(1430)^0 \rightarrow$	$(3.9 \pm 2.7) \times 10^{-5}$		—
$K^+ \pi^-$			
$K^+ \pi^+ \pi^-$ nonresonant	not seen		846
$K^+ \pi^+ \pi^- \pi^0$	$(1.21 \pm 0.09) \times 10^{-3}$		817
$K^+ \omega$	$(5.7 \pm 2.5) \times 10^{-5}$		675
$2K^+ K^-$	$(6.14 \pm 0.11) \times 10^{-5}$		550
$\phi(1020)^0 K^+$	$< 2.1 \times 10^{-5}$	$CL=90\%$	—
$K^+ \phi(1020), \phi \rightarrow K^+ K^-$	$(4.4 \pm 0.6) \times 10^{-6}$		—
$K^+ (K^+ K^-)_{S-wave}$	$(5.77 \pm 0.12) \times 10^{-5}$		550

 $\Delta C = 1$ weak neutral current (C1) modes, or Lepton Family number (LF), or Lepton number (L), or Baryon number (B) violating modes

$\pi^+ e^+ e^-$	C1	< 1.1	$\times 10^{-6} CL=90\%$	930
$\pi^+ \pi^0 e^+ e^-$		< 1.4	$\times 10^{-5} CL=90\%$	925
$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[g]	$(1.7 \pm 1.4) \times 10^{-6}$		—
$\pi^+ \mu^+ \mu^-$	C1	< 7.3	$\times 10^{-8} CL=90\%$	918
$\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$	[g]	$(1.8 \pm 0.8) \times 10^{-6}$		—
$\rho^+ \mu^+ \mu^-$	C1	< 5.6	$\times 10^{-4} CL=90\%$	757
$K^+ e^+ e^-$	[h]	< 1.0	$\times 10^{-6} CL=90\%$	870
$K^+ \pi^0 e^+ e^-$		< 1.5	$\times 10^{-5} CL=90\%$	864
$K_S^0 \pi^+ e^+ e^-$		< 2.6	$\times 10^{-5} CL=90\%$	—
$K_S^0 K^+ e^+ e^-$		< 1.1	$\times 10^{-5} CL=90\%$	—
$K^+ \mu^+ \mu^-$	[h]	< 4.3	$\times 10^{-6} CL=90\%$	856
$\pi^+ e^+ \mu^-$	LF	< 2.9	$\times 10^{-6} CL=90\%$	927
$\pi^+ e^- \mu^+$	LF	< 3.6	$\times 10^{-6} CL=90\%$	927
$K^+ e^+ \mu^-$	LF	< 1.2	$\times 10^{-6} CL=90\%$	866
$K^+ e^- \mu^+$	LF	< 2.8	$\times 10^{-6} CL=90\%$	866
$\pi^- 2e^+$	L	< 1.1	$\times 10^{-6} CL=90\%$	930
$\pi^- 2\mu^+$	L	< 2.2	$\times 10^{-8} CL=90\%$	918
$\pi^- e^+ \mu^+$	L	< 2.0	$\times 10^{-6} CL=90\%$	927
$\rho^- 2\mu^+$	L	< 5.6	$\times 10^{-4} CL=90\%$	757
$K^- 2e^+$	L	< 9	$\times 10^{-7} CL=90\%$	870
$K_S^0 \pi^- 2e^+$		< 3.3	$\times 10^{-6} CL=90\%$	863

$K^- \pi^0 2e^+$		< 8.5	$\times 10^{-6} \text{CL}=90\%$	864
$K^- 2\mu^+$	L	< 1.0	$\times 10^{-5} \text{CL}=90\%$	856
$K^- e^+ \mu^+$	L	< 1.9	$\times 10^{-6} \text{CL}=90\%$	866
$K^*(892)^- 2\mu^+$	L	< 8.5	$\times 10^{-4} \text{CL}=90\%$	703
Λe^+	L, B	< 1.1	$\times 10^{-6} \text{CL}=90\%$	602
$\bar{\Lambda} e^+$	L, B	< 6.5	$\times 10^{-7} \text{CL}=90\%$	602
$\Sigma^0 e^+$	L, B	< 1.7	$\times 10^{-6} \text{CL}=90\%$	554
$\bar{\Sigma}^0 e^+$	L, B	< 1.3	$\times 10^{-6} \text{CL}=90\%$	554

 D^0

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1864.84 \pm 0.05 \text{ MeV}$ $m_{D^\pm} - m_{D^0} = 4.822 \pm 0.015 \text{ MeV}$ Mean life $\tau = (410.1 \pm 1.5) \times 10^{-15} \text{ s}$ $c\tau = 122.9 \mu\text{m}$ **Mixing and related parameters**

$$|m_{D_1^0} - m_{D_2^0}| = (0.95^{+0.41}_{-0.44}) \times 10^{10} \hbar \text{ s}^{-1}$$

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.29^{+0.14}_{-0.18}) \times 10^{-2}$$

$$|q/p| = 0.92^{+0.12}_{-0.09}$$

$$A_\Gamma = (-0.125 \pm 0.526) \times 10^{-3}$$

$$\phi^{K_S^0 \pi \pi} = -0.09^{+0.10}_{-0.13}$$

$$K^+ \pi^- \text{ relative strong phase: } \cos \delta = 0.97 \pm 0.11$$

$$K^- \pi^+ \pi^0 \text{ coherence factor } R_{K \pi \pi^0} = 0.82 \pm 0.06$$

$$K^- \pi^+ \pi^0 \text{ average relative strong phase } \delta^{K \pi \pi^0} = (199 \pm 14)^\circ$$

$$K^- \pi^- 2\pi^+ \text{ coherence factor } R_{K 3\pi} = 0.53^{+0.18}_{-0.21}$$

$$K^- \pi^- 2\pi^+ \text{ average relative strong phase } \delta^{K 3\pi} = (125^{+22}_{-14})^\circ$$

$$D^0 \rightarrow K^- \pi^- 2\pi^+, R_{K 3\pi} (y \cos \delta^{K 3\pi} - x \sin \delta^{K 3\pi}) = (-3.0 \pm 0.7) \times 10^{-3} \text{ TeV}^{-1}$$

$$K_S^0 K^+ \pi^- \text{ coherence factor } R_{K_S^0 K \pi} = 0.70 \pm 0.08$$

$$K_S^0 K^+ \pi^- \text{ average relative strong phase } \delta^{K_S^0 K \pi} = (0 \pm 16)^\circ$$

$$K^* K \text{ coherence factor } R_{K^* K} = 0.94 \pm 0.12$$

$$K^* K \text{ average relative strong phase } \delta^{K^* K} = (-17 \pm 18)^\circ$$

CP-violation decay-rate asymmetries (labeled by the D^0 decay)

$$A_{CP}(K^+ K^-) = (-0.07 \pm 0.11)\%$$

$$A_{CP}(2K_S^0) = (0.4 \pm 1.4)\%$$

$$A_{CP}(\pi^+ \pi^-) = (0.13 \pm 0.14)\%$$

$$A_{CP}(\pi^0 \pi^0) = (0.0 \pm 0.6)\%$$

$$A_{CP}(\rho \gamma) = (6 \pm 15) \times 10^{-2}$$

$$\begin{aligned}
A_{CP}(\phi\gamma) &= (-9 \pm 7) \times 10^{-2} \\
A_{CP}(\overline{K}^*(892)^0\gamma) &= (-0.3 \pm 2.0) \times 10^{-2} \\
A_{CP}(\pi^+\pi^-\pi^0) &= (0.3 \pm 0.4)\% \\
A_{CP}(\rho(770)^+\pi^-\pi^0) &= (1.2 \pm 0.9)\% [i] \\
A_{CP}(\rho(770)^0\pi^0\pi^0) &= (-3.1 \pm 3.0)\% [i] \\
A_{CP}(\rho(770)^-\pi^+\pi^0) &= (-1.0 \pm 1.7)\% [i] \\
A_{CP}(\rho(1450)^+\pi^-\pi^0) &= (0 \pm 70)\% [i] \\
A_{CP}(\rho(1450)^0\pi^0\pi^0) &= (-20 \pm 40)\% [i] \\
A_{CP}(\rho(1450)^-\pi^+\pi^0) &= (6 \pm 9)\% [i] \\
A_{CP}(\rho(1700)^+\pi^-\pi^0) &= (-5 \pm 14)\% [i] \\
A_{CP}(\rho(1700)^0\pi^0\pi^0) &= (13 \pm 9)\% [i] \\
A_{CP}(\rho(1700)^-\pi^+\pi^0) &= (8 \pm 11)\% [i] \\
A_{CP}(f_0(980)\pi^0\pi^0) &= (0 \pm 35)\% [i] \\
A_{CP}(f_0(1370)\pi^0\pi^0) &= (25 \pm 18)\% [i] \\
A_{CP}(f_0(1500)\pi^0\pi^0) &= (0 \pm 18)\% [i] \\
A_{CP}(f_0(1710)\pi^0\pi^0) &= (0 \pm 24)\% [i] \\
A_{CP}(f_2(1270)\pi^0\pi^0) &= (-4 \pm 6)\% [i] \\
A_{CP}(\sigma(400)\pi^0\pi^0) &= (6 \pm 8)\% [i] \\
A_{CP}(\text{nonresonant } \pi^+\pi^-\pi^0) &= (-13 \pm 23)\% [i] \\
A_{CP}(a_1(1260)^+\pi^-\pi^0) &= (5 \pm 6)\% \\
A_{CP}(a_1(1260)^-\pi^+\pi^0) &= (14 \pm 18)\% \\
A_{CP}(\pi(1300)^+\pi^-\pi^0) &= (-2 \pm 15)\% \\
A_{CP}(\pi(1300)^-\pi^+\pi^0) &= (-6 \pm 30)\% \\
A_{CP}(a_1(1640)^+\pi^-\pi^0) &= (9 \pm 26)\% \\
A_{CP}(\pi_2(1670)^+\pi^-\pi^0) &= (7 \pm 18)\% \\
A_{CP}(\sigma f_0(1370)\pi^0) &= (-15 \pm 19)\% \\
A_{CP}(\sigma \rho(770)^0\pi^0) &= (3 \pm 27)\% \\
A_{CP}(2\rho(770)^0\pi^0) &= (-6 \pm 6)\% \\
A_{CP}(2f_2(1270)\pi^0) &= (-28 \pm 24)\% \\
A_{CP}(\pi^+\pi^-\pi^0\eta) \text{ in } D^0, \overline{D}^0 &\rightarrow \pi^+\pi^-\pi^0\eta = (-6 \pm 6) \times 10^{-2} \\
A_{CP}(K^+K^-\pi^0) &= (-1.0 \pm 1.7)\% \\
A_{CP}(K^*(892)^+K^-\pi^0) &= (-0.9 \pm 1.3)\% [i] \\
A_{CP}(K^*(1410)^+K^-\pi^0) &= (-21 \pm 24)\% [i] \\
A_{CP}((K^+\pi^0)_{S\text{-wave}}K^-\pi^0) &= (7 \pm 15)\% [i] \\
A_{CP}(\phi(1020)\pi^0\pi^0) &= (1.1 \pm 2.2)\% [i] \\
A_{CP}(f_0(980)\pi^0\pi^0) &= (-3 \pm 19)\% [i] \\
A_{CP}(a_0(980)^0\pi^0\pi^0) &= (-5 \pm 16)\% [i] \\
A_{CP}(f'_2(1525)\pi^0\pi^0) &= (0 \pm 160)\% [i] \\
A_{CP}(K^*(892)^-K^+\pi^0) &= (-5 \pm 4)\% [i] \\
A_{CP}(K^*(1410)^-K^+\pi^0) &= (-17 \pm 29)\% [i] \\
A_{CP}((K^-\pi^0)_{S\text{-wave}}K^+\pi^0) &= (-10 \pm 40)\% [i] \\
A_{CP}(K_S^0\pi^0) &= (-0.20 \pm 0.17)\%
\end{aligned}$$

$$\begin{aligned}
A_{CP}(K_S^0 \eta) &= (0.5 \pm 0.5)\% \\
A_{CP}(K_S^0 \eta') &= (1.0 \pm 0.7)\% \\
A_{CP}(K_S^0 \phi) &= (-3 \pm 9)\% \\
A_{CP}(K^- \pi^+) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+ \pi^-) &= (-0.9 \pm 1.4)\% \\
A_{CP}(D_{CP}(\pm 1) \rightarrow K^\mp \pi^\pm) &= (12.7 \pm 1.5)\% \\
A_{CP}(K^- \pi^+ \pi^0) &= (0.1 \pm 0.5)\% \\
A_{CP}(K^+ \pi^- \pi^0) &= (0 \pm 5)\% \\
A_{CP}(K_S^0 \pi^+ \pi^-) &= (-0.1 \pm 0.8)\% \\
A_{CP}(K^\mp \pi^\pm \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K^\mp \pi^\pm \eta &= (-1.9 \pm 1.6) \times 10^{-2} \\
A_{CP}(K_S^0 \pi^0 \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K_S^0 \pi^0 \eta &= (-3.9 \pm 3.3) \times 10^{-2} \\
A_{CP}(K^\mp \pi^\pm \pi^0 \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K^\mp \pi^\pm \pi^0 \eta &= (-8 \pm 5) \times 10^{-2} \\
A_{CP}(K^*(892)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (0.4 \pm 0.5)\% \\
A_{CP}(K^*(892)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (1 \pm 6)\% \\
A_{CP}(\bar{K}^0 \rho^0 \rightarrow K_S^0 \pi^+ \pi^-) &= (-0.1 \pm 0.5)\% \\
A_{CP}(\bar{K}^0 \omega \rightarrow K_S^0 \pi^+ \pi^-) &= (-13 \pm 7)\% \\
A_{CP}(\bar{K}^0 f_0(980) \rightarrow K_S^0 \pi^+ \pi^-) &= (-0.4 \pm 2.7)\% \\
A_{CP}(\bar{K}^0 f_2(1270) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 5)\% \\
A_{CP}(\bar{K}^0 f_0(1370) \rightarrow K_S^0 \pi^+ \pi^-) &= (-1 \pm 9)\% \\
A_{CP}(\bar{K}^0 \rho^0(1450) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 10)\% \\
A_{CP}(\bar{K}^0 f_0(600) \rightarrow K_S^0 \pi^+ \pi^-) &= (-3 \pm 5)\% \\
A_{CP}(K^*(1410)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (-2 \pm 9)\% \\
A_{CP}(K_0^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (4 \pm 4)\% \\
A_{CP}(K_0^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (12 \pm 15)\% \\
A_{CP}(K_2^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (3 \pm 6)\% \\
A_{CP}(K_2^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (-10 \pm 32)\% \\
A_{CP}(K^- \pi^+ \pi^+ \pi^-) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+ \pi^- \pi^+ \pi^-) &= (-2 \pm 4)\% \\
A_{CP}(K^+ K^- \pi^+ \pi^-) &= (1.3 \pm 1.7)\% \\
A_{CP}(K_1^*(1270)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-2.3 \pm 1.7)\% \\
A_{CP}(K_1^*(1270)^+ K^- \rightarrow K^{*0} \pi^+ K^-) &= (-1 \pm 10)\% \\
A_{CP}(K_1^*(1270)^- K^+ \rightarrow \bar{K}^{*0} \pi^- K^+) &= (-10 \pm 32)\% \\
A_{CP}(K_1^*(1270)^- K^+ \rightarrow K^+ K^- \pi^+ \pi^-) &= (1.7 \pm 3.5)\% \\
A_{CP}(K_1^*(1270)^+ K^- \rightarrow \rho^0 K^+ K^-) &= (-7 \pm 17)\% \\
A_{CP}(K_1^*(1270)^- K^+ \rightarrow \rho^0 K^- K^+) &= (10 \pm 13)\% \\
A_{CP}(K_1(1400)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-4.4 \pm 2.1)\% \\
A_{CP}(K^*(1410)^+ K^- \rightarrow K^{*0} \pi^+ K^-) &= (-20 \pm 17)\% \\
A_{CP}(K^*(1410)^- K^+ \rightarrow \bar{K}^{*0} \pi^- K^+) &= (-1 \pm 14)\% \\
A_{CP}(K^*(1680)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-17 \pm 29)\% \\
A_{CP}(K^{*0} \bar{K}^{*0} \text{ in } D^0, \bar{D}^0 \rightarrow K^{*0} \bar{K}^{*0}) &= (-5 \pm 14)\% \\
A_{CP}(K^{*0} \bar{K}^{*0} \text{ S-wave}) &= (-3.9 \pm 2.2)\% \\
A_{CP}(\phi \rho^0 \text{ in } D^0, \bar{D}^0 \rightarrow \phi \rho^0) &= (1 \pm 9)\%
\end{aligned}$$

$$\begin{aligned}
 A_{CP}(\phi \rho^0 \text{ } S\text{-wave}) &= (-3 \pm 5)\% \\
 A_{CP}(\phi \rho^0 \text{ } D\text{-wave}) &= (-37 \pm 19)\% \\
 A_{CP}(\phi(\pi^+ \pi^-)_{S\text{-wave}}) &= (6 \pm 6)\% \\
 A_{CP}(K^*(892)^0(K^- \pi^+)_{S\text{-wave}}) &= (-10 \pm 40)\% \\
 A_{CP}(K^+ K^- \pi^+ \pi^- \text{ non-resonant}) &= (8 \pm 20)\% \\
 A_{CP}((K^- \pi^+)_{P\text{-wave}}(K^+ \pi^-)_{S\text{-wave}}) &= (3 \pm 11)\% \\
 A_{CP}(K^+ K^- \mu^+ \mu^-) \text{ in } D^0, \bar{D}^0 \rightarrow K^+ K^- \mu^+ \mu^- &= (0 \pm 11)\% \\
 A_{CP}(\pi^+ \pi^- \mu^+ \mu^-) \text{ in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \mu^+ \mu^- &= (5 \pm 4)\%
 \end{aligned}$$

CP-even fractions (labeled by the D^0 decay)

$$\begin{aligned}
 \text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^0 \text{ decays} &= (97.3 \pm 1.7)\% \\
 \text{CP-even fraction in } D^0 \rightarrow K^+ K^- \pi^0 \text{ decays} &= (73 \pm 6)\% \\
 \text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- \text{ decays} &= (76.9 \pm 2.3)\% \\
 \text{CP-even fraction in } D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0 \text{ decays} &= (23.8 \pm 1.7)\% \\
 \text{CP-even fraction in } D^0 \rightarrow K^+ K^- \pi^+ \pi^- \text{ decays} &= (75 \pm 4)\%
 \end{aligned}$$

CP-violation asymmetry difference

$$\Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-0.154 \pm 0.029)\%$$

χ^2 tests of CP-violation (CPV) p-values

$$\begin{aligned}
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \pi^0 &= 4.9\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- &= (0.6 \pm 0.2)\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^- &= 96\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^0 &= 16.6\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^+ \pi^- &= 9.1\%
 \end{aligned}$$

T-violation decay-rate asymmetry

$$\begin{aligned}
 A_T(K^+ K^- \pi^+ \pi^-) &= (2.9 \pm 2.2) \times 10^{-3} [b] \\
 A_{T\text{viol}}(K_S \pi^+ \pi^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow K_S \pi^+ \pi^- \pi^0 &= (-0.3^{+1.4}_{-1.6}) \times 10^{-3}
 \end{aligned}$$

CPT-violation decay-rate asymmetry

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$\begin{aligned}
 r_V &\equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 1.46 \pm 0.07 \\
 r_2 &\equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 0.68 \pm 0.06 \\
 f_+(0) &\text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.736 \pm 0.004 \\
 f_+(0) |V_{cs}| &\text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.7166 \pm 0.0030 \\
 r_1 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = -2.40 \pm 0.16 \\
 r_2 &\equiv a_2/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 5 \pm 4 \\
 f_+(0) &\text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 0.637 \pm 0.009 \\
 f_+(0) |V_{cd}| &\text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 0.1436 \pm 0.0026 \quad (S = 1.5) \\
 r_1 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = -1.97 \pm 0.28 \quad (S = 1.4) \\
 r_2 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = -0.2 \pm 2.2 \quad (S = 1.7)
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as $\bar{K}^0$ modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Topological modes			
0-prongs	[j] (15 $\pm$ 6) %		—
2-prongs	(71 $\pm$ 6) %		—
4-prongs	[k] (14.6 $\pm$ 0.5) %		—
6-prongs	[l] (6.5 $\pm$ 1.3) $\times 10^{-4}$		—
Inclusive modes			
e^+ anything	[n] (6.49 $\pm$ 0.11) %		—
μ^+ anything	(6.8 $\pm$ 0.6) %		—
K^- anything	(54.7 $\pm$ 2.8) %	S=1.3	—
$\bar{K}^0$ anything + K^0 anything	(47 $\pm$ 4) %		—
K^+ anything	(3.4 $\pm$ 0.4) %		—
$K^*(892)^-$ anything	(15 $\pm$ 9) %		—
$\bar{K}^*(892)^0$ anything	(9 $\pm$ 4) %		—
$K^*(892)^+$ anything	< 3.6 %	CL=90%	—
$K^*(892)^0$ anything	(2.8 $\pm$ 1.3) %		—
η anything	(9.5 $\pm$ 0.9) %		—
η' anything	(2.48 $\pm$ 0.27) %		—
ϕ anything	(1.08 $\pm$ 0.04) %		—
invisibles	< 9.4 $\times 10^{-5}$	CL=90%	—
Semileptonic modes			
$K^- e^+ \nu_e$	(3.541 $\pm$ 0.034) %	S=1.3	867
$K^- \mu^+ \nu_\mu$	(3.41 $\pm$ 0.04) %		864
$K^*(892)^- e^+ \nu_e$	(2.15 $\pm$ 0.16) %		719
$K^*(892)^- \mu^+ \nu_\mu$	(1.89 $\pm$ 0.24) %		714
$K^- \pi^0 e^+ \nu_e$	(1.6 $\pm$ 1.3 / $\pm$ 0.5) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$	(1.44 $\pm$ 0.04) %		860
$(\bar{K}^0 \pi^-)_{S\text{-wave}} e^+ \nu_e$	(7.9 $\pm$ 1.7) $\times 10^{-4}$		860
$K^- \pi^+ \pi^- e^+ \nu_e$	(2.8 $\pm$ 1.4 / $\pm$ 1.1) $\times 10^{-4}$		843
$K_1(1270)^- e^+ \nu_e$	(7.6 $\pm$ 4.0 / $\pm$ 3.1) $\times 10^{-4}$		511
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.3 $\times 10^{-3}$	CL=90%	821
$(\bar{K}^*(892) \pi)^- \mu^+ \nu_\mu$	< 1.5 $\times 10^{-3}$	CL=90%	692
$\pi^- e^+ \nu_e$	(2.91 $\pm$ 0.04) $\times 10^{-3}$		927

$\pi^- \mu^+ \nu_\mu$	$(2.67 \pm 0.12) \times 10^{-3}$	S=1.3	924
$\pi^- \pi^0 e^+ \nu_e$	$(1.45 \pm 0.07) \times 10^{-3}$		922
$\rho^- e^+ \nu_e$	$(1.50 \pm 0.12) \times 10^{-3}$	S=1.9	771
$a(980)^- e^+ \nu_e, a^- \rightarrow \eta \pi^-$	$(1.33 \pm_{-0.30}^{+0.34}) \times 10^{-4}$		—
$b_1(1235)^- e^+ \nu_e, b_1^- \rightarrow \omega \pi^-$	$< 1.12 \times 10^{-4}$	CL=90%	—

Hadronic modes with one $\bar{K}$

$K^- \pi^+$	$(3.946 \pm 0.030) \%$	S=1.2	861
$K_S^0 \pi^0$	$(1.239 \pm 0.022) \%$		860
$K_L^0 \pi^0$	$(10.0 \pm 0.7) \times 10^{-3}$		860
$K_S^0 \pi^+ \pi^-$	[c] $(2.80 \pm 0.18) \%$	S=1.1	842
$K_S^0 \rho^0$	$(6.3 \pm_{-0.8}^{+0.6}) \times 10^{-3}$		674
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	$(2.0 \pm 0.6) \times 10^{-4}$		670
$K_S^0 (\pi^+ \pi^-)_{S\text{-wave}}$	$(3.3 \pm 0.8) \times 10^{-3}$		842
$K_S^0 f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$(1.20 \pm_{-0.23}^{+0.40}) \times 10^{-3}$		549
$K_S^0 f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	$(2.8 \pm_{-1.3}^{+0.9}) \times 10^{-3}$		†
$K_S^0 f_2(1270), f_2 \rightarrow \pi^+ \pi^-$	$(9 \pm_{-6}^{+10}) \times 10^{-5}$		262
$K^*(892)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(1.64 \pm_{-0.17}^{+0.14}) \%$		711
$K_0^*(1430)^- \pi^+, K_0^{*-} \rightarrow K_S^0 \pi^-$	$(2.67 \pm_{-0.33}^{+0.40}) \times 10^{-3}$		378
$K_2^*(1430)^- \pi^+, K_2^{*-} \rightarrow K_S^0 \pi^-$	$(3.4 \pm_{-1.0}^{+1.9}) \times 10^{-4}$		367
$K^*(1680)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(4.4 \pm 3.5) \times 10^{-4}$		46
$K^*(892)^+ \pi^-, K^{*+} \rightarrow K_S^0 \pi^+$	[o] $(1.13 \pm_{-0.34}^{+0.60}) \times 10^{-4}$		711
$K_0^*(1430)^+ \pi^-, K_0^{*+} \rightarrow K_S^0 \pi^+$	[o] $< 1.4 \times 10^{-5}$	CL=95%	—
$K_2^*(1430)^+ \pi^-, K_2^{*+} \rightarrow K_S^0 \pi^+$	[o] $< 3.4 \times 10^{-5}$	CL=95%	—
$K_S^0 \pi^+ \pi^-$ nonresonant	$(2.5 \pm_{-1.6}^{+6.0}) \times 10^{-4}$		842
$K^- \pi^+ \pi^0$	[c] $(14.4 \pm 0.5) \%$	S=2.0	844
$K^- \rho^+$	$(11.2 \pm 0.7) \%$		675
$K^- \rho(1700)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(8.2 \pm 1.8) \times 10^{-3}$		†
$K^*(892)^- \pi^+, K^*(892)^- \rightarrow K^- \pi^0$	$(2.31 \pm_{-0.20}^{+0.40}) \%$		711

$\bar{K}^*(892)^0 \pi^0, \bar{K}^*(892)^0 \rightarrow$ $K^- \pi^+$	$(1.95 \pm 0.24) \%$		711
$K_0^*(1430)^- \pi^+, K_0^{*-} \rightarrow$ $K^- \pi^0$	$(4.8 \pm 2.2) \times 10^{-3}$		378
$\bar{K}_0^*(1430)^0 \pi^0, \bar{K}_0^{*0} \rightarrow$ $K^- \pi^+$	$(5.9 \pm 5.0 \pm 1.6) \times 10^{-3}$		379
$K^*(1680)^- \pi^+, K^{*-} \rightarrow$ $K^- \pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$		46
$K^- \pi^+ \pi^0$ nonresonant	$(1.15 \pm 0.60 \pm 0.20) \%$		844
$K_S^0 2\pi^0$	$(9.1 \pm 1.1) \times 10^{-3}$	S=2.2	843
$K_S^0(2\pi^0)_{S-wave}$	$(2.6 \pm 0.7) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \pi^0, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(8.1 \pm 0.7) \times 10^{-3}$		711
$\bar{K}^*(1430)^0 \pi^0, \bar{K}^{*0} \rightarrow$ $K_S^0 \pi^0$	$(4 \pm 23) \times 10^{-5}$		—
$\bar{K}^*(1680)^0 \pi^0, \bar{K}^{*0} \rightarrow$ $K_S^0 \pi^0$	$(1.0 \pm 0.4) \times 10^{-3}$		—
$K_S^0 f_2(1270), f_2 \rightarrow 2\pi^0$	$(2.3 \pm 1.1) \times 10^{-4}$		—
$2K_S^0, \text{one } K_S^0 \rightarrow 2\pi^0$	$(3.2 \pm 1.1) \times 10^{-4}$		—
$K^- 2\pi^+ \pi^-$	[c] $(8.22 \pm 0.14) \%$	S=1.1	813
$K^- \pi^+ \rho^0$ total	$(6.86 \pm 0.31) \%$		609
$K^- \pi^+ \rho^0$ 3-body	$(6.1 \pm 1.6) \times 10^{-3}$		609
$\bar{K}^*(892)^0 \rho^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(1.01 \pm 0.05) \%$		416
$\bar{K}^*(892)^0 \rho^0$ transverse, $\bar{K}^{*0} \rightarrow K^- \pi^+$	$(1.2 \pm 0.4) \%$		417
$K^- a_1(1260)^+, a_1^+ \rightarrow$ $\rho^0 \pi^+$	$(4.32 \pm 0.32) \%$		327
$K_1(1270)^- \pi^+, K_1^- \rightarrow$ $K^- \pi^+ \pi^-$ total	$(3.9 \pm 0.4) \times 10^{-3}$		—
$K_1(1270)^- \pi^+, K_1^- \rightarrow$ $\bar{K}^*(892)^0 \pi^-, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(6.6 \pm 2.3) \times 10^{-4}$		484
$K^- 2\pi^+ \pi^-$ nonresonant	$(1.81 \pm 0.07) \%$		813
$K_S^0 \pi^+ \pi^- \pi^0$	[p] $(5.2 \pm 0.6) \%$		813
$K_S^0 \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$	$(1.17 \pm 0.03) \times 10^{-3}$		772
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	$(9.9 \pm 0.6) \times 10^{-3}$		670
$K^- \pi^+ 2\pi^0$	$(8.86 \pm 0.23) \%$		815
$K^- 2\pi^+ \pi^- \pi^0$	$(4.3 \pm 0.4) \%$		771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(1.3 \pm 0.6) \%$		643
$K^- \pi^+ \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	$(2.8 \pm 0.5) \%$		605
$\bar{K}^*(892)^0 \omega, \bar{K}^{*0} \rightarrow$ $K^- \pi^+, \omega \rightarrow$ $\pi^+ \pi^- \pi^0$	$(6.5 \pm 3.0) \times 10^{-3}$		410

$K_S^0 \eta \pi^0$	$(1.01 \pm 0.05) \%$	721
$K_S^0 a_0(980), a_0 \rightarrow \eta \pi^0$	$(1.20 \pm 0.28) \%$	—
$\bar{K}^{*0}(892)^0 \eta, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(2.9 \pm 0.7) \times 10^{-3}$	—
$K^- \pi^+ \eta$	$(1.88 \pm 0.05) \%$	S=1.4 721
$K^{*0}(892)^0 \eta, K^{*0} \rightarrow K^- \pi^+$	$(8.9 \pm_{-0.6}^{+0.8}) \times 10^{-3}$	—
$a_0(980)^+ K^-, a_0^+ \rightarrow \eta \pi^+$	$(7.4 \pm_{-0.7}^{+0.9}) \times 10^{-3}$	—
$K_2^{*0}(1980)^- \pi^+, K_2^{*-} \rightarrow$ $K^- \eta$	$(2.2 \pm_{-1.9}^{+1.7}) \times 10^{-4}$	—
$K^- \pi^+ \pi^0 \eta$	$(4.49 \pm 0.27) \times 10^{-3}$	656
$K_S^0 \pi^+ \pi^- \eta$	$(2.80 \pm 0.21) \times 10^{-3}$	651
$K_S^0 2\pi^0 \eta$	$(1.76 \pm 0.26) \times 10^{-3}$	656
$K_S^0 2\pi^+ 2\pi^-$	$(2.66 \pm 0.30) \times 10^{-3}$	768
$K_S^0 \rho^0 \pi^+ \pi^-, \text{ no } K^{*0}(892)^-$	$(1.1 \pm 0.7) \times 10^{-3}$	—
$K^{*0}(892)^- 2\pi^+ \pi^-,$ $K^{*0}(892)^- \rightarrow K_S^0 \pi^-,$ $\text{no } \rho^0$	$(5 \pm 7) \times 10^{-4}$	642
$K^{*0}(892)^- \rho^0 \pi^+,$ $K^{*0}(892)^- \rightarrow K_S^0 \pi^-$	$(1.6 \pm 0.6) \times 10^{-3}$	230
$K_S^0 2\pi^+ 2\pi^- \text{ nonresonant}$	$< 1.2 \times 10^{-3}$	CL=90% 768
$K^- 3\pi^+ 2\pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	713

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. These nine modes below are all corrected for unseen decays of the resonances.

$K_S^0 \eta$	$(5.08 \pm 0.13) \times 10^{-3}$	772
$K_S^0 \omega$	$(1.11 \pm 0.06) \%$	670
$K_S^0 \eta'(958)$	$(9.49 \pm 0.32) \times 10^{-3}$	565
$\bar{K}^{*0}(892)^0 \pi^+ \pi^- \pi^0$	$(1.9 \pm 0.9) \%$	643
$\bar{K}^{*0}(892)^0 \eta$	$(1.41 \pm 0.12) \%$	583
$K^- \pi^+ \omega$	$(3.1 \pm 0.6) \%$	605
$\bar{K}^{*0}(892)^0 \omega$	$(1.1 \pm 0.5) \%$	410
$K^- \pi^+ \eta'(958)$	$(6.43 \pm 0.34) \times 10^{-3}$	479
$K_S^0 \eta'(958) \pi^0$	$(2.52 \pm 0.27) \times 10^{-3}$	479
$\bar{K}^{*0}(892)^0 \eta'(958)$	$< 1.0 \times 10^{-3}$	CL=90% 119

Hadronic modes with three K's

$K_S^0 K^+ K^-$	$(4.42 \pm 0.32) \times 10^{-3}$	544
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	$(2.9 \pm 0.4) \times 10^{-3}$	—
$K^- a_0(980)^+, a_0^+ \rightarrow$ $K^+ K_S^0$	$(5.9 \pm 1.8) \times 10^{-4}$	—
$K^+ a_0(980)^-, a_0^- \rightarrow$ $K^- K_S^0$	$< 1.1 \times 10^{-4}$	CL=95% —

$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	$< 9 \times 10^{-5}$	CL=95%	—
$K_S^0 \phi, \phi \rightarrow K^+ K^-$	$(2.03 \pm 0.15) \times 10^{-3}$		520
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	$(1.7 \pm 1.1) \times 10^{-4}$		—
$3K_S^0$	$(7.5 \pm 0.7) \times 10^{-4}$	S=1.4	539
$K^+ 2K^- \pi^+$	$(2.25 \pm 0.32) \times 10^{-4}$		434
$K^+ K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(4.5 \pm 1.8) \times 10^{-5}$		†
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.7) \times 10^{-5}$		422
$\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(1.08 \pm 0.20) \times 10^{-4}$		†
$K^+ 2K^- \pi^+$ nonresonant	$(3.4 \pm 1.5) \times 10^{-5}$		434
$2K_S^0 K^\pm \pi^\mp$	$(5.9 \pm 1.3) \times 10^{-4}$		427

Pionic modes

$\pi^+ \pi^-$	$(1.453 \pm 0.024) \times 10^{-3}$	S=1.4	922
$2\pi^0$	$(8.26 \pm 0.25) \times 10^{-4}$		923
$\pi^+ \pi^- \pi^0$	$(1.49 \pm 0.06) \%$	S=2.1	907
$\rho^+ \pi^-$	$(1.01 \pm 0.04) \%$		764
$\rho^0 \pi^0$	$(3.86 \pm 0.23) \times 10^{-3}$		764
$\rho^- \pi^+$	$(5.15 \pm 0.25) \times 10^{-3}$		764
$\rho(1450)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.6 \pm 2.1) \times 10^{-5}$		—
$\rho(1450)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	$(4.5 \pm 1.9) \times 10^{-5}$		—
$\rho(1450)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	$(2.7 \pm 0.4) \times 10^{-4}$		—
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	$(6.1 \pm 1.5) \times 10^{-4}$		—
$\rho(1700)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	$(7.4 \pm 1.8) \times 10^{-4}$		—
$\rho(1700)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	$(4.8 \pm 1.1) \times 10^{-4}$		—
$f_0(980) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(3.7 \pm 0.9) \times 10^{-5}$		—
$f_0(500) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(1.22 \pm 0.22) \times 10^{-4}$		—
$f_0(1370) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(5.5 \pm 2.1) \times 10^{-5}$		—
$f_0(1500) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(5.8 \pm 1.6) \times 10^{-5}$		—
$f_0(1710) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(4.6 \pm 1.6) \times 10^{-5}$		—
$f_2(1270) \pi^0, f_2 \rightarrow \pi^+ \pi^-$	$(1.96 \pm 0.21) \times 10^{-4}$		—
$\pi^+ \pi^- \pi^0$ nonresonant	$(1.3 \pm 0.4) \times 10^{-4}$		907
$3\pi^0$	$(2.0 \pm 0.5) \times 10^{-4}$		908
$2\pi^+ 2\pi^-$	$(7.55 \pm 0.20) \times 10^{-3}$		880
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow 2\pi^+ \pi^-$ total	$(4.53 \pm 0.31) \times 10^{-3}$		—
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow \rho^0 \pi^+$ S-wave	$(3.13 \pm 0.21) \times 10^{-3}$		—
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow \rho^0 \pi^+$ D-wave	$(1.9 \pm 0.5) \times 10^{-4}$		—
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow \sigma \pi^+$	$(6.4 \pm 0.7) \times 10^{-4}$		—
$a_1(1260)^- \pi^+, a_1^- \rightarrow \rho^0 \pi^-$ S-wave	$(2.3 \pm 0.9) \times 10^{-4}$		—

$a_1(1260)^- \pi^+, a_1^- \rightarrow \sigma \pi^-$	$(6.0 \pm 3.4) \times 10^{-5}$	—
$\pi(1300)^+ \pi^-, \pi(1300)^+ \rightarrow \sigma \pi^+$	$(5.1 \pm 2.7) \times 10^{-4}$	—
$\pi(1300)^- \pi^+, \pi(1300)^- \rightarrow \sigma \pi^-$	$(2.3 \pm 2.2) \times 10^{-4}$	—
$a_1(1640)^+ \pi^-, a_1^+ \rightarrow \rho^0 \pi^+ D\text{-wave}$	$(3.2 \pm 1.6) \times 10^{-4}$	—
$a_1(1640)^+ \pi^-, a_1^+ \rightarrow \sigma \pi^+$	$(1.8 \pm 1.4) \times 10^{-4}$	—
$\pi_2(1670)^+ \pi^-, \pi_2^+ \rightarrow \pi^+ \pi^-$	$(2.0 \pm 0.9) \times 10^{-4}$	—
$f_2(1270)^0 \pi^+, f_2^0 \rightarrow \pi^+ \pi^-$		
$\pi_2(1670)^+ \pi^-, \pi_2^+ \rightarrow \sigma \pi^+$	$(2.6 \pm 1.0) \times 10^{-4}$	—
$2\rho^0$ total	$(1.85 \pm 0.13) \times 10^{-3}$	518
$2\rho^0$, parallel helicities	$(8.3 \pm 3.2) \times 10^{-5}$	—
$2\rho^0$, perpendicular helicities	$(4.8 \pm 0.6) \times 10^{-4}$	—
$2\rho^0$, longitudinal helicities	$(1.27 \pm 0.10) \times 10^{-3}$	—
$2\rho(770)^0$, <i>S</i> -wave	$(1.8 \pm 1.3) \times 10^{-4}$	—
$2\rho(770)^0$, <i>P</i> -wave	$(5.3 \pm 1.3) \times 10^{-4}$	—
$2\rho(770)^0$, <i>D</i> -wave	$(6.2 \pm 3.0) \times 10^{-4}$	—
Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$	$(1.51 \pm 0.12) \times 10^{-3}$	—
3-body total		
$\sigma \pi^+ \pi^-$	$(6.2 \pm 0.9) \times 10^{-4}$	—
$\sigma \rho(770)^0$	$(5.0 \pm 2.5) \times 10^{-4}$	—
$f_0(980) \pi^+ \pi^-, f_0 \rightarrow \pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-4}$	—
$f_2(1270) \pi^+ \pi^-, f_2 \rightarrow \pi^+ \pi^-$	$(3.7 \pm 0.6) \times 10^{-4}$	—
$2f_2(1270), f_2 \rightarrow \pi^+ \pi^-$	$(1.6 \pm 1.8) \times 10^{-4}$	—
$f_0(1370) \sigma, f_0 \rightarrow \pi^+ \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	—
$\pi^+ \pi^- 2\pi^0$	$(1.02 \pm 0.09) \%$	882
$\eta \pi^0$	[<i>q</i>] $(6.3 \pm 0.6) \times 10^{-4}$	S=1.1 846
$\omega \pi^0$	[<i>q</i>] $(1.17 \pm 0.35) \times 10^{-4}$	761
$\omega \eta$	$(1.98 \pm 0.18) \times 10^{-3}$	S=1.1 648
$2\pi^+ 2\pi^- \pi^0$	$(4.2 \pm 0.5) \times 10^{-3}$	844
$\eta \pi^+ \pi^-$	[<i>q</i>] $(1.16 \pm 0.07) \times 10^{-3}$	827
$\omega \pi^+ \pi^-$	[<i>q</i>] $(1.33 \pm 0.20) \times 10^{-3}$	738
$\omega \pi^0 \pi^0$	$< 1.10 \times 10^{-3}$	CL=90% 740
$\eta 2\pi^0$	$(3.8 \pm 1.3) \times 10^{-4}$	829
$\pi^+ \pi^- \pi^0 \eta$	$(3.23 \pm 0.22) \times 10^{-3}$	797
$3\pi^+ 3\pi^-$	$(4.3 \pm 1.2) \times 10^{-4}$	795
$\eta'(958) \pi^0$	$(9.2 \pm 1.0) \times 10^{-4}$	678
$\eta'(958) \pi^+ \pi^-$	$(4.5 \pm 1.7) \times 10^{-4}$	650
2η	$(2.11 \pm 0.19) \times 10^{-3}$	S=2.3 754

$2\eta\pi^0$	$(7.3 \pm 2.2) \times 10^{-4}$		699
3η	$< 1.3 \times 10^{-4}$	CL=90%	421
$\eta\eta'(958)$	$(1.01 \pm 0.19) \times 10^{-3}$		537

Hadronic modes with a $K\bar{K}$ pair

$K^+ K^-$	$(4.08 \pm 0.06) \times 10^{-3}$	S=1.6	791
$2K_S^0$	$(1.41 \pm 0.05) \times 10^{-4}$	S=1.1	789
$K_S^0 K^- \pi^+$	$(3.3 \pm 0.5) \times 10^{-3}$	S=1.1	739
$\bar{K}^*(892)^0 K_S^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(8.2 \pm 1.6) \times 10^{-5}$		608
$K^*(892)^+ K^-, K^{*+} \rightarrow$ $K_S^0 \pi^+$	$(1.89 \pm 0.30) \times 10^{-3}$		—
$\bar{K}^*(1410)^0 K_S^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(1.3 \pm 1.9) \times 10^{-4}$		—
$K^*(1410)^+ K^-, K^{*+} \rightarrow$ $K_S^0 \pi^+$	$(3.2 \pm 1.9) \times 10^{-4}$		—
$(K^- \pi^+)_{S\text{-wave}} K_S^0$	$(6.0 \pm 2.9) \times 10^{-4}$		739
$(K_S^0 \pi^+)_{S\text{-wave}} K^-$	$(3.9 \pm 1.0) \times 10^{-4}$		739
$a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-$	$(1.3 \pm 1.4) \times 10^{-4}$		—
$a_0(1450)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	$(2.5 \pm 2.0) \times 10^{-5}$		—
$a_2(1320)^- \pi^+, a_2^- \rightarrow$ $K_S^0 K^-$	$(5 \pm 5) \times 10^{-6}$		—
$\rho(1450)^- \pi^+, \rho^- \rightarrow K_S^0 K^-$	$(4.6 \pm 2.5) \times 10^{-5}$		—
$K_S^0 K^+ \pi^-$	$(2.17 \pm 0.34) \times 10^{-3}$	S=1.1	739
$K^*(892)^0 K_S^0, K^{*0} \rightarrow$ $K^+ \pi^-$	$(1.12 \pm 0.21) \times 10^{-4}$		608
$K^*(892)^- K^+, K^{*-} \rightarrow$ $K_S^0 \pi^-$	$(6.2 \pm 1.0) \times 10^{-4}$		—
$K^*(1410)^0 K_S^0, K^{*0} \rightarrow$ $K^+ \pi^+$	$(5 \pm 8) \times 10^{-5}$		—
$K^*(1410)^- K^+, K^{*-} \rightarrow$ $K_S^0 \pi^-$	$(2.6 \pm 2.0) \times 10^{-4}$		—
$(K^+ \pi^-)_{S\text{-wave}} K_S^0$	$(3.7 \pm 1.9) \times 10^{-4}$		739
$(K_S^0 \pi^-)_{S\text{-wave}} K^+$	$(1.4 \pm 0.6) \times 10^{-4}$		739
$a_0(980)^+ \pi^-, a_0^+ \rightarrow K_S^0 K^+$	$(6 \pm 4) \times 10^{-4}$		—
$a_0(1450)^+ \pi^-, a_0^+ \rightarrow$ $K_S^0 K^+$	$(3.2 \pm 2.5) \times 10^{-5}$		—
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow K_S^0 K^+$	$(1.1 \pm 0.6) \times 10^{-5}$		—
$K^+ K^- \pi^0$	$(3.42 \pm 0.14) \times 10^{-3}$		743
$K^*(892)^+ K^-, K^*(892)^+ \rightarrow$ $K^+ \pi^0$	$(1.52 \pm 0.07) \times 10^{-3}$		—

$K^*(892)^- K^+, K^*(892)^- \rightarrow K^- \pi^0$	$(5.4 \pm 0.4) \times 10^{-4}$	—
$(K^+ \pi^0)_{S\text{-wave}} K^-$	$(2.43 \pm 0.18) \times 10^{-3}$	743
$(K^- \pi^0)_{S\text{-wave}} K^+$	$(1.3 \pm 0.5) \times 10^{-4}$	743
$f_0(980) \pi^0, f_0 \rightarrow K^+ K^-$	$(3.6 \pm 0.6) \times 10^{-4}$	—
$\phi \pi^0, \phi \rightarrow K^+ K^-$	$(6.6 \pm 0.4) \times 10^{-4}$	—
$2K_S^0 \pi^0$	$< 5.9 \times 10^{-4}$	740
$K^+ K^- \eta$	$(5.9 \pm 1.9) \times 10^{-5}$	514
$2K_S^0 \eta$	$(1.3 \pm 0.6) \times 10^{-4}$	508
$K^+ K^- \pi^0 \pi^0$	$(6.9 \pm 0.8) \times 10^{-4}$	681
$K^+ K^- \pi^+ \pi^-$	$(2.47 \pm 0.11) \times 10^{-3}$	677
$\phi(\pi^+ \pi^-)_{S\text{-wave}}, \phi \rightarrow K^+ K^-$	$(10 \pm 5) \times 10^{-5}$	614
$(\phi \rho^0)_{S\text{-wave}}, \phi \rightarrow K^+ K^-$	$(6.9 \pm 0.6) \times 10^{-4}$	250
$(\phi \rho^0)_{P\text{-wave}}, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.9) \times 10^{-5}$	—
$(\phi \rho^0)_{D\text{-wave}}, \phi \rightarrow K^+ K^-$	$(4.2 \pm 1.4) \times 10^{-5}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{S\text{-wave}}, K^{*0} \rightarrow K^\pm \pi^\mp$	$(2.24 \pm 0.13) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{P\text{-wave}}, K^* \rightarrow K^\pm \pi^\mp$	$(1.20 \pm 0.08) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{D\text{-wave}}, K^* \rightarrow K^\pm \pi^\mp$	$(4.7 \pm 0.4) \times 10^{-5}$	—
$K^*(892)^0 (K^- \pi^+)_{S\text{-wave}}, 3\text{-body}, K^{*0} \rightarrow K^+ \pi^-$	$(1.4 \pm 0.6) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow K^{*0} \pi^+$	$(1.4 \pm 0.9) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(1.5 \pm 0.5) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow \rho^0 K^+$	$(2.2 \pm 0.6) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow \omega(782) K^+, \omega \rightarrow \pi^+ \pi^-$	$(1.5 \pm 1.2) \times 10^{-5}$	—
$K_1(1270)^- K^+, K_1^- \rightarrow \rho^0 K^-$	$(1.3 \pm 0.4) \times 10^{-4}$	—
$K_1(1400)^+ K^-, K_1^+ \rightarrow K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(4.6 \pm 0.4) \times 10^{-4}$	—
$K^*(1410)^- K^+, K^{*-} \rightarrow \bar{K}^{*0} \pi^-$	$(7.0 \pm 1.1) \times 10^{-5}$	—
$K_1(1680)^+ K^-, K_1^+ \rightarrow K^{*0} \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(8.9 \pm 3.2) \times 10^{-5}$	—
$K^+ K^- \pi^+ \pi^-$ non-resonant	$(2.7 \pm 0.6) \times 10^{-4}$	—
$2K_S^0 \pi^+ \pi^-$	$(5.3 \pm 0.9) \times 10^{-4}$	673

$K_S^0 K^- \pi^+ \pi^0$	$(1.32 \pm 0.16) \times 10^{-3}$		677
$K_S^0 K^+ \pi^- \pi^0$	$(6.5 \pm 0.7) \times 10^{-4}$		677
$K_S^0 K^- 2\pi^+ \pi^-$	$< 1.4 \times 10^{-4}$	CL=90%	595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$		600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi \pi^0$	$(1.17 \pm 0.04) \times 10^{-3}$		645
$\phi \eta$	$(1.8 \pm 0.5) \times 10^{-4}$		489
$\phi \omega$	$< 2.1 \times 10^{-3}$	CL=90%	238

Radiative modes

$\rho^0 \gamma$	$(1.82 \pm 0.32) \times 10^{-5}$		771
$\omega \gamma$	$< 2.4 \times 10^{-4}$	CL=90%	768
$\phi \gamma$	$(2.81 \pm 0.19) \times 10^{-5}$		654
$\bar{K}^*(892)^0 \gamma$	$(4.1 \pm 0.7) \times 10^{-4}$		719

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

$K^+ \ell^- \bar{\nu}_\ell$ via $\bar{D}^0$	$< 2.2 \times 10^{-5}$	CL=90%	—
K^+ or $K^*(892)^+$ $e^- \bar{\nu}_e$ via $\bar{D}^0$	$< 6 \times 10^{-5}$	CL=90%	—
$K^+ \pi^-$ DC	$(1.50 \pm 0.07) \times 10^{-4}$	S=3.0	861
$K^+ \pi^-$ via DCS	$(1.363 \pm 0.025) \times 10^{-4}$		—
$K^+ \pi^-$ via $\bar{D}^0$	$< 1.6 \times 10^{-5}$	CL=95%	861
$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \bar{D}^0$	$< 1.8 \times 10^{-4}$	CL=95%	—
$K^*(892)^+ \pi^-$, $K^{*+} \rightarrow K_S^0 \pi^+$ DC	$(1.13 \pm_{-0.34}^{+0.60}) \times 10^{-4}$		711
$K_0^*(1430)^+ \pi^-$, $K_0^{*+} \rightarrow K_S^0 \pi^+$ DC	$< 1.4 \times 10^{-5}$		—
$K_2^*(1430)^+ \pi^-$, $K_2^{*+} \rightarrow K_S^0 \pi^+$ DC	$< 3.4 \times 10^{-5}$		—
$K^+ \pi^- \pi^0$ DC	$(3.05 \pm 0.15) \times 10^{-4}$		844
$K^+ \pi^- \pi^0$ via $\bar{D}^0$	$(7.6 \pm_{-0.6}^{+0.5}) \times 10^{-4}$		—
$K^+ \pi^+ 2\pi^-$ via DCS	$(2.49 \pm 0.07) \times 10^{-4}$		—
$K^+ \pi^+ 2\pi^-$ DC	$(2.65 \pm 0.06) \times 10^{-4}$		813
$K^+ \pi^+ 2\pi^-$ via $\bar{D}^0$	$(7.9 \pm 3.0) \times 10^{-6}$		812
μ^- anything via $\bar{D}^0$	$< 4 \times 10^{-4}$	CL=90%	—

$\Delta C = 1$ weak neutral current (C1) modes, Lepton Family number (LF) violating modes, Lepton (L) or Baryon (B) number violating modes

$\gamma \gamma$	C1	$< 8.5 \times 10^{-7}$	CL=90%	932
$e^+ e^-$	C1	$< 7.9 \times 10^{-8}$	CL=90%	932
$\mu^+ \mu^-$	C1	$< 6.2 \times 10^{-9}$	CL=90%	926

$\pi^0 e^+ e^-$	$C1$	< 4	$\times 10^{-6}$	CL=90%	928
$\pi^0 \mu^+ \mu^-$	$C1$	< 1.8	$\times 10^{-4}$	CL=90%	915
$\eta e^+ e^-$	$C1$	< 3	$\times 10^{-6}$	CL=90%	852
$\eta \mu^+ \mu^-$	$C1$	< 5.3	$\times 10^{-4}$	CL=90%	838
$\pi^+ \pi^- e^+ e^-$	$C1$	< 7	$\times 10^{-6}$	CL=90%	922
$\rho^0 e^+ e^-$	$C1$	< 1.0	$\times 10^{-4}$	CL=90%	771
$\pi^+ \pi^- \mu^+ \mu^-$	$C1$	(9.6 ± 1.2)	$\times 10^{-7}$		894
$\pi^+ \pi^- \mu^+ \mu^-$ (non-res)		< 5.5	$\times 10^{-7}$	CL=90%	—
$\rho^0 \mu^+ \mu^-$	$C1$	< 2.2	$\times 10^{-5}$	CL=90%	754
$\omega e^+ e^-$	$C1$	< 6	$\times 10^{-6}$	CL=90%	768
$\omega \mu^+ \mu^-$	$C1$	< 8.3	$\times 10^{-4}$	CL=90%	751
$K^- K^+ e^+ e^-$	$C1$	< 1.1	$\times 10^{-5}$	CL=90%	791
$\phi e^+ e^-$	$C1$	< 5.2	$\times 10^{-5}$	CL=90%	654
$K^- K^+ \mu^+ \mu^-$	$C1$	(1.54 ± 0.32)	$\times 10^{-7}$		710
$K^- K^+ \mu^+ \mu^-$ (non-res)		< 3.3	$\times 10^{-5}$	CL=90%	—
$\phi \mu^+ \mu^-$	$C1$	< 3.1	$\times 10^{-5}$	CL=90%	631
$\overline{K}^0 e^+ e^-$		$[h] < 2.4$	$\times 10^{-5}$	CL=90%	866
$\overline{K}^0 \mu^+ \mu^-$		$[h] < 2.6$	$\times 10^{-4}$	CL=90%	852
$K^- \pi^+ e^+ e^-$, $675 < m_{ee} < 875$ MeV		(4.0 ± 0.5)	$\times 10^{-6}$		—
$K^- \pi^+ e^+ e^-$, $1.005 < m_{ee} < 1.035$ GeV		< 5	$\times 10^{-7}$	CL=90%	—
$\overline{K}^*(892)^0 e^+ e^-$		$[h] < 4.7$	$\times 10^{-5}$	CL=90%	719
$K^- \pi^+ \mu^+ \mu^-$	$C1$	< 3.59	$\times 10^{-4}$	CL=90%	829
$K^- \pi^+ \mu^+ \mu^-$, $675 < m_{\mu\mu} < 875$ MeV		(4.2 ± 0.4)	$\times 10^{-6}$		—
$\overline{K}^*(892)^0 \mu^+ \mu^-$		$[h] < 2.4$	$\times 10^{-5}$	CL=90%	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	$C1$	< 8.1	$\times 10^{-4}$	CL=90%	863
$\mu^\pm e^\mp$	LF	$[r] < 1.3$	$\times 10^{-8}$	CL=90%	929
$\pi^0 e^\pm \mu^\mp$	LF	$[r] < 8.0$	$\times 10^{-7}$	CL=90%	924
$\eta e^\pm \mu^\mp$	LF	$[r] < 2.25$	$\times 10^{-6}$	CL=90%	848
$\pi^+ \pi^- e^\pm \mu^\mp$	LF	$[r] < 1.71$	$\times 10^{-6}$	CL=90%	911
$\rho^0 e^\pm \mu^\mp$	LF	$[r] < 5.0$	$\times 10^{-7}$	CL=90%	767
$\omega e^\pm \mu^\mp$	LF	$[r] < 1.71$	$\times 10^{-6}$	CL=90%	764
$K^- K^+ e^\pm \mu^\mp$	LF	$[r] < 1.00$	$\times 10^{-6}$	CL=90%	754
$\phi e^\pm \mu^\mp$	LF	$[r] < 5.1$	$\times 10^{-7}$	CL=90%	648
$\overline{K}^0 e^\pm \mu^\mp$	LF	$[r] < 1.74$	$\times 10^{-6}$	CL=90%	863
$K^- \pi^+ e^\pm \mu^\mp$	LF	$[r] < 1.90$	$\times 10^{-6}$	CL=90%	848
$\overline{K}^*(892)^0 e^\pm \mu^\mp$	LF	$[r] < 1.25$	$\times 10^{-6}$	CL=90%	714
$2\pi^- 2e^+$	L	< 9.1	$\times 10^{-7}$	CL=90%	922
$2\pi^- 2\mu^+$	L	< 1.52	$\times 10^{-6}$	CL=90%	894
$K^- \pi^- 2e^+$	L	< 5.0	$\times 10^{-7}$	CL=90%	861
$K^- \pi^- 2\mu^+$	L	< 5.3	$\times 10^{-7}$	CL=90%	829
$2K^- 2e^+$	L	< 3.4	$\times 10^{-7}$	CL=90%	791

$2K^- 2\mu^+$	L	< 1.0	$\times 10^{-7}$	CL=90%	710
$\pi^- \pi^- e^+ \mu^+$	L	< 3.06	$\times 10^{-6}$	CL=90%	911
$K^- \pi^- e^+ \mu^+$	L	< 2.10	$\times 10^{-6}$	CL=90%	848
$2K^- e^+ \mu^+$	L	< 5.8	$\times 10^{-7}$	CL=90%	754
$p e^-$	L, B	$[s] < 1.0$	$\times 10^{-5}$	CL=90%	696
$\bar{p} e^+$	L, B	$[t] < 1.1$	$\times 10^{-5}$	CL=90%	696

 $D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

 I, J, P need confirmation.Mass $m = 2006.85 \pm 0.05$ MeV ($S = 1.1$) $m_{D^{*0}} - m_{D^0} = 142.014 \pm 0.030$ MeV ($S = 1.5$)Full width $\Gamma < 2.1$ MeV, CL = 90% $\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^0$	$(64.7 \pm 0.9) \%$	43
$D^0 \gamma$	$(35.3 \pm 0.9) \%$	137

 $D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

 I, J, P need confirmation.Mass $m = 2010.26 \pm 0.05$ MeV $m_{D^*(2010)^+} - m_{D^+} = 140.603 \pm 0.015$ MeV $m_{D^*(2010)^+} - m_{D^0} = 145.4258 \pm 0.0017$ MeVFull width $\Gamma = 83.4 \pm 1.8$ keV $D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	$(67.7 \pm 0.5) \%$	39
$D^+ \pi^0$	$(30.7 \pm 0.5) \%$	38
$D^+ \gamma$	$(1.6 \pm 0.4) \%$	136

 $D_0^*(2300)$

$$I(J^P) = \frac{1}{2}(0^+)$$

was $D_0^*(2400)$ Mass $m = 2343 \pm 10$ MeV ($S = 1.5$)Full width $\Gamma = 229 \pm 16$ MeV

$D_0^*(2300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D \pi^\pm$	seen	411

$D_1(2420)$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 2422.1 \pm 0.6$ MeV ($S = 1.7$) $m_{D_1(2420)^0} - m_{D^{*+}} = 411.8 \pm 0.6$ MeV ($S = 1.7$) $m_{D_1(2420)^\pm} - m_{D_1(2420)^0} = 4 \pm 4$ MeVFull width $\Gamma = 31.3 \pm 1.9$ MeV ($S = 2.8$) $\bar{D}_1(2420)$ modes are charge conjugates of modes below. **$D_1(2420)$ DECAY MODES**Fraction (Γ_i/Γ) p (MeV/c)

$D^*(2007)^0 \pi$	seen	359
-------------------	------	-----

 $D_1(2430)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 2412 \pm 9$ MeVFull width $\Gamma = 314 \pm 29$ MeV **$D_1(2430)^0$ DECAY MODES**Fraction (Γ_i/Γ) p (MeV/c)

$D^*(2010)^+ \pi^-$	seen	345
---------------------	------	-----

 $D_2^*(2460)$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $J^P = 2^+$ assignment strongly favored.Mass $m = 2461.1^{+0.7}_{-0.8}$ MeV ($S = 6.2$) $m_{D_2^*(2460)^0} - m_{D^+} = 591.5^{+0.7}_{-0.8}$ MeV ($S = 5.9$) $m_{D_2^*(2460)^0} - m_{D^{*+}} = 450.9^{+0.7}_{-0.8}$ MeV ($S = 5.9$) $m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7$ MeVFull width $\Gamma = 47.3 \pm 0.8$ MeV ($S = 1.5$) $\bar{D}_2^*(2460)$ modes are charge conjugates of modes below. **$D_2^*(2460)$ DECAY MODES**Fraction (Γ_i/Γ) p (MeV/c)

$D \pi^-$	seen	509
$D^*(2010) \pi^-$	seen	389

$D_3^*(2750)$

$$I(J^P) = \frac{1}{2}(3^-)$$

Mass $m = 2763.1 \pm 3.2$ MeV (S = 2.1)Full width $\Gamma = 66 \pm 5$ MeV

$D_3^*(2750)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D\pi$	seen	743
$D^+\pi^-$	seen	739
$D^0\pi^\pm$	seen	743
$D^*\pi$	seen	639
$D^{*+}\pi^-$	seen	639

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^-2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] Submodes of the $D^+ \rightarrow K^-2\pi^+\pi^0$ and $K_S^0 2\pi^+\pi^-$ modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+\ell^+\ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 *Review*, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^-2\pi^+\pi^-$, $K^-2\pi^+\pi^-\pi^0$, $\bar{K}^0 2\pi^+2\pi^-$, $K^+2K^-\pi^+$, $2\pi^+2\pi^-$, $2\pi^+2\pi^-\pi^0$, $K^+K^-\pi^+\pi^-$, and $K^+K^-\pi^+\pi^-\pi^0$, branching fractions.
- [l] This is the sum of our $K^-3\pi^+2\pi^-$ and $3\pi^+3\pi^-$ branching fractions.

- [n] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.17 ± 0.17 %.
- [o] This is a doubly Cabibbo-suppressed mode.
- [p] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [q] This branching fraction includes all the decay modes of the resonance in the final state.
- [r] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [s] This limit is for either D^0 or $\bar{D}^0$ to $p e^-$.
- [t] This limit is for either D^0 or $\bar{D}^0$ to $\bar{p} e^+$.