

LIGHT UNFLAVORED MESONS

($S = C = B = 0$)

For $I = 1$ (π , b , ρ , a): $u\bar{d}$, $(u\bar{u}-d\bar{d})/\sqrt{2}$, $d\bar{u}$;
for $I = 0$ (η , η' , h , h' , ω , ϕ , f , f'): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

$\pi^\pm$

$$J^G(J^P) = 1^-(0^-)$$

Mass $m = 139.57039 \pm 0.00018$ MeV ($S = 1.8$)

Mean life $\tau = (2.6033 \pm 0.0005) \times 10^{-8}$ s ($S = 1.2$)

$$c\tau = 7.8045 \text{ m}$$

$\pi^\pm \rightarrow \ell^\pm \nu \gamma$ form factors [a]

$$F_V = 0.0254 \pm 0.0017$$

$$F_A = 0.0119 \pm 0.0001$$

$$F_V \text{ slope parameter } a = 0.10 \pm 0.06$$

$$R = 0.059^{+0.009}_{-0.008}$$

π^- modes are charge conjugates of the modes below.

For decay limits to particles which are not established, see the section on Searches for Axions and Other Very Light Bosons.

π^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$\mu^+ \nu_\mu$	[b] (99.98770 $\pm$ 0.00004) %		30
$\mu^+ \nu_\mu \gamma$	[c] (2.00 $\pm$ 0.25) $\times 10^{-4}$		30
$e^+ \nu_e$	[b] (1.230 $\pm$ 0.004) $\times 10^{-4}$		70
$e^+ \nu_e \gamma$	[c] (7.39 $\pm$ 0.05) $\times 10^{-7}$		70
$e^+ \nu_e \pi^0$	(1.036 $\pm$ 0.006) $\times 10^{-8}$		4
$e^+ \nu_e e^+ e^-$	(3.2 $\pm$ 0.5) $\times 10^{-9}$		70
$\mu^+ \nu_\mu \nu \bar{\nu}$	< 9	$\times 10^{-6}$ 90%	30
$e^+ \nu_e \nu \bar{\nu}$	< 1.6	$\times 10^{-7}$ 90%	70
Lepton Family number (LF) or Lepton number (L) violating modes			
$\mu^+ \bar{\nu}_e$	L [d] < 1.5	$\times 10^{-3}$ 90%	30
$\mu^+ \nu_e$	LF [d] < 8.0	$\times 10^{-3}$ 90%	30
$\mu^- e^+ e^+ \nu$	LF < 1.6	$\times 10^{-6}$ 90%	30



$$I^G(J^{PC}) = 1^-(0^-\,+)$$

$$\text{Mass } m = 134.9768 \pm 0.0005 \text{ MeV} \quad (S = 1.1)$$

$$m_{\pi^\pm} - m_{\pi^0} = 4.5936 \pm 0.0005 \text{ MeV}$$

$$\text{Mean life } \tau = (8.43 \pm 0.13) \times 10^{-17} \text{ s} \quad (S = 1.2)$$

$$c\tau = 25.3 \text{ nm}$$

For decay limits to particles which are not established, see the appropriate Search sections (A^0 (axion) and Other Light Boson (X^0) Searches, etc.).

π^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)	
2γ	$(98.823\pm0.034) \%$	$S=1.5$	67	
$e^+ e^- \gamma$	$(1.174\pm0.035) \%$	$S=1.5$	67	
γ positronium	$(1.82 \pm 0.29) \times 10^{-9}$		67	
$e^+ e^+ e^- e^-$	$(3.34 \pm 0.16) \times 10^{-5}$		67	
$e^+ e^-$	$(6.46 \pm 0.33) \times 10^{-8}$		67	
4γ	< 2	$\times 10^{-8}$ CL=90%	67	
$\nu \bar{\nu}$	[e] < 2.7	$\times 10^{-7}$ CL=90%	67	
$\nu_e \bar{\nu}_e$	< 1.7	$\times 10^{-6}$ CL=90%	67	
$\nu_\mu \bar{\nu}_\mu$	< 1.6	$\times 10^{-6}$ CL=90%	67	
$\nu_\tau \bar{\nu}_\tau$	< 2.1	$\times 10^{-6}$ CL=90%	67	
$\gamma \nu \bar{\nu}$	< 1.9	$\times 10^{-7}$ CL=90%	67	
Charge conjugation (C) or Lepton Family number (LF) violating modes				
3γ	C	< 3.1	$\times 10^{-8}$ CL=90%	67
$\mu^+ e^-$	LF	< 3.8	$\times 10^{-10}$ CL=90%	26
$\mu^- e^+$	LF	< 3.4	$\times 10^{-9}$ CL=90%	26
$\mu^+ e^- + \mu^- e^+$	LF	< 3.6	$\times 10^{-10}$ CL=90%	26



$$I^G(J^{PC}) = 0^+(0^-\,+)$$

$$\text{Mass } m = 547.862 \pm 0.017 \text{ MeV}$$

$$\text{Full width } \Gamma = 1.31 \pm 0.05 \text{ keV}$$

C-nonconserving decay parameters

$$\pi^+ \pi^- \pi^0 \quad \text{left-right asymmetry} = (0.09^{+0.11}_{-0.12}) \times 10^{-2}$$

$$\pi^+ \pi^- \pi^0 \quad \text{sextant asymmetry} = (0.12^{+0.10}_{-0.11}) \times 10^{-2}$$

$$\pi^+ \pi^- \pi^0 \quad \text{quadrant asymmetry} = (-0.09 \pm 0.09) \times 10^{-2}$$

$$\pi^+ \pi^- \gamma \quad \text{left-right asymmetry} = (0.9 \pm 0.4) \times 10^{-2}$$

$$\pi^+ \pi^- \gamma \quad \beta (D\text{-wave}) = -0.02 \pm 0.07 \quad (S = 1.3)$$

CP-nonconserving decay parameters

$$\pi^+ \pi^- e^+ e^- \text{ decay-plane asymmetry } A_\phi = (-0.6 \pm 3.1) \times 10^{-2}$$

Other decay parameters $\pi^0\pi^0\pi^0$ Dalitz plot $\alpha = -0.0288 \pm 0.0012$ ($S = 1.1$)Parameter Λ in $\eta \rightarrow \ell^+\ell^-\gamma$ decay = $0.716 \pm 0.011 \text{ GeV}/c^2$

η DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Neutral modes			
neutral modes	$(72.12 \pm 0.34) \%$	$S=1.2$	—
2γ	$(39.41 \pm 0.20) \%$	$S=1.1$	274
$3\pi^0$	$(32.68 \pm 0.23) \%$	$S=1.1$	179
$\pi^0 2\gamma$	$(2.56 \pm 0.22) \times 10^{-4}$		257
$2\pi^0 2\gamma$	$< 1.2 \times 10^{-3}$	CL=90%	238
4γ	$< 2.8 \times 10^{-4}$	CL=90%	274
invisible	$< 1.0 \times 10^{-4}$	CL=90%	—
Charged modes			
charged modes	$(27.89 \pm 0.29) \%$	$S=1.2$	—
$\pi^+\pi^-\pi^0$	$(22.92 \pm 0.28) \%$	$S=1.2$	174
$\pi^+\pi^-\gamma$	$(4.22 \pm 0.08) \%$	$S=1.1$	236
$e^+e^-\gamma$	$(6.9 \pm 0.4) \times 10^{-3}$	$S=1.3$	274
$\mu^+\mu^-\gamma$	$(3.1 \pm 0.4) \times 10^{-4}$		253
e^+e^-	$< 7 \times 10^{-7}$	CL=90%	274
$\mu^+\mu^-$	$(5.8 \pm 0.8) \times 10^{-6}$		253
$2e^+2e^-$	$(2.40 \pm 0.22) \times 10^{-5}$		274
$\pi^+\pi^-e^+e^-(\gamma)$	$(2.68 \pm 0.11) \times 10^{-4}$		235
$e^+e^-\mu^+\mu^-$	$< 1.6 \times 10^{-4}$	CL=90%	253
$2\mu^+2\mu^-$	$< 3.6 \times 10^{-4}$	CL=90%	161
$\mu^+\mu^-\pi^+\pi^-$	$< 3.6 \times 10^{-4}$	CL=90%	113
$\pi^+e^-\bar{\nu}_e + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90%	256
$\pi^+\pi^-2\gamma$	$< 2.1 \times 10^{-3}$		236
$\pi^+\pi^-\pi^0\gamma$	$< 5 \times 10^{-4}$	CL=90%	174
$\pi^0\mu^+\mu^-\gamma$	$< 3 \times 10^{-6}$	CL=90%	210

**Charge conjugation (C), Parity (P),
Charge conjugation $\times$ Parity (CP), or
Lepton Family number (LF) violating modes**

$\pi^0\gamma$	C	$[f] < 9$	$\times 10^{-5}$	CL=90%	257
$\pi^+\pi^-$	P, CP	< 4.4	$\times 10^{-6}$	CL=90%	236
$2\pi^0$	P, CP	< 3.5	$\times 10^{-4}$	CL=90%	238
$2\pi^0\gamma$	C	< 5	$\times 10^{-4}$	CL=90%	238
$3\pi^0\gamma$	C	< 6	$\times 10^{-5}$	CL=90%	179
3γ	C	< 1.6	$\times 10^{-5}$	CL=90%	274
$4\pi^0$	P, CP	< 6.9	$\times 10^{-7}$	CL=90%	40
$\pi^0e^+e^-$	C	$[g] < 8$	$\times 10^{-6}$	CL=90%	257

$\pi^0 \mu^+ \mu^-$	C	$[g] < 5$	$\times 10^{-6}$	CL=90%	210
$\mu^+ e^- + \mu^- e^+$	LF	< 6	$\times 10^{-6}$	CL=90%	264

 $f_0(500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

also known as σ ; was $f_0(600)$

See the review on "Scalar Mesons below 2 GeV."

Mass (T-Matrix Pole $\sqrt{s}$) = (400–550)– i (200–350) MeV

Mass (Breit-Wigner) = 400 to 800 MeV

Full width (Breit-Wigner) = 100 to 800 MeV

$f_0(500)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi$	seen	–
$\gamma\gamma$	seen	–

 $\rho(770)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

See the note in $\rho(770)$ Particle Listings.Mass $m = 775.26 \pm 0.23$ MeVFull width $\Gamma = 149.1 \pm 0.8$ MeV

$\rho(770)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi\pi$	~ 100	%	363
$\rho(770)^\pm$ decays			
$\pi^\pm \gamma$	(4.5 ± 0.5) $\times 10^{-4}$	S=2.2	375
$\pi^\pm \eta$	< 6	CL=84%	152
$\pi^\pm \pi^+ \pi^- \pi^0$	< 2.0	CL=84%	254
$\rho(770)^0$ decays			
$\pi^+ \pi^- \gamma$	(9.9 ± 1.6) $\times 10^{-3}$		362
$\pi^0 \gamma$	(4.7 ± 0.8) $\times 10^{-4}$	S=1.7	376
$\eta \gamma$	(3.00 ± 0.21) $\times 10^{-4}$		194
$\pi^0 \pi^0 \gamma$	(4.5 ± 0.8) $\times 10^{-5}$		363
$\mu^+ \mu^-$	[h] (4.55 ± 0.28) $\times 10^{-5}$		373
$e^+ e^-$	[h] (4.72 ± 0.05) $\times 10^{-5}$		388
$\pi^+ \pi^- \pi^0$	($1.01^{+0.54}_{-0.36} \pm 0.34$) $\times 10^{-4}$		323
$\pi^+ \pi^- \pi^+ \pi^-$	(1.8 ± 0.9) $\times 10^{-5}$		251
$\pi^+ \pi^- \pi^0 \pi^0$	(1.6 ± 0.8) $\times 10^{-5}$		257
$\pi^0 e^+ e^-$	< 1.2	CL=90%	376

$\omega(782)$

$$J^{PC} = 0^-(1^--)$$

Mass $m = 782.66 \pm 0.13$ MeV ($S = 2.0$)Full width $\Gamma = 8.68 \pm 0.13$ MeV

$\omega(782)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi^+\pi^-\pi^0$	$(89.2 \pm 0.7) \%$		327
$\pi^0\gamma$	$(8.34 \pm 0.26) \%$	S=2.1	380
$\pi^+\pi^-$	$(1.53^{+0.11}_{-0.13}) \%$	S=1.2	366
neutrals (excluding $\pi^0\gamma$)	$(7^{+7}_{-5}) \times 10^{-3}$	S=1.1	—
$\eta\gamma$	$(4.5 \pm 0.4) \times 10^{-4}$	S=1.1	200
$\pi^0 e^+ e^-$	$(7.7 \pm 0.6) \times 10^{-4}$		380
$\pi^0 \mu^+ \mu^-$	$(1.34 \pm 0.18) \times 10^{-4}$	S=1.5	349
$e^+ e^-$	$(7.39 \pm 0.19) \times 10^{-5}$	S=1.7	391
$\pi^+\pi^-\pi^0\pi^0$	$< 2 \times 10^{-4}$	CL=90%	262
$\pi^+\pi^-\gamma$	$< 3.6 \times 10^{-3}$	CL=95%	366
$\pi^+\pi^-\pi^+\pi^-$	$< 1 \times 10^{-3}$	CL=90%	256
$\pi^0\pi^0\gamma$	$(6.7 \pm 1.1) \times 10^{-5}$		367
$\eta\pi^0\gamma$	$< 3.3 \times 10^{-5}$	CL=90%	162
$\mu^+\mu^-$	$(7.4 \pm 1.8) \times 10^{-5}$		377
3γ	$< 1.9 \times 10^{-4}$	CL=95%	391

Charge conjugation (C) violating modes

$\eta\pi^0$	C	$< 2.1 \times 10^{-4}$	CL=90%	162
$2\pi^0$	C	$< 2.2 \times 10^{-4}$	CL=90%	367
$3\pi^0$	C	$< 2.3 \times 10^{-4}$	CL=90%	330
invisible		$< 7 \times 10^{-5}$	CL=90%	—

 $\eta'(958)$

$$J^{PC} = 0^+(0^-+)$$

Mass $m = 957.78 \pm 0.06$ MeVFull width $\Gamma = 0.188 \pm 0.006$ MeV

$\eta'(958)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\pi^+\pi^-\eta$	$(42.5 \pm 0.5) \%$		232
$\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	$(29.5 \pm 0.4) \%$		165
$\pi^0\pi^0\eta$	$(22.4 \pm 0.5) \%$		239
$\omega\gamma$	$(2.52 \pm 0.07) \%$		159
$\omega e^+ e^-$	$(2.0 \pm 0.4) \times 10^{-4}$		159
$\gamma\gamma$	$(2.307 \pm 0.033) \%$		479
$3\pi^0$	$(2.50 \pm 0.17) \times 10^{-3}$		430

$\mu^+ \mu^- \gamma$	$(1.13 \pm 0.28) \times 10^{-4}$			467
$\pi^+ \pi^- \mu^+ \mu^-$	$< 2.9 \times 10^{-5}$	90%		401
$\pi^+ \pi^- \pi^0$	$(3.61 \pm 0.17) \times 10^{-3}$			428
$(\pi^+ \pi^- \pi^0)$ S-wave	$(3.8 \pm 0.5) \times 10^{-3}$			428
$\pi^\mp \rho^\pm$	$(7.4 \pm 2.3) \times 10^{-4}$			106
$\pi^0 \rho^0$	$< 4 \%$	90%		111
$2(\pi^+ \pi^-)$	$(8.4 \pm 0.9) \times 10^{-5}$			372
$\pi^+ \pi^- 2\pi^0$	$(1.8 \pm 0.4) \times 10^{-4}$			376
$2(\pi^+ \pi^-)$ neutrals	$< 1 \%$	95%		—
$2(\pi^+ \pi^-) \pi^0$	$< 1.8 \times 10^{-3}$	90%		298
$2(\pi^+ \pi^-) 2\pi^0$	$< 1 \%$	95%		197
$3(\pi^+ \pi^-)$	$< 3.1 \times 10^{-5}$	90%		189
$K^\pm \pi^\mp$	$< 4 \times 10^{-5}$	90%		334
$\pi^+ \pi^- e^+ e^-$	$(2.4^{+1.3}_{-1.0}) \times 10^{-3}$			458
$\pi^+ e^- \nu_e + \text{c.c.}$	$< 2.1 \times 10^{-4}$	90%		469
$\gamma e^+ e^-$	$(4.91 \pm 0.27) \times 10^{-4}$			479
$\pi^0 \gamma \gamma$	$(3.20 \pm 0.24) \times 10^{-3}$			469
$\pi^0 \gamma \gamma (\text{non resonant})$	$(6.2 \pm 0.9) \times 10^{-4}$			—
$\eta \gamma \gamma$	$< 1.33 \times 10^{-4}$	90%		322
$4\pi^0$	$< 4.94 \times 10^{-5}$	90%		380
$e^+ e^-$	$< 5.6 \times 10^{-9}$	90%		479
invisible	$< 6 \times 10^{-4}$	90%		—

**Charge conjugation (C), Parity (P),
Lepton family number (LF) violating modes**

$\pi^+ \pi^-$	P, CP	$< 1.8 \times 10^{-5}$	90%	458
$\pi^0 \pi^0$	P, CP	$< 4 \times 10^{-4}$	90%	459
$\pi^0 e^+ e^-$	C [g]	$< 1.4 \times 10^{-3}$	90%	469
$\eta e^+ e^-$	C [g]	$< 2.4 \times 10^{-3}$	90%	322
3γ	C	$< 1.0 \times 10^{-4}$	90%	479
$\mu^+ \mu^- \pi^0$	C [g]	$< 6.0 \times 10^{-5}$	90%	445
$\mu^+ \mu^- \eta$	C [g]	$< 1.5 \times 10^{-5}$	90%	273
$e \mu$	LF	$< 4.7 \times 10^{-4}$	90%	473

$f_0(980)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Scalar Mesons below 2 GeV."

Mass $m = 990 \pm 20$ MeV

Full width $\Gamma = 10$ to 100 MeV

$f_0(980)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi \pi$	seen	476

$K\bar{K}$	seen	36
$\gamma\gamma$	seen	495

 $a_0(980)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

See the review on "Scalar Mesons below 2 GeV."

Mass $m = 980 \pm 20$ MeVFull width $\Gamma = 50$ to 100 MeV

$a_0(980)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta\pi$	seen	319
$K\bar{K}$	seen	†
$\eta'\pi$	seen	†
$\rho\pi$	not seen	137
$\gamma\gamma$	seen	490

 $\phi(1020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 1019.461 \pm 0.016$ MeVFull width $\Gamma = 4.249 \pm 0.013$ MeV ($S = 1.1$)

$\phi(1020)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
K^+K^-	(49.2 $\pm$ 0.5) %	S=1.3	127
$K_L^0 K_S^0$	(34.0 $\pm$ 0.4) %	S=1.3	110
$\rho\pi^+ \pi^+ \pi^- \pi^0$	(15.24 $\pm$ 0.33) %	S=1.2	—
$\eta\gamma$	(1.303 $\pm$ 0.025) %	S=1.2	363
$\pi^0\gamma$	(1.32 $\pm$ 0.06) $\times 10^{-3}$		501
$\ell^+ \ell^-$	—		510
$e^+ e^-$	(2.974 $\pm$ 0.034) $\times 10^{-4}$	S=1.3	510
$\mu^+ \mu^-$	(2.86 $\pm$ 0.19) $\times 10^{-4}$		499
$\eta e^+ e^-$	(1.08 $\pm$ 0.04) $\times 10^{-4}$		363
$\pi^+ \pi^-$	(7.3 $\pm$ 1.3) $\times 10^{-5}$		490
$\omega\pi^0$	(4.7 $\pm$ 0.5) $\times 10^{-5}$		171
$\omega\gamma$	< 5 %	CL=84%	209
$\rho\gamma$	< 1.2 $\times 10^{-5}$	CL=90%	215
$\pi^+ \pi^- \gamma$	(4.1 $\pm$ 1.3) $\times 10^{-5}$		490
$f_0(980)\gamma$	(3.22 $\pm$ 0.19) $\times 10^{-4}$	S=1.1	29
$\pi^0\pi^0\gamma$	(1.12 $\pm$ 0.06) $\times 10^{-4}$		492
$\pi^+ \pi^- \pi^+ \pi^-$	(3.9 $^{+2.8}_{-2.2}$) $\times 10^{-6}$		410
$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 4.6 $\times 10^{-6}$	CL=90%	342

$\pi^0 e^+ e^-$	$(1.33^{+0.07}_{-0.10}) \times 10^{-5}$		501
$\pi^0 \eta \gamma$	$(7.27 \pm 0.30) \times 10^{-5}$	S=1.5	346
$a_0(980) \gamma$	$(7.6 \pm 0.6) \times 10^{-5}$		39
$K^0 \bar{K}^0 \gamma$	$< 1.9 \times 10^{-8}$	CL=90%	110
$\eta'(958) \gamma$	$(6.22 \pm 0.21) \times 10^{-5}$		60
$\eta \pi^0 \pi^0 \gamma$	$< 2 \times 10^{-5}$	CL=90%	293
$\mu^+ \mu^- \gamma$	$(1.4 \pm 0.5) \times 10^{-5}$		499
$\rho \gamma \gamma$	$< 1.2 \times 10^{-4}$	CL=90%	215
$\eta \pi^+ \pi^-$	$< 1.8 \times 10^{-5}$	CL=90%	288
$\eta \mu^+ \mu^-$	$< 9.4 \times 10^{-6}$	CL=90%	321
$\eta U \rightarrow \eta e^+ e^-$	$< 1 \times 10^{-6}$	CL=90%	—
invisible	$< 1.7 \times 10^{-4}$	CL=90%	—

Lepton Family number (LF) violating modes

$e^\pm \mu^\mp$	LF	$< 2 \times 10^{-6}$	CL=90%	504
-----------------	----	----------------------	--------	-----

 $h_1(1170)$

$$I^G(J^{PC}) = 0^-(1^+ -)$$

Mass $m = 1166 \pm 6$ MeVFull width $\Gamma = 375 \pm 35$ MeV

$h_1(1170)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho \pi$	seen	305

 $b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^+ -)$$

Mass $m = 1229.5 \pm 3.2$ MeV (S = 1.6)Full width $\Gamma = 142 \pm 9$ MeV (S = 1.2)

$b_1(1235)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\omega \pi$	seen		348
[D/S amplitude ratio = 0.277 ± 0.027]			
$\pi^\pm \gamma$	$(1.6 \pm 0.4) \times 10^{-3}$		607
$\eta \rho$	seen		†
$\pi^+ \pi^+ \pi^- \pi^0$	< 50 %	84%	535
$K^*(892)^\pm K^\mp$	seen		†
$(K\bar{K})^\pm \pi^0$	< 8 %	90%	248
$K_S^0 K_L^0 \pi^\pm$	< 6 %	90%	235
$K_S^0 K_S^0 \pi^\pm$	< 2 %	90%	235
$\phi \pi$	< 1.5 %	84%	147

$a_1(1260)$ [i]

$$I^G(J^{PC}) = 1^-(1^{++})$$

Mass $m = 1230 \pm 40$ MeV [i]Full width $\Gamma = 250$ to 600 MeV

$a_1(1260)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
3π	seen	577
$(\rho\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi$	seen	353
$(\rho\pi)_{D\text{-wave}}, \rho \rightarrow \pi\pi$	seen	353
$(\rho(1450)\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi$	seen	†
$(\rho(1450)\pi)_{D\text{-wave}}, \rho \rightarrow \pi\pi$	seen	†
$f_0(500)\pi, f_0 \rightarrow \pi\pi$	seen	—
$f_0(980)\pi, f_0 \rightarrow \pi\pi$	not seen	179
$f_0(1370)\pi, f_0 \rightarrow \pi\pi$	seen	†
$f_2(1270)\pi, f_2 \rightarrow \pi\pi$	seen	†
$\pi^+\pi^-\pi^0$	seen	576
$\pi^0\pi^0\pi^0$	not seen	577
$KK\pi$	seen	250
$K^*(892)K$	seen	†
$\pi\gamma$	seen	608

 $f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 1275.5 \pm 0.8$ MeVFull width $\Gamma = 186.7^{+2.2}_{-2.5}$ MeV ($S = 1.4$)

$f_2(1270)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\pi\pi$	$(84.2^{+2.9}_{-0.9})\%$	$S=1.1$	623
$\pi^+\pi^-\pi^0$	$(7.7^{+1.1}_{-3.2})\%$	$S=1.2$	563
$K\bar{K}$	$(4.6^{+0.5}_{-0.4})\%$	$S=2.7$	404
$2\pi^+2\pi^-$	$(2.8 \pm 0.4)\%$	$S=1.2$	560
$\eta\eta$	$(4.0 \pm 0.8) \times 10^{-3}$	$S=2.1$	326
$4\pi^0$	$(3.0 \pm 1.0) \times 10^{-3}$		565
$\gamma\gamma$	$(1.42 \pm 0.24) \times 10^{-5}$	$S=1.4$	638
$\eta\pi\pi$	$< 8 \times 10^{-3}$	CL=95%	478
$K^0K^-\pi^+ + \text{c.c.}$	$< 3.4 \times 10^{-3}$	CL=95%	293
e^+e^-	$< 6 \times 10^{-10}$	CL=90%	638

$f_1(1285)$

$$J^{PC} = 0^+(1^+ +)$$

Mass $m = 1281.9 \pm 0.5$ MeV (S = 1.8)Full width $\Gamma = 22.7 \pm 1.1$ MeV (S = 1.5)

$f_1(1285)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
4π	$(32.7 \pm 1.9) \%$	S=1.2	568
$\pi^0 \pi^0 \pi^+ \pi^-$	$(21.8 \pm 1.3) \%$	S=1.2	566
$2\pi^+ 2\pi^-$	$(10.9 \pm 0.6) \%$	S=1.2	563
$\rho^0 \pi^+ \pi^-$	$(10.9 \pm 0.6) \%$	S=1.2	336
$\rho^0 \rho^0$	seen		†
$4\pi^0$	$< 7 \times 10^{-4}$	CL=90%	568
$\eta \pi^+ \pi^-$	$(35 \pm 15) \%$		479
$\eta \pi \pi$	$(52.2 \pm 2.0) \%$	S=1.2	482
$a_0(980) \pi$ [ignoring $a_0(980) \rightarrow K \bar{K}$]	$(38 \pm 4) \%$		238
$\eta \pi \pi$ [excluding $a_0(980) \pi$]	$(14 \pm 4) \%$		482
$K \bar{K} \pi$	$(9.0 \pm 0.4) \%$	S=1.1	308
$K \bar{K}^*(892)$	not seen		†
$\pi^+ \pi^- \pi^0$	$(3.0 \pm 0.9) \times 10^{-3}$		603
$\rho^\pm \pi^\mp$	$< 3.1 \times 10^{-3}$	CL=95%	390
$\gamma \rho^0$	$(6.1 \pm 1.0) \%$	S=1.7	406
$\phi \gamma$	$(7.4 \pm 2.6) \times 10^{-4}$		236
$e^+ e^-$	$< 9.4 \times 10^{-9}$	CL=90%	641

 $\eta(1295)$

$$J^{PC} = 0^+(0^- +)$$

See the review on "Pseudoscalar and pseudovector mesons in the 1400 MeV region."

Mass $m = 1294 \pm 4$ MeV (S = 1.6)Full width $\Gamma = 55 \pm 5$ MeV

$\eta(1295)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta \pi^+ \pi^-$	seen	487
$a_0(980) \pi$	seen	248
$\eta \pi^0 \pi^0$	seen	490
$\eta(\pi\pi)_{S\text{-wave}}$	seen	—

 $\pi(1300)$

$$J^{PC} = 1^-(0^- +)$$

Mass $m = 1300 \pm 100$ MeV [?]Full width $\Gamma = 200$ to 600 MeV

$\pi(1300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	404
$\pi(\pi\pi)_{S\text{-wave}}$	seen	—

 $a_2(1320)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

Mass $m = 1318.2 \pm 0.6$ MeV (S = 1.2)Full width $\Gamma = 107 \pm 5$ MeV [1]

$a_2(1320)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
3π	$(70.1 \pm 2.7) \%$	S=1.2	624
$\eta\pi$	$(14.5 \pm 1.2) \%$		535
$\omega\pi\pi$	$(10.6 \pm 3.2) \%$	S=1.3	366
$K\bar{K}$	$(4.9 \pm 0.8) \%$		437
$\eta'(958)\pi$	$(5.5 \pm 0.9) \times 10^{-3}$		288
$\pi^\pm\gamma$	$(2.91 \pm 0.27) \times 10^{-3}$		652
$\gamma\gamma$	$(9.4 \pm 0.7) \times 10^{-6}$		659
e^+e^-	$< 5 \times 10^{-9}$	CL=90%	659

 $f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Scalar Mesons below 2 GeV."

Mass $m = 1200$ to 1500 MeVFull width $\Gamma = 200$ to 500 MeV

$f_0(1370)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi$	seen	672
4π	seen	617
$4\pi^0$	seen	617
$2\pi^+2\pi^-$	seen	612
$\pi^+\pi^-2\pi^0$	seen	615
$\rho\rho$	seen	†
$2(\pi\pi)_{S\text{-wave}}$	seen	—
$\pi(1300)\pi$	seen	†
$a_1(1260)\pi$	seen	35
$\eta\eta$	seen	411
$K\bar{K}$	seen	475
$K\bar{K}n\pi$	not seen	†
6π	not seen	508
$\omega\omega$	not seen	†

$\gamma\gamma$	seen	685
e^+e^-	not seen	685

 $\pi_1(1400)$

$$I^G(J^{PC}) = 1^-(1^--^+)$$

See the review on "Non- $q\bar{q}$ Mesons." See also $\pi_1(1600)$.

Mass $m = 1354 \pm 25$ MeV ($S = 1.8$)

Full width $\Gamma = 330 \pm 35$ MeV

$\pi_1(1400)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta\pi^0$	seen	557
$\eta\pi^-$	seen	556
$\rho(770)\pi$	not seen	442

 $\eta(1405)$

$$I^G(J^{PC}) = 0^+(0^--^+)$$

See the review on "Pseudoscalar and Pseudovector Mesons in the 1400 MeV Region." See also $\eta(1475)$.

Mass $m = 1408.8 \pm 2.0$ MeV ($S = 2.2$)

Full width $\Gamma = 50.1 \pm 2.6$ MeV ($S = 1.7$)

$\eta(1405)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$K\bar{K}\pi$	seen		424
$\eta\pi\pi$	seen		562
$a_0(980)\pi$	seen		345
$\eta(\pi\pi)_{S\text{-wave}}$	seen		—
$f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$	not seen		—
$f_0(980)\eta$	seen		†
4π	seen		639
$\rho\rho$	<58 %	99.85%	†
$\rho^0\gamma$	seen		491
$K^*(892)K$	seen		123

 $h_1(1415)$

$$I^G(J^{PC}) = 0^-(1^+-)$$

was $h_1(1380)$

Mass $m = 1416 \pm 8$ MeV ($S = 1.5$)

Full width $\Gamma = 90 \pm 15$ MeV

$f_1(1420)$

$$J^{PC} = 0^+(1^{++})$$

See the review on "Pseudoscalar and Pseudovector Mesons in the 1400 MeV Region."

$$\text{Mass } m = 1426.3 \pm 0.9 \text{ MeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma = 54.5 \pm 2.6 \text{ MeV}$$

$f_1(1420)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K \bar{K} \pi$	seen	438
$K \bar{K}^*(892) + \text{c.c.}$	seen	163
$\eta \pi \pi$	possibly seen	573
$\phi \gamma$	seen	349

 $\omega(1420)$ ^[k]

$$J^{PC} = 0^-(1^{--})$$

$$\text{Mass } m = 1410 \pm 60 \text{ MeV} \quad [j]$$

$$\text{Full width } \Gamma = 290 \pm 190 \text{ MeV} \quad [j]$$

$\omega(1420)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho \pi$	seen	480
$\omega \pi \pi$	seen	437
$b_1(1235) \pi$	seen	112
$e^+ e^-$	seen	705

 $a_0(1450)$

$$J^{PC} = 1^-(0^{++})$$

See the review on "Scalar Mesons below 2 GeV."

$$\text{Mass } m = 1474 \pm 19 \text{ MeV}$$

$$\text{Full width } \Gamma = 265 \pm 13 \text{ MeV}$$

$a_0(1450)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi \eta$	0.093 ± 0.020	627
$\pi \eta'(958)$	0.033 ± 0.017	410
$K \bar{K}$	0.082 ± 0.028	547
$\omega \pi \pi$	DEFINED AS 1	484
$a_0(980) \pi \pi$	seen	342
$\gamma \gamma$	seen	737

$\rho(1450)$

$$J^{PC} = 1^{+}(1^{-}-)$$

See the note in $\rho(1450)$ Particle Listings.

Mass $m = 1465 \pm 25$ MeV [j]

Full width $\Gamma = 400 \pm 60$ MeV [j]

$\rho(1450)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi$	seen	720
$\pi^+\pi^-$	seen	719
4π	seen	669
e^+e^-	seen	732
$\eta\rho$	seen	311
$a_2(1320)\pi$	not seen	55
$K\bar{K}$	seen	541
K^+K^-	seen	541
$K\bar{K}^*(892) + \text{c.c.}$	possibly seen	229
$\eta\gamma$	seen	630
$f_0(500)\gamma$	not seen	—
$f_0(980)\gamma$	not seen	398
$f_0(1370)\gamma$	not seen	92
$f_2(1270)\gamma$	not seen	177

 $\eta(1475)$

$$J^{PC} = 0^{+}(0^{-}+)$$

See the review on "Pseudoscalar and Pseudovector Mesons in the 1400 MeV Region." See also $\eta(1405)$.

Mass $m = 1475 \pm 4$ MeV ($S = 1.4$)

Full width $\Gamma = 90 \pm 9$ MeV ($S = 1.6$)

$\eta(1475)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}\pi$	seen	477
$K\bar{K}^*(892) + \text{c.c.}$	seen	244
$a_0(980)\pi$	seen	396
$\gamma\gamma$	seen	738
$K_S^0 K_S^0 \eta$	possibly seen	†
$\gamma\phi(1020)$	possibly seen	385

$f_0(1500)$

$$J^{PC} = 0^{+}(0^{+}+)$$

See the reviews on "Scalar Mesons below 2 GeV" and on "Non- $q\bar{q}$ Mesons".

Mass $m = 1506 \pm 6$ MeV ($S = 1.4$)

Full width $\Gamma = 112 \pm 9$ MeV

$f_0(1500)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	P (MeV/c)
$\pi\pi$	$(34.5 \pm 2.2) \%$	1.2	741
$\pi^+\pi^-$	seen		740
$2\pi^0$	seen		741
4π	$(48.9 \pm 3.3) \%$	1.2	692
$4\pi^0$	seen		692
$2\pi^+2\pi^-$	seen		687
$2(\pi\pi)_{S\text{-wave}}$	seen		—
$\rho\rho$	seen		†
$\pi(1300)\pi$	seen		145
$a_1(1260)\pi$	seen		219
$\eta\eta$	$(6.0 \pm 0.9) \%$	1.1	517
$\eta\eta'(958)$	$(2.2 \pm 0.8) \%$	1.4	20
$K\bar{K}$	$(8.5 \pm 1.0) \%$	1.1	569
$\gamma\gamma$	not seen		753

 $f'_2(1525)$

$$J^{PC} = 0^{+}(2^{+}+)$$

Mass $m = 1517.4 \pm 2.5$ MeV ($S = 2.8$)

Full width $\Gamma = 86 \pm 5$ MeV ($S = 2.2$)

$f'_2(1525)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	P (MeV/c)
$K\bar{K}$	$(87.6 \pm 2.2) \%$	1.1	576
$\eta\eta$	$(11.6 \pm 2.2) \%$	1.1	525
$\pi\pi$	$(8.3 \pm 1.6) \times 10^{-3}$		747
$\gamma\gamma$	$(9.5 \pm 1.1) \times 10^{-7}$	1.1	759

 $\pi_1(1600)$

$$J^{PC} = 1^{-}(1^{-}+)$$

See the review on "Non- $q\bar{q}$ Mesons" and a note in PDG 06, Journal of Physics **G33** 1 (2006). See also $\pi_1(1400)$.

Mass $m = 1661^{+15}_{-11}$ MeV ($S = 1.2$)

Full width $\Gamma = 240 \pm 50$ MeV ($S = 1.7$)

$\pi_1(1600)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi\pi$	seen	803
$\rho^0\pi^-$	seen	641
$f_2(1270)\pi^-$	not seen	318
$b_1(1235)\pi$	seen	357
$\eta'(958)\pi^-$	seen	543
$f_1(1285)\pi$	seen	314

 $a_1(1640)$

$$J^{PC} = 1^-(1^{++})$$

Mass $m = 1655 \pm 16$ MeV ($S = 1.2$)Full width $\Gamma = 254 \pm 40$ MeV ($S = 1.8$)

$a_1(1640)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi\pi\pi$	seen	800
$f_2(1270)\pi$	seen	314
$\sigma\pi$	seen	—
$\rho\pi$ S -wave	seen	638
$\rho\pi$ D -wave	seen	638
$\omega\pi\pi$	seen	607
$f_1(1285)\pi$	seen	309
$a_1(1260)\eta$	not seen	†

 $\eta_2(1645)$

$$J^{PC} = 0^+(2^{-+})$$

Mass $m = 1617 \pm 5$ MeVFull width $\Gamma = 181 \pm 11$ MeV

$\eta_2(1645)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$a_2(1320)\pi$	seen	242
$K\bar{K}\pi$	seen	580
$K^*\bar{K}$	seen	404
$\eta\pi^+\pi^-$	seen	685
$a_0(980)\pi$	seen	499
$f_2(1270)\eta$	not seen	†

 $\omega(1650)$ [1]

$$J^{PC} = 0^-(1^{--})$$

Mass $m = 1670 \pm 30$ MeV [1]Full width $\Gamma = 315 \pm 35$ MeV [1]

$\omega(1650)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	647
$\rho(1450)\pi$	seen	145
$\omega\pi\pi$	seen	617
$\omega\eta$	seen	500
e^+e^-	seen	835
$\pi^0\gamma$	not seen	830

 $\omega_3(1670)$

$$J^{PC} = 0^-(3^--)$$

Mass $m = 1667 \pm 4$ MeVFull width $\Gamma = 168 \pm 10$ MeV

$\omega_3(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\rho\pi$	seen	645
$\omega\pi\pi$	seen	615
$b_1(1235)\pi$	possibly seen	361

 $\pi_2(1670)$

$$J^{PC} = 1^-(2^-+)$$

Mass $m = 1670.6^{+2.9}_{-1.2}$ MeV ($S = 1.3$)Full width $\Gamma = 258^{+8}_{-9}$ MeV ($S = 1.2$)

$\pi_2(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
3π	$(95.8 \pm 1.4) \%$		808
$f_2(1270)\pi$	$(56.3 \pm 3.2) \%$		327
$\rho\pi$	$(31 \pm 4) \%$		647
$\sigma\pi$	$(10 \pm 4) \%$		—
$\pi(\pi\pi)_{S\text{-wave}}$	$(8.7 \pm 3.4) \%$		—
$\pi^\pm\pi^+\pi^-$	$(53 \pm 4) \%$		806
$K\bar{K}^*(892) + \text{c.c.}$	$(4.2 \pm 1.4) \%$		453
$\omega\rho$	$(2.7 \pm 1.1) \%$		302
$\pi^\pm\gamma$	$(7.0 \pm 1.2) \times 10^{-4}$		829
$\gamma\gamma$	$< 2.8 \times 10^{-7}$	90%	835
$\eta\pi$	$< 5 \%$		739
$\pi^\pm 2\pi^+ 2\pi^-$	$< 5 \%$		735
$\rho(1450)\pi$	$< 3.6 \times 10^{-3}$	97.7%	145
$b_1(1235)\pi$	$< 1.9 \times 10^{-3}$	97.7%	364
$f_1(1285)\pi$	possibly seen		322
$a_2(1320)\pi$	not seen		291

$\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^--)$$

Mass $m = 1680 \pm 20$ MeV [1]Full width $\Gamma = 150 \pm 50$ MeV [1]

$\phi(1680)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K \bar{K}^*(892) + \text{c.c.}$	seen	462
$K_S^0 K \pi$	seen	621
$K \bar{K}$	seen	680
$e^+ e^-$	seen	840
$\omega \pi \pi$	not seen	623
$K^+ K^- \pi^+ \pi^-$	seen	544
$\eta \phi$	seen	290
$\eta \gamma$	seen	751

 $\rho_3(1690)$

$$I^G(J^{PC}) = 1^+(3^--)$$

Mass $m = 1688.8 \pm 2.1$ MeVFull width $\Gamma = 161 \pm 10$ MeV (S = 1.5)

$\rho_3(1690)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	p (MeV/c)
4π	(71.1 $\pm$ 1.9) %		790
$\pi^\pm \pi^+ \pi^- \pi^0$	(67 $\pm$ 22) %		787
$\omega \pi$	(16 $\pm$ 6) %		655
$\pi \pi$	(23.6 $\pm$ 1.3) %		834
$K \bar{K} \pi$	(3.8 $\pm$ 1.2) %		629
$K \bar{K}$	(1.58 $\pm$ 0.26) %	1.2	685
$\eta \pi^+ \pi^-$	seen		727
$\rho(770) \eta$	seen		520
$\pi \pi \rho$	seen		633
$a_2(1320) \pi$	seen		307
$\rho \rho$	seen		335

 $\rho(1700)$

$$I^G(J^{PC}) = 1^+(1^--)$$

See the note in $\rho(1700)$ Particle Listings.Mass $m = 1720 \pm 20$ MeV [1] ($\eta \rho^0$ and $\pi^+ \pi^-$ modes)Full width $\Gamma = 250 \pm 100$ MeV [1] ($\eta \rho^0$ and $\pi^+ \pi^-$ modes)

$\rho(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$2(\pi^+\pi^-)$	seen	803
$\rho\pi\pi$	seen	653
$\rho^0\pi^+\pi^-$	seen	651
$\rho^\pm\pi^\mp\pi^0$	seen	652
$a_1(1260)\pi$	seen	404
$h_1(1170)\pi$	seen	450
$\pi(1300)\pi$	seen	349
$\rho\rho$	seen	372
$\pi^+\pi^-$	seen	849
$\pi\pi$	seen	849
$K\bar{K}^*(892) + \text{c.c.}$	seen	496
$\eta\rho$	seen	545
$a_2(1320)\pi$	not seen	334
$K\bar{K}$	seen	704
e^+e^-	seen	860
$\pi^0\omega$	seen	674
$\pi^0\gamma$	not seen	855

 $a_2(1700)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

Mass $m = 1698 \pm 40$ MeVFull width $\Gamma = 265 \pm 60$ MeV

$a_2(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta\pi$	$(3.6 \pm 1.1) \%$	754
$\gamma\gamma$	$(1.13 \pm 0.30) \times 10^{-6}$	849
$\rho\pi$	seen	664
$f_2(1270)\pi$	seen	350
$K\bar{K}$	$(1.9 \pm 1.2) \%$	691
$\omega\pi^-\pi^0$	seen	634
$\omega\rho$	seen	338

 $f_0(1710)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Non- $q\bar{q}$ Mesons."Mass $m = 1704 \pm 12$ MeVFull width $\Gamma = 123 \pm 18$ MeV

$f_0(1710)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}$	seen	694
$\eta\eta$	seen	652
$\pi\pi$	seen	841
$\gamma\gamma$	seen	852
$\omega\omega$	seen	337

 $\pi(1800)$

$$I^G(J^{PC}) = 1^-(0^--)$$

$$\text{Mass } m = 1810^{+9}_{-11} \text{ MeV} \quad (S = 2.2)$$

$$\text{Full width } \Gamma = 215^{+7}_{-8} \text{ MeV}$$

$\pi(1800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\pi^+\pi^-\pi^-$	seen	878
$f_0(500)\pi^-$	seen	—
$f_0(980)\pi^-$	seen	624
$f_0(1370)\pi^-$	seen	366
$f_0(1500)\pi^-$	not seen	247
$\rho\pi^-$	not seen	731
$\eta\eta\pi^-$	seen	660
$a_0(980)\eta$	seen	471
$a_2(1320)\eta$	not seen	†
$f_2(1270)\pi$	not seen	441
$f_0(1370)\pi^-$	not seen	366
$f_0(1500)\pi^-$	seen	247
$\eta\eta'(958)\pi^-$	seen	373
$K_0^*(1430)K^-$	seen	†
$K^*(892)K^-$	not seen	568

 $\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

$$\text{Mass } m = 1854 \pm 7 \text{ MeV}$$

$$\text{Full width } \Gamma = 87^{+28}_{-23} \text{ MeV} \quad (S = 1.2)$$

$\phi_3(1850)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}$	seen	785
$K\bar{K}^*(892) + \text{c.c.}$	seen	602

$\eta_2(1870)$

$$I^G(J^{PC}) = 0^+(2^-+)$$

Mass $m = 1842 \pm 8$ MeVFull width $\Gamma = 225 \pm 14$ MeV **$\eta_2(1870)$ DECAY MODES**

	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma\gamma$	seen	921

 $\pi_2(1880)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

Mass $m = 1874^{+26}_{-5}$ MeV ($S = 1.6$)Full width $\Gamma = 237^{+33}_{-30}$ MeV ($S = 1.2$) **$f_2(1950)$**

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 1936 \pm 12$ MeV ($S = 1.3$)Full width $\Gamma = 464 \pm 24$ MeV **$f_2(1950)$ DECAY MODES**

	Fraction (Γ_i/Γ)	p (MeV/c)
$K^*(892)\bar{K}^*(892)$	seen	377
$\pi^+\pi^-$	seen	958
$\pi^0\pi^0$	seen	959
4π	seen	921
$\eta\eta$	seen	798
$K\bar{K}$	seen	833
$\gamma\gamma$	seen	968
$p\bar{p}$	seen	238

 $a_4(1970)$

$$I^G(J^{PC}) = 1^-(4^{++})$$

was $a_4(2040)$ Mass $m = 1967 \pm 16$ MeV ($S = 2.1$)Full width $\Gamma = 324^{+15}_{-18}$ MeV **$a_4(1970)$ DECAY MODES**

	Fraction (Γ_i/Γ)	p (MeV/c)
$K\bar{K}$	seen	851
$\pi^+\pi^-\pi^0$	seen	959
$\rho\pi$	seen	825
$f_2(1270)\pi$	seen	559

$\omega \pi^- \pi^0$	seen	801
$\omega \rho$	seen	601
$\eta \pi$	seen	902
$\eta'(958) \pi$	seen	743

 $f_2(2010)$

$$J^{PC} = 0^+(2^{++})$$

Mass $m = 2011^{+60}_{-80}$ MeVFull width $\Gamma = 202 \pm 60$ MeV

$f_2(2010)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi \phi$	seen	†
$K \bar{K}$	seen	876

 $f_4(2050)$

$$J^{PC} = 0^+(4^{++})$$

Mass $m = 2018 \pm 11$ MeV (S = 2.1)Full width $\Gamma = 237 \pm 18$ MeV (S = 1.9)

$f_4(2050)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\omega \omega$	seen	637
$\pi \pi$	$(17.0 \pm 1.5) \%$	1000
$K \bar{K}$	$(6.8^{+3.4}_{-1.8}) \times 10^{-3}$	880
$\eta \eta$	$(2.1 \pm 0.8) \times 10^{-3}$	848
$4\pi^0$	$< 1.2 \%$	964
$a_2(1320) \pi$	seen	567

 $\phi(2170)$

$$J^{PC} = 0^-(1^{--})$$

Mass $m = 2159 \pm 17$ MeV [J] (S = 1.4)Full width $\Gamma = 137 \pm 16$ MeV [J]

$\phi(2170)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$e^+ e^-$	seen	1080
$\phi f_0(980)$	seen	395
$K^+ K^- f_0(980) \rightarrow$ $K^+ K^- \pi^+ \pi^-$	seen	—
$K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen	—
$K^{*0} K^\pm \pi^\mp$	not seen	759
$K^*(892)^0 \bar{K}^*(892)^0$	not seen	609

$f_2(2300)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2297 \pm 28$ MeVFull width $\Gamma = 149 \pm 40$ MeV

$f_2(2300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi\phi$	seen	529
$K\bar{K}$	seen	1037
$\gamma\gamma$	seen	1149

 $f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2345^{+50}_{-40}$ MeVFull width $\Gamma = 322^{+70}_{-60}$ MeV

$f_2(2340)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi\phi$	seen	580
$\eta\eta$	seen	1037

NOTES

- [a] See the review on “Form Factors for Radiative Pion and Kaon Decays” for definitions and details.
- [b] Measurements of $\Gamma(e^+\nu_e)/\Gamma(\mu^+\nu_\mu)$ always include decays with γ 's, and measurements of $\Gamma(e^+\nu_e\gamma)$ and $\Gamma(\mu^+\nu_\mu\gamma)$ never include low-energy γ 's. Therefore, since no clean separation is possible, we consider the modes with γ 's to be subreactions of the modes without them, and let $[\Gamma(e^+\nu_e) + \Gamma(\mu^+\nu_\mu)]/\Gamma_{\text{total}} = 100\%$.
- [c] See the $\pi^\pm$ Particle Listings for the energy limits used in this measurement; low-energy γ 's are not included.
- [d] Derived from an analysis of neutrino-oscillation experiments.
- [e] Astrophysical and cosmological arguments give limits of order 10^{-13} .
- [f] Forbidden by angular momentum conservation.
- [g] C parity forbids this to occur as a single-photon process.
- [h] The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+\mu^-) = \Gamma(\rho^0 \rightarrow e^+e^-) \times 0.99785$.
- [i] See the “Note on $a_1(1260)$ ” in the $a_1(1260)$ Particle Listings in PDG 06, Journal of Physics **G33** 1 (2006).

[j] Our estimate. See the Particle Listings for details.

[k] See also the $\omega(1650)$.

[l] See also the $\omega(1420)$.