

# K<sub>1</sub>(1270)


$$I(J^P) = \frac{1}{2}(1^+)$$

## K<sub>1</sub>(1270) MASS

VALUE (MeV)

DOCUMENT ID

**1253±7 OUR AVERAGE** Includes data from the 4 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.



### PRODUCED BY K<sup>-</sup>, BACKWARD SCATTERING, HYPERON EXCHANGE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

**1275±10**    700    GAVILLET    78    HBC    +    4.2 K<sup>-</sup> p → Ξ<sup>-</sup> (Kππ)<sup>+</sup>

### PRODUCED BY K BEAMS

VALUE (MeV) DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

**1270±10**    <sup>1</sup> DAUM    81C CNTR    -    63 K<sup>-</sup> p → K<sup>-</sup> 2π p

• • • We do not use the following data for averages, fits, limits, etc. • • •

- ~ 1276    <sup>2</sup> TORNQVIST    82B    RVUE
- ~ 1300    VERGEEST    79    HBC    -    4.2 K<sup>-</sup> p → ( $\bar{K}\pi\pi$ )<sup>-</sup> p
- 1289±25    <sup>3</sup> CARNEGIE    77    ASPK    ±    13 K<sup>±</sup> p → (Kππ)<sup>±</sup> p
- ~ 1300    BRANDENB...    76    ASPK    ±    13 K<sup>±</sup> p → (Kππ)<sup>±</sup> p

~ 1270	OTTER	76	HBC	-	10,14,16	$K^- p \rightarrow (\bar{K}\pi\pi)^- p$
1260	DAVIS	72	HBC	+	12	$K^+ p$
1234±12	FIRESTONE	72B	DBC	+	12	$K^+ d$

<sup>1</sup> Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

<sup>2</sup> From a unitarized quark-model calculation.

<sup>3</sup> From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

## PRODUCED BY BEAMS OTHER THAN K MESONS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	----------------

The data in this block is included in the average printed for a previous datablock.

<b>1248.1 ± 3.3 ± 1.4</b>		GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1289.81 ± 0.56 ± 1.66	894k	AAIJ	18A1	LHCB	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
1279 ± 10	25k	<sup>1</sup> ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
1294 ± 10	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
1300	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
1242 + <sup>9</sup> / <sub>-10</sub>		<sup>2</sup> ASTIER	69	HBC	$\bar{p} p$
1300	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

<sup>1</sup> Systematic errors not estimated.

<sup>2</sup> This was called the C meson.

## PRODUCED IN $\tau$ LEPTON DECAYS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	------------	----------------

The data in this block is included in the average printed for a previous datablock.


<b>1254 ± 33 ± 34</b>	7k	ASNER	00B	CLEO	$\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
-----------------------	----	-------	-----	------	---

## $K_1(1270)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
--------------------	--------------------

**90 ± 20 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.

**101 ± 12 OUR AVERAGE** Includes data from the 4 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.



**PRODUCED BY  $K^-$ , BACKWARD SCATTERING, HYPERON EXCHANGE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

<b>75 ± 15</b>	700	GAVILLET	78	HBC	+	4.2 $K^- p \rightarrow \Xi^- K \pi \pi$
----------------	-----	----------	----	-----	---	---

**PRODUCED BY  $K$  BEAMS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

<b>90 ± 8</b>	<sup>1</sup> DAUM	81C	CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
~ 150	VERGEEST	79	HBC	-	4.2 $K^- p \rightarrow (\bar{K} \pi \pi)^- p$
150 ± 71	<sup>2</sup> CARNEGIE	77	ASPK	±	13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$
~ 200	BRANDENB...	76	ASPK	±	13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$
120	DAVIS	72	HBC	+	12 $K^+ p$
188 ± 21	FIRESTONE	72B	DBC	+	12 $K^+ d$

<sup>1</sup> Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

<sup>2</sup> From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

**PRODUCED BY BEAMS OTHER THAN  $K$  MESONS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

<b>119.5 ± 5.2 ± 6.7</b>	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
--------------------------	-------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

116.11 ± 1.65 ± 2.96	894k	AAIJ	18A1	LHCB	$D^0 \rightarrow K^{\mp} \pi^{\pm} \pi^{\pm} \pi^{\mp}$
131 ± 21	25k	<sup>1</sup> ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
66 ± 15	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
127 <sup>+7</sup> / <sub>-25</sub>		ASTIER	69	HBC	$\bar{p} p$
60	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

<sup>1</sup> Systematic errors not estimated.

### PRODUCED IN $\tau$ LEPTON DECAYS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	------------	----------------

The data in this block is included in the average printed for a previous datablock.

<b>260<sup>+90</sup>/<sub>-70</sub> ± 80</b>	7k	ASNER	00B	CLEO	$\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
--	----	-------	-----	------	---

### $K_1(1270)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $K\rho$	(38 ± 13) %	2.2
$\Gamma_2$ $K_0^*(1430)\pi$	(28 ± 4) %	
$\Gamma_3$ $K^*(892)\pi$	(21 ± 10) %	2.2
$\Gamma_4$ $K\omega$	(11.0 ± 2.0) %	
$\Gamma_5$ $K f_0(1370)$	( 3.0 ± 2.0) %	
$\Gamma_6$ $\gamma K^0$	seen	

### $K_1(1270)$ PARTIAL WIDTHS

#### $\Gamma(K\rho)$ $\Gamma_1$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

57 ± 5	MAZZUCATO 79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
75 ± 6	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

#### $\Gamma(K_0^*(1430)\pi)$ $\Gamma_2$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

26 ± 6	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
--------	--------------	------	---	--

#### $\Gamma(K^*(892)\pi)$ $\Gamma_3$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 11	MAZZUCATO 79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
2 ± 2	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$\Gamma(K\omega)$   $\Gamma_4$ 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4 \pm 4$	MAZZUCATO 79	HBC	+	$4.2 K^- p \rightarrow \Xi^- (K\pi\pi)^+$
$24 \pm 3$	CARNEGIE 77B	ASPK	$\pm$	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$

 $\Gamma(K f_0(1370))$   $\Gamma_5$ 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$22 \pm 5$	CARNEGIE 77B	ASPK	$\pm$	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
------------	--------------	------	-------	--

 $\Gamma(\gamma K^0)$   $\Gamma_6$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

<b><math>73.2 \pm 6.1 \pm 28.3</math></b>	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$
---	-----------------	------	-----------------------------

 **$K_1(1270)$  BRANCHING RATIOS** $\Gamma(K\rho)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.38 \pm 0.13</math> OUR FIT</b>	Error includes scale factor of 2.2.		
---	-------------------------------------	--	--

<b><math>0.42 \pm 0.06</math></b>	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-----------------------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.584 \pm 0.043$	<sup>2</sup> GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
dominant	RODEBACK	81	HBC $4 \pi^- p \rightarrow \Lambda K 2\pi$

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.28 \pm 0.04</math></b>	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-----------------------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0201 \pm 0.0064$	<sup>2</sup> GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
---------------------	--------------------	----	---

 $\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.21 \pm 0.10</math> OUR FIT</b>	Error includes scale factor of 2.2.		
---	-------------------------------------	--	--

<b><math>0.16 \pm 0.05</math></b>	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-----------------------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.171 \pm 0.023$	<sup>2</sup> GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
-------------------	--------------------	----	---

 $\Gamma(K^*(892)\pi)/\Gamma(K\rho)$   $\Gamma_3/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.56 \pm 0.29</math> OUR FIT</b>	Error includes scale factor of 2.2.		
---	-------------------------------------	--	--

<b><math>0.99 \pm 0.15 \pm 0.18</math></b>	ABLIKIM	21U	BES3 $D_s^+ \rightarrow \bar{K}_1(1270)^0 K^+$
--	---------	-----	--

 $\Gamma(K\omega)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b><math>0.11 \pm 0.02</math></b>	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-----------------------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.225 \pm 0.052$	<sup>2</sup> GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
-------------------	--------------------	----	---

$\Gamma(K\omega)/\Gamma(K\rho)$  $\Gamma_4/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.30	95	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
-------	----	----------	----	-----	--

 $\Gamma(K f_0(1370))/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.03±0.02</b>	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
------------------	-------------------	-----	--

**D-wave/S-wave RATIO FOR  $K_1(1270) \rightarrow K^*(892)\pi$** 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>1.0±0.7</b>	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
----------------	-------------------	-----	--

<sup>1</sup> Average from low and high  $t$  data.<sup>2</sup> Assuming that decays are saturated by the  $K\rho$ ,  $K_0^*(1430)\pi$ ,  $K^*(892)\pi$ ,  $K\omega$  decay modes and neglecting interference between them. The values  $B(\omega \rightarrow \pi^+\pi^-) = (1.53^{+0.11}_{-0.13})\%$  and  $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10)\%$  are used. Systematic uncertainties not estimated. **$K_1(1270)$  REFERENCES**

ABLIKIM	21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18AI	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
GENG	07	PR D75 014017	L.S. Geng <i>et al.</i>	
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
RODEBACK	81	ZPHY C9 9	S. Rodeback <i>et al.</i>	(CERN, CDEF, MADR+)
MAZZUCATO	79	NP B156 532	M. Mazzucato <i>et al.</i>	(CERN, ZEEM, NIJM+)
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
GAVILLET	78	PL 76B 517	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+) JP
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
CARNEGIE	77B	PL 68B 287	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
OTTER	76	NP B106 77	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
CRENNELL	72	PR D6 1220	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)
ASTIER	69	NP B10 65	A. Astier <i>et al.</i>	(CDEF, CERN, IPNP, LVP) IJP
CRENNELL	67	PRL 19 44	D.J. Crennell <i>et al.</i>	(BNL) I